

Engagement in HIV care among young female sex workers in Zimbabwe

Sue Napierala, PhD, Women's Global Health Imperative, RTI International; San Francisco, CA,  
United States

Sungai Tafadzwa Chabata, MSc, Center for Sexual Health and HIV/AIDS Research  
(CeSHHAR); Harare, Zimbabwe

Elizabeth Fearon, PhD, London School of Hygiene and Tropical Medicine; London, United  
Kingdom

Calum Davey, PhD, London School of Hygiene and Tropical Medicine; London, United  
Kingdom

James Hargreaves, PhD, London School of Hygiene and Tropical Medicine; London, United  
Kingdom

Joanna Busza, MSc, London School of Hygiene and Tropical Medicine; London, United  
Kingdom

Phillis Mushati, MSc, Center for Sexual Health and HIV/AIDS Research (CeSHHAR); Harare,  
Zimbabwe

Sibongile Mtetwa, MSc, Center for Sexual Health and HIV/AIDS Research (CeSHHAR);  
Harare, Zimbabwe

Tarisai Chiyaka, MSc, Center for Sexual Health and HIV/AIDS Research (CeSHHAR); Harare, Zimbabwe

Owen Mugurungi, MD, Ministry of Health and Child Care, Harare, Zimbabwe

Dagmar Hanisch, MPH, United Nations Population Fund; Harare, Zimbabwe

Karin Hatzold, MD, Population Services International, Harare, Zimbabwe

Andrew Phillips, PhD, University College London; London, United Kingdom

Frances M Cowan, MD, Liverpool School of Tropical Medicine; Liverpool, United Kingdom;  
Center for Sexual Health and HIV/AIDS Research (CeSHHAR); Harare, Zimbabwe

Corresponding author: Sue Napierala

351 California Street

San Francisco, CA 94104

Phone: +1 (415) 848 1384

Fax: +1 (415) 848 1300

snapierala@rti.org

Parts of this manuscript have been previously presented in 2015 at the 8th IAS Conference on HIV pathogenesis, Treatment & Prevention, Vancouver, Canada.

Conflicts of Interest and Source of Funding: The SAPPH-IRe trial baseline survey was funded by United Nations Population Fund via Zimbabwe's Integrated Support Fund which receives funds from DfID, Irish Aid and Swedish SIDA. A small amount of funding for survey work was from GIZ. The authors declare that they have no competing interests.

Running head: Engagement in care among young sex workers

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and build upon the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

## Abstract

**Introduction:** Young female sex workers (FSW) are at greater HIV risk than their older counterparts. Yet the extent of their engagement with HIV services is largely unknown. We compared engagement among FSW aged 18-24 with those  $\geq 25$  years.

**Methods:** We used respondent-driven sampling to recruit FSW from 14 communities in Zimbabwe from November-December 2013. We collected data on demographics, behaviour, service uptake, and HIV and viral load testing. Data were pooled and weighted using RDS-2 estimation. We analyzed HIV care cascade variables by age group. To identify potential drivers of younger FSW service use, we explored factors associated with knowing one's HIV status.

**Results:** Among 2617 participants, mean age was 31 years and 26% were 18-24. Over half initiated sex work before age 25. Overall HIV prevalence was 59%, but was lower among younger FSW (35% vs 67%,  $p < 0.01$ ). Younger HIV-infected FSW were significantly less engaged at each step of the care cascade. Among younger FSW reporting ART use, 62% had an undetectable viral load compared with 79% in older FSW. In multivariable regression, young FSW encouraged to have an HIV test by another FSW (AOR=2.54; 95% confidence interval

[CI]:1.44-4.50), and those with no recent clients (AOR=4.31; 95%CI: 1.30-14.33) were more likely to report knowing their status.

Conclusions: The high proportion of FSW initiating sex work before age 25 and their lower engagement in HIV services highlights the importance of considering this vulnerable population in HIV programming. Implementing targeted services tailored to the unique needs of young FSW is a public health imperative.

Keywords: sex workers, young people, HIV care continuum, Zimbabwe, key and vulnerable populations

## **Introduction**

Young women who sell sex are highly vulnerable, and at considerably greater risk for HIV acquisition than their older counterparts.(1) A substantial proportion of female sex workers (FSW) begin selling sex as young women, with data indicating 20-40% of FSW globally initiated sex work as adolescents <20 years.(2) Young FSW experience both vulnerabilities of youth, including lack of knowledge, poorly developed life skills, lack of financial autonomy and limited access to health facilities,(3) as well as those associated with being FSW, such as stigma, discrimination, criminalization and violence.(4-6) Compared to older FSW, studies have shown that younger FSW have increased numbers of partners, less power to negotiate condom use, increased susceptibility to violence, as well as increased biological susceptibility to HIV acquisition.(7-9) There has been one study using programmatic data from Zimbabwe's National

Sex Worker Programme reporting that annual HIV incidence may be as high as 10% among FSW  $\leq 25$  years of age, as compared to 6% in those  $\geq 36$  years.(10)

A 2014 UNAIDS technical brief provides examples of successful programs, highlighting considerations for targeting services to young FSW.(1) However, few programs currently address the unique needs of young FSW, and empirical data on how best to deliver these evidence-based interventions in a way that is acceptable and accessible to this population remain sparse.(11, 12) Furthermore, particularly for young women, sex work is often a fluid concept, with women having different interpretations of, and identification with, sex work.(13, 14) During the period of sex work initiation, young women may be at especially high risk of new infection and least likely to access both general and FSW targeted services.(11, 12) The extent to which young FSW are engaged with HIV services is unknown.

In Zimbabwe sex work is illegal, however FSW have been identified as a key population and included in Zimbabwe's National HIV and AIDS Strategic Plan since 2006.(15-17) In 2009, following a national FSW situational analysis in Zimbabwe,(18) the National Sex Worker Programme, *Sisters with a Voice*, was established in five sites. By 2013 the *Sisters* program had expanded to 36 sites, covering all provinces of Zimbabwe. Services are based on guidance from the World Health Organization (WHO)(19) and include dedicated FSW clinics offering HIV testing and counselling, reproductive health services, condom provision and health education supported by trained peer educators, and a program of participatory activities to build community empowerment. As of December 2017, the *Sisters* program had seen approximately 65,000 women at around 155,000 visits. In 2013 we conducted surveys of FSW in 14 sites around Zimbabwe as part of the Sisters Antiretroviral therapy Programme for Prevention of HIV-an Integrated Response (SAPPH-IRe) trial (Pan African Trials Registry

[PACTR201312000722390]); a cluster randomized trial to determine the effectiveness of an enhanced community-based intervention to increase uptake, retention in care and adherence to antiretroviral-based prevention and treatment among FSW. In this analysis, we compared engagement in services and the HIV care cascade among FSW aged 18-24 years compared with those  $\geq 25$  years of age. We also explored factors associated with young FSWs' engagement in HIV services. We focused specifically on knowledge of HIV status, as this represents the biggest drop off in the cascade, particularly among younger women.

## Methods

Between November and December 2013, 2722 FSW were recruited using respondent-driven sampling (RDS) from 14 *Sisters* sites in Zimbabwe. RDS was used as it was not feasible to assemble a sampling frame of the intended target population. RDS is a recommended sampling strategy for hard-to-reach populations.(20) The results are reported using the STROBE-RDS reporting guidelines.(21) Detailed methods are described elsewhere (22), but in brief: eligible participants were  $\geq 18$  years and currently working as a FSW, defined as having exchanged sex for money, goods or services in the past 30 days, and had lived at the site for at least the previous 6 months. At each site we conducted geographic and social mapping in order to identify 6-8 'seed' participants. Seeds were purposefully selected to represent all sub-populations within the sex worker communities at each site. Seeds were interviewed and given two recruitment coupons to pass on to other sex workers in their social network. When women receiving a coupon attended for the interview they were then given two coupons to pass on to other FSW they knew who worked in that location, and had not previously been recruited to the survey. Participants were given \$5 to compensate for loss of earnings during the interview, and \$2 for each peer recruited. In all 14 sites a maximum of six iterations, or 'waves', of this process were

performed, including initial seeds. Approximately 200 FSW were recruited into the study per site. All participants completed an interviewer-administered questionnaire to collect data on demographics, sexual behaviour, sex work, HIV testing history and serostatus, uptake of HIV services and antiretroviral therapy (ART) use. All women provided a finger-prick dried blood spot sample for HIV testing. If this test was positive viral load (VL) was measured using the same sample.

### *Measures*

Younger FSW were defined as less than 25 years of age (WHO upper age range for defining youth and young people), and older FSW were defined as 25 years of age and older. Finger prick blood samples were collected as dried blood spot (DBS) and used for HIV antibody testing (AniLabsystems EIA kit; AniLabsystems Ltd, OyToilette 3, FIN-01720, Finland). If HIV antibodies were detected then the sample was tested for HIV viral load using NucliSENS EasyQ HIV-1 v2.0, to quantify the viral load. An undetectable viral load, i.e. viral suppression, was defined as a viral load of <1000 copies/mL. Knowing one's HIV status was defined as reporting having previously received an HIV positive test result, or having received an HIV-negative test result in the past 6 months prior to the survey. Mental health was assessed using the validated Patient Health Questionnaire (PHQ-9), a set of nine questions about mental state in the previous two weeks.<sup>(23)</sup> A score of nine or more out of the 20-item assessment indicated depressive disorder.

### *Statistical analyses*

Performance of the RDS surveys and RDS analytic methods have been reported elsewhere. We pooled data from across the 14 survey sites. We applied the svy command in Stata to replicate what the RDS-2 analysis package performs, probability weighting participants by the inverse of their network size, i.e. the number of women that each individual could have recruited, but adapted it for use with data pooled across sites. We normalized inverse degree weights by dividing them by the sum of inverse degrees at each site because the degree distribution differed across sites and we did not wish to weight women up or down on the basis of where they were from. As recommended, seeds were dropped in conducting the RDS estimation and regression analyses.(24) A fixed effect term for study site was included in regression analyses. All analyses were conducted using Stata 14.

Characteristics of participants, and variables of importance along the HIV care cascade, were analyzed descriptively, and stratified by younger versus older age. Differences in distribution of categorical variables by age group were tested using chi-squared tests. To further understand potential drivers of engagement in HIV services among younger FSW, we explored factors associated with knowing one's HIV status, as defined above. We included in our model demographic, behavioral and health factors which could conceivably influence engagement in services. We estimated the odds ratio (OR) and 95% CI of factors associated with knowing one's status using logistic regression models. Adjusted Wald tests were used to calculate p-values. Factors associated with the outcome at  $p \leq 0.20$  on a Wald test in univariable regression were entered into a multivariable regression model. A hierarchical model was used, first fitted on sociodemographic and then behavioral and health variables, which could be on the causal pathway between sociodemographic factors and the outcome. (25) Variables were retained in the



final model if independently associated with the outcome at the  $p \leq 0.10$  level, or if their inclusion altered the effect estimate of any variables in the model by  $\geq 10\%$  (i.e. were possible confounders).

### *Ethics*

This research was reviewed and approved by the Medical Research Council Zimbabwe, Research Council of Zimbabwe, University College London, the London School of Hygiene and Tropical Medicine, and RTI International prior to initiating research activities. All participants provided written informed consent for study participation.

### **Results**

A total of 2,722 participants were recruited over six waves from the 14 sites. The ninety seeds were dropped from the analysis. An additional 15 participants were missing recruiter information and were therefore treated as seeds and dropped from the analysis, leaving a total of 2617 participants.

Mean age of participants was 31 years (range 18-65), and 24% were under 25 years. Table 1 shows RDS-weighted baseline characteristics overall and stratified by younger versus older age. Overall 1450 (42%) women reported initiating sex work before age 25, with 11% starting before 18. Younger FSW had a mean duration of sex work of 2.6 years, versus 8.0 years for older FSW (data not shown). Younger FSW were more likely to have never been married ( $p < 0.01$ ) and less likely to have had children ( $p < 0.01$ ). They were more educated ( $p < 0.01$ ), more likely to report no religion ( $p < 0.01$ ), and to rate their overall health as good or very good ( $p < 0.01$ ).

HIV prevalence among the whole population was 59% and was lower among younger FSW (35% vs 67%,  $p < 0.01$ ; Table 2). Virtually everyone (99%) reported knowing where to get an HIV test. However, younger FSW were more likely to report having had an HIV test in the past 6 months (73% vs 52%,  $p < 0.01$ ) and this remained true when restricting to HIV-negative participants (77% vs 68%,  $p = 0.01$ ; data not shown). However, among FSW living with HIV, younger FSW were less likely to report knowing their HIV status (38% vs 69%,  $p < 0.01$ ). Among those who reported knowing their HIV-positive status, younger FSW were moderately less likely to report ART use (55% vs 69%,  $p = 0.06$ ). A high proportion of women on ART in both age groups reported 100% ART adherence (83% vs 88%). However, only 62% of younger FSW reporting ART use had a viral load  $< 1000$  copies/mL, compared with 79% in older FSW ( $p = 0.06$ ). In both groups a proportion of those who knew their HIV-positive status and reported no ART use had a viral load  $< 1000$  copies/mL, including 9% among younger FSW and 20% in older FSW ( $p = 0.09$ ).

Figure 1 shows the HIV care cascades among younger and older FSW, respectively. Each step on the cascade was statistically significantly different between the two age groups. Comparing the younger vs older age groups at each stop of the cascade, 38% vs 69% knew their positive status. Among these, 55% vs 68% were on ART, and of those 62% vs 79% were virally suppressed, respectively. Among all HIV-infected younger FSW, just 13% knew their status, were on ART, and had an undetectable viral load, compared with 37% of older FSW. Among all HIV-infected FSW, 1% of younger FSW and 4% of older FSW reported not being on ART but were virally suppressed.

Among young FSW, in univariable logistic regression analysis of factors associated with knowing their HIV status (Table 3), women reporting no clients in the previous week (odds ratio [OR]=3.87; 95% confidence interval [CI]: 1.17-12.85), as compared to 1-4 clients, were more likely to know their status. Young FSW who reported being encouraged to have an HIV test by another FSW in the past month were more likely to know their status (OR=2.41; 95%CI: 1.37-4.23). In multivariable analysis, these variables remained significantly associated with young FSW knowing their HIV status: having no clients in the last week (adjusted odds ratio [AOR] =4.31; 95%CI: 1.30-14.33), and those encouraged to have an HIV test by another FSW in the past month (AOR=2.54; 95%CI:1.44-4.50).

## **Discussion**

This study compares engagement with HIV services and progression along the care cascade between younger FSW <25 years and older FSW  $\geq$ 25 years. Few comparative data exist which estimate younger FSW engagement along the cascade. In our work, roughly a quarter of all FSW sampled were aged 18-24, and over half of all FSW reported initiating sex work before the age of 25. Accurate global estimates of the number of young people engaged in sex work are lacking,<sup>(1)</sup> however our findings support estimates that 20-40% of FSW globally initiated sex work as adolescents.<sup>(2)</sup> HIV prevalence was high (59%) overall, but as expected was lower among younger FSW, who are likely to have initiated sex work more recently. While recent testing was more common among younger FSW, HIV-positive young FSW were less likely to report knowing their HIV status. This may reflect high HIV incidence and more recent HIV infection, which has been documented in other research among FSW in Zimbabwe.<sup>(10)</sup> Fewer younger FSW reported taking ART, even amongst those who knew their status. This could again reflect more recent infection and/or slower HIV progression, or it could indicate lower

engagement in HIV care more generally. Differential social desirability bias, whereby younger women may be less comfortable disclosing their positive HIV status or initiation onto ART, may also contribute to these findings. Challenges in uptake of and adherence to HIV services for FSW have been well-documented.(26, 27) There are additional challenges facing younger women, including lack of knowledge, limited social/economic empowerment, and autonomy.(28-30) In addition, there may be competition and mistrust between older and younger FSW, which may result in younger women failing to access services designed for adults.(12, 31) These additional barriers may need to be specifically addressed in order to achieve more equitable engagement of young FSW in HIV services.

We conducted a regression analysis of factors associated with reporting knowing one's HIV status, in order to identify potential areas for intervention. The strongest predictor of young FSW knowing their HIV status was being encouraged to have an HIV test by another FSW. This finding lends support to the growing body of research on the positive impact of social support and community empowerment among FSW in HIV prevention and care.(4, 32) This is an important consideration for future HIV programming for this population, and may be particularly relevant for younger FSW who are the most difficult to access within the population of FSW as a whole.

Among young FSW who reported that they were positive, 74% were on ART and 62% of those on ART had a viral load <1000 copies/mL. At the time of this survey "test and treat" had not been introduced as national policy (although was the policy for women self-identifying as sex workers), therefore many newly HIV-infected young FSW may not have been identified as eligible for treatment. With introduction of universal "test and treat" in Zimbabwe we would expect the proportion on ART to increase, and this should be evaluated in future research.

However the low rate of viral suppression among those on ART has implications for resistance and resistance transmission, and is a priority to address among these young women.

This research represents a robust analysis among a large number of FSW across 14 diverse sites across Zimbabwe. We used RDS, a technique designed for sampling hard-to-reach populations. With that in mind, this research has limitations. We applied a lower age limit of 18 years. It is likely that those under 18 years are the most vulnerable and least engaged in HIV services among FSW, and they were not represented in this research. We used RDS because it was not possible to assemble a sampling frame of the target population. RDS uses a model of sampling probability to weight observations to approximate a random sample. This model makes a number of assumptions that might not have been met, some of which cannot be investigated.(33)

However, we have investigated potential RDS biases relating to estimation of HIV care cascade indicators in this study previously, and judged that the process had worked well.(22) Because participants recruit each other, it is difficult to document refusal rates, which may bias the sample. . Knowledge of HIV status and ART use were self-reported, and thus subject to reporting bias, and this bias may also be differential by age. However, HIV status and viral load at the time of the survey were biologically measured. The use of dried blood spot samples may have decreased sensitivity as compared to plasma, but our previously published validation study demonstrated good agreement.(34)

There is little research on young FSW and their engagement in HIV services, and there are currently few interventions targeting their specific needs, and even fewer that have been taken to scale.(1) The transition into sex work is likely an especially important time for young women, when self-identification as a FSW is low, risk is very high, negotiation skills are generally lower, and capacity to access targeted services weak. The fact that over half of women in our study

reported initiating sex work before 25 years of age highlights the importance of considering the needs of this vulnerable population in HIV programming. Despite the availability of a national network of dedicated FSW clinics in Zimbabwe, similar reported attendance at the clinics and similar sociodemographic characteristics between younger and older FSW, our research demonstrated that engagement at each step along the HIV care cascade was significantly lower among younger FSW as compared to their older counterparts. A number of challenges have been identified in supporting HIV prevention and care for younger FSW, particularly so for those under aged 18 who are not represented in this study.(35) These include difficulty in identifying young women who sell sex, particularly as initiation into sex work may be gradual and they may not self-identify as FSW. For those under 18, there may be a tension between harm reduction and child protection approaches, whereby young women may not want to present for services for fear of laws around the criminalization of sex work or child trafficking. Stigma around sex work and HIV remain significant barriers to engagement in services. (36) Young FSW are often highly mobile, making service provision challenging.(28) These represent just some of the challenges of delivering HIV programmes for young FSW.

## **Conclusions**

We have a growing body of research demonstrating possible strategies to increase engagement of young FSW in HIV services. An appropriate package of services will likely include combination strategies including biological and structural interventions which are tailored to the priorities and needs of adolescent girls and young women. A critical component of these programmes is social cohesion, bringing girls together as a community and empowering them. This needs to support other activities. Harm reduction and condom access are vital. Formative research in Zimbabwe suggests that education, training and skills-building opportunities are necessary, both to support

less risky behaviour and increased economic empowerment, and/or to provide alternatives to sex work for those who desire them. Young FSW may operate in different situations compared to older FSW, and therefore need to be supported on a case-by-case basis according to their individual needs.

Within the *Sisters With a Voice* program in Zimbabwe, we have already begun implementing additional services targeting younger FSW, and young peer educators have been recruited at all sites. In 2016 the Determined, Resilient, Empowered, AIDS-free, Mentored, Safe (DREAMS) intervention was also launched in Zimbabwe and includes working with adolescent girls and young women who sell sex in six of 63 districts. DREAMS is a multi-component intervention utilizing structural, behavioural, and biomedical approaches to reduce HIV amongst the highest risk adolescent girls and young women. These services have since been expanded with funding from Global Fund for AIDS TB and Malaria. Despite the myriad challenges, implementing targeted services, tailored to address the unique and varying needs of young women who sell sex is a public health imperative.

## References

1. UNAIDS, WHO. HIV and Young People Who Sell Sex: A technical brief. Geneva, Switzerland; 2015.
2. Silverman JG. Adolescent female sex workers: invisibility, violence and HIV. *Archives of disease in childhood*. 2011;96(5):478-81.
3. Napierala Mavedzenge SM, Doyle AM, Ross DA. HIV prevention in young people in sub-Saharan Africa: a systematic review. *J Adolesc Health*. 2011;49(6):568-86.

4. Shannon K, Strathdee SA, Goldenberg SM, Duff P, Mwangi P, Rusakova M, et al. Global epidemiology of HIV among female sex workers: influence of structural determinants. *Lancet*. 2015;385(9962):55-71.
5. Decker MR, Crago AL, Chu SK, Sherman SG, Seshu MS, Buthelezi K, et al. Human rights violations against sex workers: burden and effect on HIV. *Lancet*. 2015;385(9963):186-99.
6. Das P, Horton R. Bringing sex workers to the centre of the HIV response. *Lancet*. 2015;385(9962):3-4.
7. McClure C, Chandler C, Bissell S. Responses to HIV in sexually exploited children or adolescents who sell sex. *Lancet*. 2015;385(9963):97-9.
8. Dellar RC, Dlamini S, Karim QA. Adolescent girls and young women: key populations for HIV epidemic control. *J Int AIDS Soc*. 2015;18(2 Suppl 1):19408.
9. Bekker LG, Johnson L, Wallace M, Hosek S. Building our youth for the future. *J Int AIDS Soc*. 2015;18(2 Suppl 1):20027.
10. Hargreaves JR, Mtetwa S, Davey C, Dirawo J, Chidiya S, Benedikt C, et al. Cohort analysis of programme data to estimate HIV incidence and uptake of HIV-related services among female sex workers in Zimbabwe, 2009-14. *J Acquir Immune Defic Syndr*. 2015.
11. Busza J, Mtetwa S, Mapfumo R, Hanisch D, Wong-Gruenwald R, Cowan F. Underage and underserved: reaching young women who sell sex in Zimbabwe. *AIDS Care*. 2016;28 Suppl 2:14-20.
12. Delany-Moretlwe S, Cowan FM, Busza J, Bolton-Moore C, Kelley K, Fairlie L. Providing comprehensive health services for young key populations: needs, barriers and gaps. *J Int AIDS Soc*. 2015;18(2 Suppl 1):19833.



13. Stoebenau K, Heise L, Wamoyi J, Bobrova N. Revisiting the understanding of "transactional sex" in sub-Saharan Africa: A review and synthesis of the literature. *Soc Sci Med.* 2016;168:186-97.
14. Busza J. For love or money: the role of sexual exchange in young people's sexual relationships. In: Ingham R, Aggleton P, editors. *Promoting Young People's Sexual Health.* London and New York: Routledge; 2006. p. 134-53.
15. National AIDS Council, Ministry of Health and Child Care. *Zimbabwe National HIV and AIDS Strategic Plan (ZNASP) 2006-2010.*; 2006.
16. National AIDS Council, Ministry of Health and Child Care. *Zimbabwe National HIV and AIDS Strategic Plan (ZNASP) 2011-2015.*; 2011.
17. National AIDS Council, Ministry of Health and Child Care. *Zimbabwe National HIV and AIDS Strategic Plan (ZNASP) 2015-2018: Commitment towards fast tracking 90.90.90 targets by 2020 and ending aids by 2030.*; 2015.
18. Corbett EL. Unpublished data from 'Readiness assessment and pilot testing of supervised self-testing for HIV amongst communities and hospital patients and staff in urban Blantyre, Malawi'. 2010.
19. WHO U, UNAIDS, nswp. *Prevention and Treatment of HIV and other Sexually Transmitted Infections for Sex Workers in Low- and Middle-income Countries: Recommendations for a public health approach.* Geneva; 2012.
20. Heckathorn DD. Respondent-driven sampling: a new approach to the study of hidden populations. *Social problems.* 1997;44(2):174-99.

21. White RG, Hakim AJ, Salganik MJ, Spiller MW, Johnston LG, Kerr L, et al. Strengthening the Reporting of Observational Studies in Epidemiology for respondent-driven sampling studies: "STROBE-RDS" statement. *J Clin Epidemiol.* 2015;68(12):1463-71.
22. Cowan FM, Davey CB, Fearon E, Mushati P, Dirawo J, Cambiano V, et al. The HIV Care Cascade Among Female Sex Workers in Zimbabwe: Results of a Population-Based Survey From the Sisters Antiretroviral Therapy Programme for Prevention of HIV, an Integrated Response (SAPPH-IRe) Trial. *J Acquir Immune Defic Syndr.* 2017;74(4):375-82.
23. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606-13.
24. Volz E, Heckathorn DD. Probability based estimation theory for respondent driven sampling. *Journal of official statistics.* 2008;24(1):79.
25. Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol.* 1997;26(1):224-7.
26. Schwartz S, Lambert A, Phaswana-Mafuya N, Kose Z, McIngana M, Holland C, et al. Engagement in the HIV care cascade and barriers to antiretroviral therapy uptake among female sex workers in Port Elizabeth, South Africa: findings from a respondent-driven sampling study. *Sex Transm Infect.* 2017;93(4):290-6.
27. Ma PHX, Chan ZCY, Loke AY. The Socio-Ecological Model Approach to Understanding Barriers and Facilitators to the Accessing of Health Services by Sex Workers: A Systematic Review. *AIDS Behav.* 2017;21(8):2412-38.
28. Busza J, Mtetwa S, Chirawu P, Cowan F. Triple jeopardy: adolescent experiences of sex work and migration in Zimbabwe. *Health Place.* 2014;28:85-91.

29. Zhang XD, Temmerman M, Li Y, Luo W, Luchters S. Vulnerabilities, health needs and predictors of high-risk sexual behaviour among female adolescent sex workers in Kunming, China. *Sex Transm Infect.* 2013;89(3):237-44.
30. Inguane C, Horth RZ, Miranda AE, Young PW, Sathane I, Cummings BE, et al. Socio-demographic, Behavioral and Health Characteristics of Underage Female Sex Workers in Mozambique: The Need to Protect a Generation from HIV Risk. *AIDS Behav.* 2015;19(12):2184-93.
31. Beyrer C, Baral S, Kerrigan D, El-Bassel N, Bekker LG, Celentano DD. Expanding the space: inclusion of most-at-risk populations in HIV prevention, treatment, and care services. *J Acquir Immune Defic Syndr.* 2011;57 Suppl 2:S96-9.
32. Kerrigan D, Kennedy CE, Morgan-Thomas R, Reza-Paul S, Mwangi P, Win KT, et al. A community empowerment approach to the HIV response among sex workers: effectiveness, challenges, and considerations for implementation and scale-up. *Lancet.* 2015;385(9963):172-85.
33. Gile KJ, Handcock MS. Respondent-Driven Sampling: An Assessment of Current Methodology. *Sociol Methodol.* 2010;40(1):285-327.
34. Napierala Mavedzenge S, Davey C, Chirenje T, Mushati P, Mtetwa S, Dirawo J, et al. Finger Prick Dried Blood Spots for HIV Viral Load Measurement in Field Conditions in Zimbabwe. *PLoS One.* 2015;10(5):e0126878.
35. Steen R, Jana S, Reza-Paul S, Richter M. Trafficking, sex work, and HIV: efforts to resolve conflicts. *The Lancet.* 385(9963):94-6.
36. Hargreaves J, Busza J, Mushati P, Fearon E, Cowan F. Overlapping HIV and sex-work Stigma among female sex-workers recruited to 14 respondent-driven sampling surveys across Zimbabwe, 2013. *AIDS Care.* 2017;In press.

**Figure 1a-b.** Cascade of care for HIV-positive female sex workers a) <25 years, and b)  $\geq$ 25 years in Zimbabwe

ACCEPTED

**Table 1:** Baseline characteristics of study participants, weighted (N=2617)

Characteristic	Age at survey		Total n(%)	Comparison p-value
	18-24 years (N=641) n(%)	≥25 years (N=1976) n(%)		
Age at start of sex work				<0.001
	<18 years	159 (24.4)	139 (6.3)	298 (11.1)
	18-24 years	482 (75.6)	670 (30.1)	1,152 (42.1)
	25-29 years	n/a	614 (34.9)	614 (25.7)
	30-39 years	n/a	479 (25.2)	479 (18.6)
	≥40 years	n/a	74 (3.5)	74 (2.6)
Number of years in sex work				<0.001
	0-1 years	189 (36.3)	151 (9.8)	340 (16.8)
	2-4 years	368 (52.1)	587 (31.1)	955 (36.6)
	5-8 years	77 (10.5)	538 (25.2)	615 (21.3)
	≥9 years	7 (1.1)	700 (34.0)	707 (25.3)
Marital status				<0.001
	Divorced/separated	378 (59.9)	1,259 (62.5)	1,637 (61.9)
	Widowed	19 (2.4)	461 (25.6)	480 (19.5)
	Never been married	241 (37.3)	238 (10.8)	479 (17.8)
	Married/living together as if married	3 (0.4)	18 (1.0)	21 (0.9)
Number of children				<0.001
	0	191 (32.8)	294 (17.1)	485 (21.2)
	1	374 (57.6)	1,043 (51.6)	1,417 (53.2)
	≥2	76 (9.6)	639 (31.4)	715 (25.6)
Highest level of education				<0.001
	No formal schooling	9 (1.8)	87 (5.1)	96 (4.3)
	Some primary school	140 (23.5)	596 (35.5)	736 (32.3)
	Some secondary school	275 (45.4)	653 (31.5)	928 (35.2)
	Completed secondary or more	215 (29.3)	625 (27.9)	840 (28.3)
Religion				0.002
	Christian	318 (50.8)	1,161 (58.7)	1,479 (56.6)
	Other	47 (8.3)	223 (11.6)	270 (10.7)
	No religion	275 (40.9)	590 (29.7)	865 (32.7)
Proportion of income generated through sex work				0.01
	<25%	35 (6.1)	177 (9.5)	212 (8.6)
	25%-50%	56 (10.2)	243 (14.0)	299 (13.0)
	>50%-99%	157 (25.6)	516 (25.3)	673 (25.4)
	100%	393 (58.1)	1,040 (51.2)	1,433 (53.0)
Venue for client recruitment				0.32
	Bars/nightclubs/entertainment venue	471 (71.3)	1,366 (68.6)	1,837 (69.3)
	By telephone	35 (7.3)	121 (6.3)	156 (6.6)
	In the market place/street	98 (16.2)	347 (19.5)	445 (18.6)
	Other	26 (5.3)	96 (5.6)	122 (5.5)

Number of clients in the last week					0.02
	0	36 (6.1)	160 (9.6)	196 (8.7)	
	1-4	228 (41.6)	798 (43.8)	1,026 (43.3)	
	5-9	161 (25.2)	481 (24.2)	642 (24.4)	
	≥10	216 (27.1)	537 (22.4)	753 (23.7)	
Amount charged per client					0.04
	≤\$2	5 (1.4)	16 (1.1)	21 (1.2)	
	\$2-5	380 (56.4)	1,141 (61.4)	1,521 (60.1)	
	>\$5-10	207 (34.2)	693 (33.6)	900 (33.7)	
	>\$10	38 (8.1)	78 (3.9)	116 (5.0)	
Alcohol consumption in the past 12 months					0.02
	Never	222 (38.2)	780 (43.7)	1,002 (42.3)	
	Once a month or less	63 (10.5)	200 (11.1)	263 (10.9)	
	2-4 times per month	88 (13.0)	304 (15.2)	392 (14.6)	
	2-3 times per week	100 (14.7)	238 (10.7)	338 (11.7)	
	4 or more times per week	168 (23.6)	451 (19.4)	619 (20.5)	
Symptoms of depressive disorder*					0.17
	Yes	253 (41.2)	878 (45.8)	1131 (44.6)	
	No	387 (58.8)	1084 (54.2)	1471 (55.4)	
Personal health rating					<0.001
	Very good	115 (20.3)	188 (10.3)	303 (12.9)	
	Good	300 (50.5)	778 (40.0)	1,078 (42.8)	
	Fair	182 (22.9)	725 (35.2)	907 (31.9)	
	Poor	41 (6.3)	272 (14.6)	313 (12.4)	

\*As per the Patient Health Questionnaire (PHQ-9)

**Table 2:** Variables of importance along the HIV prevention and care cascades, by age group

Characteristic	Age at survey		Total n(%)	Comparison p-value
	18-24 years n(%)	≥25 years n(%)		
<i>Testing and testing history</i>				
Knows where to get an HIV test				0.23
	Yes	634 (99.0)	1,970 (99.5)	2,604 (99.4)
	No	6 (1.0)	6 (0.5)	12 (0.6)
Months since last HIV test				<0.001
	< 6 months	459 (72.7)	1,009 (51.5)	1,468 (57.2)
	6-12 months	50 (7.2)	210 (11.6)	260 (10.4)
	>12 months	63 (10.2)	576 (28.3)	639 (23.5)
	Never tested	58 (9.9)	147 (8.6)	205 (9.0)
Result of most recent HIV test, among those who have tested				<0.001
	Positive	106 (16.0)	960 (52.0)	1,066 (42.5)
	Negative	475 (84.0)	855 (48.0)	1,330 (57.5)
Ever told anyone the results of HIV test(s), among those who have tested				0.49
	Yes	458 (81.7)	1,487 (79.9)	1,945 (80.4)
	No	124 (18.3)	341 (20.1)	465 (19.6)
<i>HIV status</i>				
HIV status on day of survey				<0.001
	Positive	229 (34.7)	1,311 (67.4)	1,540 (58.7)
	Negative	411 (65.3)	648 (32.6)	1,059 (41.3)
Among those HIV-positive, knows HIV status <sup>†</sup>				<0.001
	Yes	97 (37.9)	913 (68.5)	1,010 (63.6)
	No	132 (62.1)	398 (31.6)	530 (36.4)
<i>Care-seeking behavior among those who know they are HIV-positive</i>				
Currently on ART				0.06
	Yes	49 (55.4)	625 (68.9)	674 (67.7)
	No	47 (44.6)	288 (31.1)	335 (32.3)
If not on ART, why not				0.47
	Not eligible for ART/advised to wait	38 (74.8)	244 (83.3)	282 (82.2)
	Eligible but cannot access ART	0 (0.0)	5 (2.1)	5 (1.8)
	Does not want to take ART	4 (16.7)	11 (3.8)	15 (5.4)
	Eligibility for ART has not been evaluated	2 (8.5)	22 (10.8)	24 (10.6)
Among those on ART, where is ART being accessed				0.27
	Hospital	30 (63.4)	308 (51.3)	358 (52.2)
	Dedicated sex worker (Sisters) clinic	0 (0.0)	5 (1.1)	8 (1.0)
	Government clinic	18 (35.9)	289 (45.7)	307 (45.0)
	Antenatal clinic	1 (0.7)	19 (1.9)	20 (1.8)
	Private Doctor	0 (0.0)	2 (0.0)	2 (0.0)
<i>Viral suppression among those who know they are HIV-positive</i>				
Among those reporting being on ART, ART is taken at the exact time prescribed				0.84
	Less than half of the time	1 (0.1)	10 (0.9)	11 (0.9)

Half of the time or more, but not most of the time	1 (0.4)	13 (1.7)	14 (1.6)	
Most of the time	5 (16.3)	60 (9.2)	65 (9.7)	
All of the time	42 (83.2)	539 (88.1)	581 (87.8)	
Among those reporting being on ART, viral load is undetectable (<1000 copies/ml)				0.06
Yes	23 (62.1)	463 (78.7)	486 (77.5)	
No	25 (37.9)	150 (21.3)	175 (22.6)	
Among those reporting not being on ART, viral load is undetectable (<1000 copies/ml)				0.09
Yes	9 (8.7)	66 (19.7)	75 (18.3)	
No	38 (91.3)	219 (80.3)	257 (81.7)	

<sup>†</sup>Defined as having reported receiving a previous HIV+ result



**Table 3:** Factors associated with knowing HIV status - defined as having received a positive result or having had a negative HIV test within the last 6 months prior to the survey - among young FSW (N=641)

Characteristic	Number of women knowing status (n=492) n(%)	Crude OR (95% CI)	p-value	Adjusted OR (95% CI)	p-value
Age			0.46		
18-19 years	78 (17.5)	ref			
20-24 years	414 (82.5)	1.27 (0.68-2.37)			
Number of years in sex work			0.30		
0-1 years	144 (38.1)	ref			
2-4 years	281 (49.8)	0.64 (0.36-1.12)			
5-8 years	60 (10.6)	0.79 (0.35-1.82)			
≥9 years	7 (1.5)	-			
Marital status			1.00		
Divorced/separated	293 (59.8)	ref			
Widowed	16 (2.3)	1.00 (0.21-4.89)			
Never been married	181 (37.4)	1.04 (0.59-1.83)			
Married/living together as if married	2 (0.4)	1.37 (0.07-27.31)			
Highest level of education			0.64		
No formal schooling	7 (1.8)	ref			
Some primary school	99 (21.8)	1.02 (0.19-5.34)			
Some secondary school	216 (45.8)	1.46 (0.29-7.43)			
Completed secondary or more	168 (30.7)	1.56 (0.29-8.26)			
Religion			0.64		
Christian	240 (49.8)	ref			
Other	31 (8.0)	0.81 (0.34-1.91)			
No religion	220 (42.2)	1.21 (0.67-2.19)			
Proportion of income generated through sex work			0.36		
<25%	25 (5.3)	ref			
25%-50%	41 (9.8)	1.52 (0.39-5.89)			
>50%-99%	128 (28.2)	2.79 (0.70-11.11)			
100%	298 (56.7)	1.46 (0.46-4.71)			
Venue for client recruitment			0.59		
Bars/nightclubs/entertainment venue	359 (70.2)	ref			
By telephone	27 (7.6)	1.35 (0.47-3.89)			
In the market place/street	76 (16.2)	1.22 (0.54-2.73)			
Other	20 (6.0)	2.22 (0.67-7.37)			
Number of clients in the last week			0.07		0.04
1-4	167 (41.7)	ref		ref	
5-9	127 (23.3)	0.70 (0.35-1.39)		0.64 (0.31-1.32)	
≥10	167 (27.5)	1.11 (0.58-2.12)		1.07 (0.56-2.02)	
0	31 (7.5)	3.87 (1.17-12.85)		4.31 (1.30-14.33)	

Condom use with clients in the past month				0.36
	Never	66 (10.0)	ref	
	Rarely - about 25% of the time	12 (1.7)	2.41 (0.43-13.64)	
	Some of the time - about 50% of the time	34 (7.6)	2.75 (0.75-10.04)	
	Most of the time - about 75% of the time	56 (14.3)	1.30 (0.49-3.49)	
	Always - 100% of the time	278 (60.8)	1.94 (0.785-4.47)	
	Don't wish to answer	46 (5.7)	0.95 (0.29-3.15)	
Was encouraged to have an HIV test by another FSW in the past month				0.002
	No	125 (25.7)	ref	ref
	Yes	365 (74.3)	2.41 (1.37-4.23)	2.54 (1.44-4.50)
Alcohol consumption in the past 12 months				0.54
	Never	176 (38.8)	ref	
	Once a month or less	48 (10.1)	0.75 (0.28-1.98)	
	2-4 times per month	75 (14.6)	1.51 (0.60-3.78)	
	2-3 times per week	71 (13.5)	0.64 (0.27-1.49)	
	4 or more times per week	122 (23.1)	0.94 (0.48-1.84)	
Symptoms of depressive disorder <sup>†</sup>				0.87
	No	290 (58.8)	ref	
	Yes	201 (41.2)	0.95 (0.54-1.68)	
Personal health rating				0.29
	Very good	91 (20.4)	ref	
	Good	245 (51.8)	1.18 (0.53-2.66)	
	Fair	121 (20.6)	0.74 (0.32-1.71)	
	Poor	35 (7.2)	2.05 (0.56-7.52)	

<sup>†</sup>As per the Patient Health Questionnaire (PHQ-9)

**Figure 1a-b.** Cascade of care for HIV-positive female sex workers a) <25 years, and b) ≥25 years in Zimbabwe

