

Molecular characterisation and taxon assemblage typing of giardiasis in primary school children living close to the shoreline of Lake Albert, Uganda

Hajri Al-Shehri^{a,b}, E. James LaCourse^a, Otto Klimach^a, Narcis B. Kabatereine^c, J. Russell Stothard^{a,*}

^a Department of Parasitology, Liverpool School of Tropical Medicine, Liverpool L3 5QA, UK

^b Ministry of Health, Asir District, Kingdom of Saudi Arabia

^c Vector Control Division, Ministry of Health, Kampala, Uganda

ARTICLE INFO

Article history:

Received 17 May 2018

Received in revised form 9 September 2018

Accepted 10 September 2018

Keywords:

Giardia duodenalis

Real-time PCR

Assemblage B

β -giardin

Wasting

ABSTRACT

As part of an epidemiological survey for gastrointestinal parasites in school children across five primary schools on the shoreline of Lake Albert, the prevalence of giardiasis was 87.0% ($n = 254$) as determined by real-time PCR analysis of faecal samples with a genus-specific *Giardia* 18S rDNA probe. Faecal samples were further characterised with taxon assemblage-specific triose phosphate isomerase (TPI) Taqman® probes and by sequence characterisation of the β -giardin gene. While less sensitive than the 18S rDNA assay, general prevalence by TPI probes was 52.4%, with prevalence by taxon assemblage of 8.3% (assemblage A), 35.8% (assemblage B) and 8.3% co-infection (A & B assemblages). While assemblage B was dominant across the sample, proportions of assemblages A and B, and co-infections thereof, varied by school and by age of child; mixed infections were particularly common at Runga school (OR = 6.9 [95% CI; 2.5, 19.3]) and in children aged 6 and under (OR = 2.7 [95% CI; 1.0, 7.3]). Infection with assemblage B was associated with underweight children (OR = 2.0 [95% CI; 1.0, 3.9]). The presence of each assemblage was also confirmed by sequence analysis of the β -giardin gene finding sub-assemblage All and further genetic diversity within assemblage B. To better explore the local epidemiology of giardiasis and its impact on child health, additional sampling of school children with assemblage typing would be worthwhile.

© 2018 The Authors. Published by Elsevier Ltd on behalf of World Federation of Parasitologists. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The binucleate flagellated protozoan *Giardia duodenalis* (syn. *G. lamblia*, *G. intestinalis*) is a common gastrointestinal parasite able to infect a variety of mammals (Adam, 2001; Helmy et al., 2014). Where sanitation and hygiene are poor, these parasites can cause acute and/or chronic giardiasis across all ages (Wegayehu et al., 2016; Muhsen and Levine, 2012; Rogawski et al., 2017; Tellevik et al., 2015). While levels of endemicity of giardiasis may vary across the world, it can be common in children living within low and middle income countries (Laishram et al., 2012; Muhsen and Levine, 2012); for example, in Uganda giardiasis can be particularly rife (Al-Shehri et al., 2016; Fuhrmann et al., 2016), but its effect on child health is not fully appreciated but in Rwanda nearby, the very high prevalence of *G. duodenalis* in children aged 5 and under, was associated with being underweight (Ignatius et al., 2012).

* Corresponding author.

E-mail address: russell.stothard@lstmed.ac.uk. (J.R. Stothard).

There are eight distinct groups or taxonomic assemblages (A to H) within *Giardia* currently recognised (Sprong et al., 2009; Almeida et al., 2010; Takumi et al., 2012). Assemblages A and B are typically held most responsible for human infections, with the latter assemblage associated with zoonotic transmission (Almeida et al., 2010; Feng and Xiao, 2011; Vanni et al., 2012; Asher et al., 2014; Thompson and Ash, 2016); each assemblage can be further divided into sub-assemblages, e.g. A: AI, AII & AIII and B: BIII and BIV on the basis of sequence variation within molecular markers e.g. glutamate dehydrogenase (GDH), β -giardin, small subunit ribosomal DNA (18S rDNA), and triose phosphate isomerase (TPI) (Durigan et al., 2014; Karim et al., 2015; Minetti et al., 2015). Despite efforts to investigate specific assemblages with disease symptoms and severity, there is no absolute association to date (Sprong et al., 2009; Thompson and Ash, 2016).

In Uganda, general investigations on the epidemiology of giardiasis are increasing (Nizeyi et al., 1999; Graczyk et al., 2002; Nizeyi et al., 2002; Johnston et al., 2010), although only a single study has employed molecular methods of characterisation (Ankarklev et al., 2012). Ankarklev et al. (2012) investigated associations between taxon assemblages and *Helicobacter pylori* infection in apparently healthy children aged 0–12 living in Kampala, the capital. Assemblage B was found dominant and a risk factor for *H. pylori* infection (Ankarklev et al., 2012) and like in other parts of the world, assemblage B was more associated with symptomatic infections (Pelayo et al., 2008; Puebla et al., 2014).

To shed light on the taxonomic assemblages of *Giardia* within school children living on the shoreline of Lake Albert, we undertook a molecular characterisation of previously characterised stool samples as reported by Al-Shehri et al. (Al-Shehri et al., 2016). Faecal samples were further characterised with assemblage-specific TaqMan® TPI probes and the presence of each taxon assemblage confirmed by sequence analysis of the β -giardin gene. Associations between taxon assemblage and collected epidemiological data were explored.

2. Materials and methods

2.1. Faecal material and epidemiological information

Faecal samples were available for further molecular analysis (see below) that were initially collected within the epidemiological survey of 254 school children from five primary schools (Bugoigo, Runga, Walakuba, Biiso and Busingiro) as reported by Al-Shehri et al. (Al-Shehri et al., 2016). Each sampled child underwent an epidemiological questionnaire and clinical examination; data on socio-demographical aspects and standard biometry were recorded (height with a clinical stadiometer, model 214; SECA, Hanover, MD and weight by weighing scales with a model 803; SECA, Hanover, MD). Heights and weights were used to assess stunting, height-for-age Z-score (HAZ), and wasting, weight-for-age Z-score (WAZ). Children were defined as stunted if their height-for-age Z score was $-2 \leq SD$ and underweight if their weight-for-age Z score was $-2 \leq SD$ (WHO, 2007). Finger-prick blood was collected from each child and tested for haemoglobin levels by HemoCue® portable haemoglobin photometer (HemoCue, CA 92630, USA). Children were considered anaemic if haemoglobin levels were below 115 g/L (WHO, 2011).

During the surveys, all sampled stools were tested for faecal occult blood (Mission Test, Acon Laboratories, San Diego, CA, USA) but owing to a limited supply of rapid diagnostic tests (RDTs), only stools collected from Bugoigo and Runga were tested in-field with Quik-Chek RDTs for giardiasis and cryptosporidiosis (GIADIA/CRYPTOSPORIDIUM Quik-Chek, Alere, Galway, Ireland). Stools were then stored in absolute ethanol for later DNA analysis.

2.2. Molecular profiling of *G. duodenalis* assemblages

After transfer to the UK and each faecal sample was spiked with Phocine Herpes Virus to act as an internal control for genomic DNA extraction and amplification performance of later real-time PCR assays. Genomic DNA was extracted, and detection of *Giardia* 18S rDNA was performed using TaqMan® assay following primers, probes and protocols of Verweij et al. (Verweij et al., 2004). These extractions were again retested with a duplex real-time PCR assay with assemblage-specific A and B probes using the TPI locus (Elwin et al., 2014). The real-time PCR analysis of faecal extractions from each school was completed in separate PCR plates that each contained negative and positive controls; a negative control (without genomic DNA template) of extraction elution buffer (10 mM Tris-HCl [pH 8], 1 mM EDTA) and a positive control (with reference genomic *Giardia* DNA template) from a heavily infected individual excreting approximately 1000 cysts per gram of faeces as estimated by microscopy. As a further quality control, reamplification of 10% of samples was undertaken to assess assemblage assay reliability. Assays were performed in a Chromo-4 with Opticon monitor™ version 3.1. (Bio-Rad, UK). The infection was determined according to C_t values; for the 18S rDNA TaqMan® assay no-infection was $C_t \geq 40$ and positive infection $C_t \leq 39$ while for assemblages-specific probes was $C_t \leq 45$.

To further confirm assemblage A and B, the β -giardin gene was amplified from samples from six children using nested PCR following protocols of Minetti et al. (Minetti et al., 2015). PCR products were purified using the QIAquick® PCR purification kit (QIAGEN Ltd.) and were sequenced in both directions by Sanger sequencing. Nucleotide sequences and chromatograms were analysed and edited using Geneious software (Vejlsovej55, 8600 Silkeborg, Denmark). Sequences from this study were aligned with each other and reference sequences downloaded from GenBank (listed below). The assemblages and sub-assemblages at each locus were identified by BLAST searches against the following reference sequences: β -giardin (accession nos. X14185.1–AI, AY072723.1–AII, DQ650649.1–AIII, AY072726.1–BIII, AY072725.1–BIV).

2.3. Statistical analyses

Statistical analysis was performed using Minitab Ltd.® (Brandon Court, Unit E1-E2 Coventry CV3 2TE UK). Binary logistic regression tests were performed to compare data from each school and as well as risk variables as an independent indicator to assess any associations with specific assemblages.

3. Results

Out of the 254 samples examined, 221 tested positive (87.0%) by targeting *Giardia* 18S rDNA assay while 133 (52.3%) tested positive with TPI assemblage-specific probes. Across Bugoigo and Runga schools, the prevalence of giardiasis by Quik-Chek RDT was 41.6%. Of the 133-tested positive by TPI probes, 21 samples were positive for assemblage A (15.8%) only, 91 positives for assemblage B (68.4%) only and 21 positives for both assemblage A and B (15.8%), mixed assemblage infections.

Across these samples assemblage, A was less common than assemblage B, an approximate ratio of 1: 2.7, with assemblage B dominant. To ascertain if there was any amplification bias in assemblage detection, Fig. 1A shows a bivariate plot of Ct values for *Giardia* 18S rDNA TaqMan® probe and the corresponding Ct value of assemblage A TPI TaqMan® probe ($18S\ rDNA = 0.203 + 0.6991\ TPI$, with R-squared 34.91% ($P < 0.005$), and positive correlation ($r = 0.60$)); Fig. 1B shows bivariate plot for assemblage B ($18S\ rDNA = 0.228 + 0.6947\ TPI$, with R-squared 28.39% ($P < 0.005$), and positive correlation ($r = 0.54$)). The performance of each TaqMan® assay appeared equivalent. Of note, however, is that mixed assemblage infections appear more common at Runga school where the local prevalence of assemblage A was also much higher.

Table 2 shows epidemiological associations cross-tabulated against available assemblage information. Most notable is the association of mixed assemblages in younger children (OR = 2.7 [95% CI; 1.0, 7.3]) that assemblage B was associated with the presence of faecal occult blood (OR = 2.2 [95% CI; 1.0, 5.2]). It appeared that there was also a significant association of infection with assemblage B and children of lower weight-for-age, i.e. wasting (OR = 2.0 [95% CI; 1.0, 3.9]).

Table 3 details the point mutations with the six representative samples for the β -giardin gene, finding an exact match with sub-assemblage AII and no sequence within the three sample inspected. By contrast, each of the three samples for assemblage B was different and did not match either BIII and BIV precisely. The sequence from Sample 102 is particularly notable as there appeared to be allelic variation within the TPI gene as evidenced by split-peak chromatograms of A/G or T/C at three locations present within this region (see Annex Supplemental Fig. 1).

4. Discussion

The high prevalence of giardiasis reported here by real-time PCR with the 18S rDNA probe analysis (87.0%) demonstrates that children living on the shoreline of Lake Albert are at very high risk of both acute and more likely, chronic infections. The high burden of giardiasis was also corroborated in field by the Quik-Chek RDT at Runga and Bugoigo schools confirming that some 41.6% of children were patently shedding copious amounts of *Giardia* cysts within their stools. It is unsurprising perhaps that the levels are so high since this lakeshore environment has very poor local sanitation and water hygiene, as well as being hyper-endemic for intestinal schistosomiasis, another waterborne disease (Al-Shehri et al., 2016). Nonetheless the prevalence of giardiasis here is much elevated in comparison to other parts of the world (Thompson and Smith, 2001), although in Rwanda over 60% of rural children have been shown to be infected with *Giardia* by molecular typing methods (Ignatius et al., 2014). More broadly, the diagnostic sensitivity of real-time PCR methods is known to be superior to alternative diagnostic methods, often




Fig. 1. Bivariate plot of Ct values obtained for each sample using *Giardia* TaqMan® 18S rDNA versus Ct value of assemblage-specific TaqMan® TPI probe with dashed lines showing the 95% prediction interval. Fig. 1A. Using assemblage A probe; Fig. 1B Using assemblage B probe.

Table 1

Prevalence (%) of *G. duodenalis* and assemblages across all five schools by real-time PCR; the odds ratio of assemblages A, B or A/B by school compared against the total given (with 95% confidence limits).

School	Giardia TaqMan® 18S rDNA probe		Assemblage (A & B) TaqMan® TPI probe				
	Number of positives % (x/y)	95% CL	Number of positives % (x/y)	95% CL	A % (x/y) OR [95% CI]	B % (x/y) OR [95% CI]	AB % (x/y) OR [95% CI]
Bugoigo	94.5% (52/55)	[85.8–98.6]	56.3% (31/55)	[43.1–69.0]	5.4% (3/55) 0.6 [0.2, 2.5]	43.6% (24/55) 1.4 [0.8, 2.8]	7.2% (4/55) 0.9 [0.3, 3.1]
Runga	94.1% (48/51)	[84.8–98.5]	72.5% (37/51)	[59.2–83.4]	15.6% (8/51) 4.7 [1.7, 13.3]	37.2% (19/51) 2.0 [1.0, 4.3]	19.6% (10/51) 6.9 [2.5, 19.3]
Walukuba	88.0% (44/50)	[76.7–95.0]	40.0% (20/50)	[27.2–54.0]	2.0% (1/50) 0.1 [0.0, 1.2]	32.0% (16/50) 0.6 [0.3, 1.8]	6.0% (3/50) 0.5 [0.1, 1.8]
Biiso	84.0% (42/50)	[71.9–92.3]	54.0% (27/50)	[40.2–67.4]	14.0% (7/50) 2.1 [0.8, 5.9]	36.0% (18/50) 1.0 [0.5, 2.1]	4.0% (2/50) 0.4 [0.1, 2.1]
Busingiro	72.9% (35/48)	[59.1–84.0]	37.5% (18/48)	[24.7–51.8]	4.1% (2/48) 0.3 [0.1, 1.5]	29.1% (14/48) 0.5 [0.3, 1.1]	4.1% (2/48) 0.3 [0.1, 1.5]
All	87.0% (221/254)	[82.4–90.7]	52.4% (133/254)	[46.2–58.5]	8.3% (21/254)	35.8% (91/254)	8.3% (21/254)

revealing giardiasis to be more pervasive (Gotfred-Rasmussen et al., 2016), and also creates opportunities for investigations of (sub)assemblage transmission dynamics (Thompson and Ash, 2016).

Given the multi-copy nature of the 18S rDNA against the lower copy number of TPI, the diagnostic sensitivity of TPI probes is lower, such that just under a half of the infected cases detected by 18S rDNA were missed. It has been stated previously that the detection limit of *Giardia* 18S rDNA probe assay is approaching 10 pg DNA/μL (Jaros et al., 2011), presumably that of TPI assay is much higher (Elwin et al., 2014) such that assemblage typing of 'light' intensity infections is not always possible. A similar level of diagnostic discordance has been observed elsewhere (Ignatius et al., 2014) which hopefully does not lead to a systematic bias in general reporting of each assemblage, as evidenced by Ct values in Fig. 1, but rather that typing parasites with assemblage-specific primers is not possible when shedding cysts are too few in number.

Nonetheless, in this sample assemblage B dominates upon comparison to assemblage A. Notably this 1:2.7 ratio varied by school with Runga having a greater proportion of assemblage A, as well as co-infection with assemblage B thereof, see Table 1, and more broadly, there appeared to be some interesting epidemiological associations by assemblage, see Table 2. Although there was no association with gender, younger children appeared to harbour a greater proportion of mixed assemblage infections than older counterparts (OR = 2.7 [95% CI; 1.0, 7.3]). There was also an indication that faecal occult blood was associated with assemblage B (OR = 2.2 [95% CI; 1.0, 5.2]) and in children being underweight (OR = 2.0 [95% CI; 1.0, 3.9]). These findings add to the general debate on the health consequences of giardiasis with particular emphasis on assemblage B, which also appears more genetically heterogeneous than assemblage A here (Thompson and Ash, 2016).

It is an interesting observation that of the six samples subjected to sequence analysis of β-giardin, the three samples selected from assemblage A were identical and could be further unequivocally assigned to sub-lineage AII, which has been reported in

Table 2

Analysis of potential epidemiological associations by binary logistic regression with *Giardia* assemblages A, B or A/B co-infection.

Epidemiological factors	Assemblage (A, B & AB)® TaqMan probe (TPI)					
	Infected with A	OR [95% CI]	Infected with B	OR [95% CI]	Infected with AB	OR [95% CI]
Gender						
Male	10	1.0 [0.4, 2.7]	44	1.0 [0.6, 1.9]	12	1.5 [0.6, 3.9]
Female	11	0.9 [0.4, 2.4]	47	0.9 [0.5, 1.6]	9	0.6 [0.3, 1.6]
Age group						
5 to 6	11	1.5 [0.6, 3.8]	38	0.9 [0.6, 1.7]	14	2.7 [1.0, 7.3]
7 to 8	9	0.9 [0.4, 2.5]	34	0.7 [0.5, 1.4]	6	0.5 [0.2, 1.5]
9 to 10	1	0.2 [0.0, 2.3]	19	1.5 [0.7, 3.1]	1	0.2 [0.0, 2.3]
Faecal occult blood (FOB)						
Negative	18	0.6 [0.2, 2.4]	74	0.4 [0.2, 1.0]	17	0.4 [0.1, 1.5]
Positive	3	1.6 [0.4, 6.6]	17	2.2 [1.0, 5.2]	4	2.3 [0.7, 8.2]
Height-for-age Z score, mean						
−2 > SD height-for-age Z score	17	1.3 [0.4, 4.3]	63	0.7 [0.4, 1.3]	14	0.6 [0.2, 1.7]
−2 ≤ SD height-for-age Z score	4	0.7 [0.2, 2.4]	28	1.4 [0.8, 2.6]	7	1.5 [0.6, 4.3]
Weight-for-age Z score, mean						
−2 > SD weight-for-age Z score	20	3.9 [0.5, 31.2]	65	0.4 [0.3, 1.0]	16	0.6 [0.2, 1.9]
−2 ≤ SD weight-for-age Z score	1	0.2 [0.0, 2.0]	26	2.0 [1.0, 3.9]	5	1.5 [0.5, 4.8]
Anaemia (<115 Hg/L)						
Negative	9	0.4 [0.1, 1.4]	46	0.5 [0.3, 1.2]	9	0.3 [0.1, 1.0]
Positive	6	2.3 [0.7, 7.5]	23	1.7 [0.8, 3.7]	8	3.1 [1.1, 9.4]
Not determined	6	–	22	–	4	–

Table 3
Single nucleotide polymorphisms within β -giardin of *Giardia duodenalis*.

Assemblage	Isolate/Genbank number	Nucleotide position								
A isolates	Beta-giardin (bg)	284	383	407	473	491	563	593	596	611
AI	X14185.1	C	T	T	T	A	G	T	C	A
AII	AY072723.1	C	T	T	T	A	G	T	T	A
AIII	DQ650649.1	T	C	C	C	G	A	C	C	G
	Sample 9	C	T	T	T	A	G	T	T	A
	Sample 22	C	T	T	T	A	G	T	T	A
	Sample 103	C	T	T	T	A	G	T	T	A
B isolates		170	176	188	233	287	314	317	398	
BIII	AY072726.1	C	A	A	G	C	C	C	C	
BIV	AY072725.1	T	A	A	A	T	T	T	T	
	Sample 24	C	A	A	A	C	T	T	C	
	Sample 104	C	A	A	A	C	C	T	C	
	Sample 102	C	A/G	A/G	A	C	T/C	T	C	

other studies (Cacciò and Ryan, 2008; Plutzer et al., 2010; Cacciò and Sprong, 2010; Ryan and Cacciò, 2013; Beck et al., 2012; Zhang et al., 2012). By contrast, of the three samples selected from assemblage B, there were each different, see Table 3, and none matched exactly either BIII or BIV sub-assemblages. Most notable are the point mutations at positions 176, 188 and 314, where split-peak chromatograms were observed (see Annex). This is indicative of mixed amplicon templates inferring putative allelic variation within the TPI locus. The genomic complexity of *Giardia* is complex, being binucleate and sometimes aneuploid (Aguilar et al., 2016) which might infer sample 102 was either a mixed co-infection of two independent B lineages or contains a single infection lineage with an unusual genomic TPI variant. Nonetheless, there is greater diversity within assemblage B and with further genetic profiling would reveal additional variants which might point towards currently unknown heterogeneities in local transmission cycles. For example, there is numerous livestock e.g. cattle and goats, that regularly enter into the lake and while drinking openly defecate into the water which may add to raised zoonotic potential in such domestic water directly drawn from the lake.

To conclude, additional sampling of school children would be worthwhile if putative associations between assemblage B and detrimental health outcomes reported here are to be fully verified statistically. Furthermore to better monitor local transmission cycles of *Giardia*, we encourage future studies that track each assemblage within local livestock and undertake environmental sampling of lake water where domestic water is drawn.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.parepi.2018.e00074>.

Acknowledgements

We wish to thank the VCD field team for their support during the fieldwork and Dr. Jaco Verweij for technical assistance in real-time PCR. HA is grateful to receive sponsored PhD studentship from the Ministry of Health, Kingdom of Saudi Arabia.

References

- Adam, R.D., 2001. Biology of *Giardia lamblia*. Clin. Microbiol. Rev. 14 (3), 447–475.
- Aguilar, J.M., et al., 2016. Evidence of heterozygosity and recombinant alleles in single cysts of *Giardia duodenalis*. Revista Brasileira De Parasitologia Veterinaria 25 (2), 187–195.
- Almeida, A., Pozio, E., Cacciò, S.M., 2010. Genotyping of *Giardia duodenalis* cysts by new real-time PCR assays for detection of mixed infections in human samples. Appl. Environ. Microbiol. 76 (6), 1895–1901.
- Al-Shehri, H., et al., 2016. An extensive burden of giardiasis associated with intestinal schistosomiasis and anaemia in school children on the shoreline of Lake Albert, Uganda. Trans. R. Soc. Trop. Med. Hyg. 110 (10), 597–603.
- Ankarklev, J., et al., 2012. Common coinfections of *Giardia intestinalis* and *Helicobacter pylori* in non-symptomatic Ugandan children. PLoS Negl. Trop. Dis. 6 (8), e1780.
- Asher, A.J., et al., 2014. Distribution of *Giardia duodenalis* assemblages A and B among children living in a remote indigenous community of the Northern Territory, Australia. PLoS One 9 (11), e112058.
- Beck, R., et al., 2012. Genotyping *Giardia duodenalis* isolates from dogs: lessons from a multilocus sequence typing study. Vector-Borne and Zoonotic Diseases 12 (3), 206–213.
- Cacciò, S.M., Ryan, U., 2008. Molecular epidemiology of giardiasis. Mol. Biochem. Parasitol. 160 (2), 75–80.
- Cacciò, S.M., Sprong, H., 2010. *Giardia duodenalis*: genetic recombination and its implications for taxonomy and molecular epidemiology. Exp. Parasitol. 124 (1), 107–112.
- Durigan, M., et al., 2014. Genetic diversity of *Giardia duodenalis*: multilocus genotyping reveals zoonotic potential between clinical and environmental sources in a metropolitan region of Brazil. PLoS One 9 (12), e115489.
- Elwin, K., et al., 2014. *Giardia duodenalis* typing from stools: a comparison of three approaches to extracting DNA, and validation of a probe-based real-time PCR typing assay. J. Med. Microbiol. 63 (1), 38–44.
- Feng, Y., Xiao, L., 2011. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. Clin. Microbiol. Rev. 24 (1), 110–140.
- Fuhrmann, S., et al., 2016. Risk of intestinal parasitic infections in people with different exposures to waste water and fecal sludge in Kampala, Uganda: a cross-sectional study. PLoS Negl. Trop. Dis. 10 (3), e0004469.
- Gotfred-Rasmussen, H., et al., 2016. Comparison of sensitivity and specificity of 4 methods for detection of *Giardia duodenalis* in feces: immunofluorescence and PCR are superior to microscopy of concentrated iodine-stained samples. Diagn. Microbiol. Infect. Dis. 84 (3), 187–190.
- Graczyk, T.K., et al., 2002. Anthrozoönotic *Giardia duodenalis* genotype (assemblage) A infections in habitats of free-ranging human-habituated gorillas, Uganda. J. Parasitol. 88 (5), 905–909.

- Helmy, Y.A., et al., 2014. Epidemiology of *Giardia duodenalis* infection in ruminant livestock and children in the Ismailia province of Egypt: insights by genetic characterization. *Parasit. Vectors* 7 (1), 321.
- Ignatius, R., et al., 2012. High prevalence of *Giardia duodenalis* Assemblage B infection and association with underweight in Rwandan children. *PLoS Negl. Trop. Dis.* 6 (6), e1677 2012.
- Ignatius, R., et al., 2014. Detection of *Giardia duodenalis* assemblage A and B isolates by immunochromatography in stool samples from Rwandan children. *Clin. Microbiol. Infect.* 20 (10), O783–O785.
- Jaros, D., et al., 2011. Detection of *Giardia intestinalis* assemblages A, B and D in domestic cats from Warsaw, Poland. *Pol. J. Microbiol.* 60, 259–263.
- Johnston, A.R., et al., 2010. Molecular epidemiology of cross-species *Giardia duodenalis* transmission in western Uganda. *PLoS Negl. Trop. Dis.* 4 (5), e683.
- Karim, M.R., et al., 2015. Multi-locus analysis of *Giardia duodenalis* from non-human primates kept in zoos in China: geographical segregation and host-adaptation of assemblage B isolates. *Infect. Genet. Evol.* 30, 82–88.
- Laishram, S., Kang, G., Ajjampur, S.S.R., 2012. Giardiasis: a review on assemblage distribution and epidemiology in India. *Indian J. Gastroenterol.* 31 (1), 3–12.
- Minetti, C., et al., 2015. Determination of *Giardia duodenalis* assemblages and multi-locus genotypes in patients with sporadic giardiasis from England. *Parasit. Vectors* 8 (1), 444.
- Muhsen, K., Levine, M.M., 2012. A systematic review and meta-analysis of the association between *Giardia lamblia* and endemic pediatric diarrhea in developing countries. *Clin. Infect. Dis.* 55, S271–S293.
- Nizeyi, J.B., et al., 1999. *Cryptosporidium* sp. and *Giardia* sp. infections in mountain gorillas (*Gorilla gorilla beringei*) of the Bwindi Impenetrable National Park, Uganda. *J. Parasitol.* 85, 1084–1088.
- Nizeyi, J., Cranfield, M., Graczyk, T., 2002. Cattle near the Bwindi Impenetrable National Park, Uganda, as a reservoir of *Cryptosporidium parvum* and *Giardia duodenalis* for local community and free-ranging gorillas. *Parasitol. Res.* 88 (4), 380–385.
- Pelayo, L., et al., 2008. *Giardia* infections in Cuban children: the genotypes circulating in a rural population. *Ann. Trop. Med. Parasitol.* 102 (7), 585–595.
- Plutzer, J., Ongerth, J., Karanis, P., 2010. *Giardia* taxonomy, phylogeny and epidemiology: facts and open questions. *Int. J. Hyg. Environ. Health* 213 (5), 321–333.
- Puebla, L.J., et al., 2014. Correlation of *Giardia duodenalis* assemblages with clinical and epidemiological data in Cuban children. *Infect. Genet. Evol.* 23, 7–12.
- Rogawski, E.T., et al., 2017. Determinants and impact of *Giardia* infection in the first 2 years of life in the MAL-ED birth cohort. *Journal of the Pediatric Infectious Diseases Society.* 6 (2), 153–160.
- Ryan, U., Cacciò, S.M., 2013. Zoonotic potential of *Giardia*. *Int. J. Parasitol.* 43 (12–13), 943–956.
- Sprong, H., Cacciò, S.M., van der Giessen, J.W., 2009. Identification of zoonotic genotypes of *Giardia duodenalis*. *PLoS Negl. Trop. Dis.* 3 (12), e558.
- Takumi, K., et al., 2012. Population-based analyses of *Giardia duodenalis* is consistent with the clonal assemblage structure. *Parasit. Vectors* 5 (1), 168.
- Tellevik, M.G., et al., 2015. Prevalence of *Cryptosporidium parvum/hominis*, *Entamoeba histolytica* and *Giardia lamblia* among young children with and without diarrhea in Dar es Salaam, Tanzania. *PLoS Negl. Trop. Dis.* 9 (10).
- Thompson, R.C.A., Ash, A., 2016. Molecular epidemiology of *Giardia* and *Cryptosporidium* infections. *Infect. Genet. Evol.* 40, 315–323.
- Thompson, R., Smith, A., 2001. Zoonotic enteric protozoa. *Vet. Parasitol.* 182 (1), 70–78.
- Vanni, I., et al., 2012. Detection of *Giardia duodenalis* assemblages A and B in human feces by simple, assemblage-specific PCR assays. *PLoS Negl. Trop. Dis.* 6 (8), e1776.
- Verweij, J.J., et al., 2004. Simultaneous detection of *Entamoeba histolytica*, *Giardia lamblia*, and *Cryptosporidium parvum* in fecal samples by using multiplex real-time PCR. *J. Clin. Microbiol.* 42 (3), 1220–1223.
- Wegayehu, T., et al., 2016. Multilocus genotyping of *Giardia duodenalis* isolates from children in Oromia Special Zone, Central Ethiopia. *BMC Microbiol.* 16 (1), 89.
- WHO, 2007. Growth Reference Data for 5–19 Years. World Health Organization, Geneva.
- WHO, 2011. Haemoglobin Concentrations for the Diagnosis of Anaemia and Assessment of Severity. Download from: <http://www.who.int/vmnis/indicators/haemoglobin.pdf> (2015).
- Zhang, W., et al., 2012. Genetic characterizations of *Giardia duodenalis* in sheep and goats in Heilongjiang Province, China and possibility of zoonotic transmission. *PLoS Negl. Trop. Dis.* 6 (9), e1826.