

Aetiology and outcomes of sepsis in adults in sub-Saharan Africa: a systematic review and meta-analysis

Joseph M LEWIS^{1,2}, Nicholas FEASEY^{1,2}, Jamie RYLANCE^{1,2}

1 Liverpool School of Tropical Medicine, Liverpool, UK

2 Malawi Liverpool Wellcome Clinical Research Programme, Blantyre, Malawi

Corresponding author:

Joseph M Lewis

Liverpool School of Tropical Medicine

Pembroke Place

Liverpool

L3 5QA

E: joseph.lewis@lstmed.ac.uk

T: +44 7969 419910

Nicholas Feasey

E: Nicholas.feasey@lstmed.ac.uk

Jamie Rylance

E: Jamie.rylance@lstmed.ac.uk

Aetiology and outcomes of sepsis in adults in sub-Saharan Africa: a systematic review and meta-analysis

ABSTRACT

Background

Aetiology and outcomes of sepsis in sub-Saharan Africa (sSA) are poorly described; we performed a systematic review and meta-analysis to summarise available data.

Methods

Systematic searches of PubMed and Scopus were undertaken to identify prospective studies recruiting adults (>13 years) with community-acquired sepsis in sSA post-2000. Random effects meta-analysis of in-hospital and 30-day mortality was undertaken and available aetiology data also summarized by random effects meta analysis.

Results

15 studies of 2800 participants were identified. Inclusion criteria were heterogeneous. The majority of patients were HIV infected, and *M. tuberculosis* was the most common cause of blood stream infection where sought. Pooled in-hospital mortality for Sepsis-2 defined sepsis and severe sepsis were 19% (95% CI 12-29%) and 39% (95% CI 30-47%) respectively and sepsis mortality was associated with the proportion of HIV-infected participants. Mortality and morbidity data beyond 30 days were absent.

Conclusions

Sepsis in sSA is dominated by HIV and tuberculosis, with poor outcomes. Optimal antimicrobial strategies, including the role of tuberculosis treatment, are unclear.

Long term outcome data are lacking. Standardised sepsis diagnostic criteria that are easily applied in low resource settings are needed to establish an evidence base for sepsis management in sSA.

Key words: Sepsis, Africa south of the Sahara, Bloodstream Infection, Tuberculosis, HIV

Aetiology and outcomes of sepsis in adults in sub-Saharan Africa: a systematic review and meta-analysis

INTRODUCTION

Sepsis, defined most recently as a syndrome of life-threatening organ dysfunction due to a dysregulated host response to infection[1], is common worldwide and carries a high mortality: recent estimates suggest 19.4 million yearly cases and 5.3 million deaths[2]. In high income settings, outcomes are improving, due in part to a comprehensive application of an expanding evidence base for early recognition, rapid administration of appropriate antimicrobials, and aggressive fluid resuscitation paired with intensive monitoring of physiology and provision for organ support[3,4]. In low resource settings including sub-Saharan Africa (sSA), data are limited but some studies have identified high mortality[5]. It is clear that sepsis protocols developed in high-income settings should not simply be exported unchanged to sSA: aggressive fluid resuscitation has been shown to be harmful in one RCT in adults and one in children[6,7] and caution is warranted before proposing fluid management guidelines for sSA.

The paucity of data presents challenges in proposing sepsis management specific to sSA. Firstly, defining sepsis for clinical practice or research is problematic. Recent Sepsis-3 guidelines (the third iteration of the international consensus diagnostic definitions of sepsis) suggest operationalising the diagnosis of sepsis using the sequential organ failure assessment (SOFA) score[1]. Applying this score to resource-limited settings is difficult due to patchy availability of variables, particularly

laboratory values. The bedside “quick SOFA” (qSOFA) score can identify patients at a higher risk of death, but is a screening rather than a diagnostic tool[8–10]. Previous iterations of the guidelines defined sepsis using the systemic inflammatory response score (SIRS) in the presence of a suspicion of infection, with severe sepsis defined by the addition of organ dysfunction[11], but SIRS, while applicable at the bedside, has been criticised for its lack of discriminatory power[12].

Secondly, the optimal clinical management of sepsis in sSA is unknown. Early, appropriate antimicrobials improve outcomes in high-income settings[13,14] and it is likely that this is a transferable recommendation to sSA. Certainly, it seems unlikely that rapid administration of antimicrobials will adversely affect outcomes in the way aggressive fluid resuscitation does, and may represent an important first step in improving outcomes. However, in the absence of robust sepsis aetiology data, what constitutes “appropriate” empirical antimicrobial chemotherapy remains an open question in sSA. Tuberculosis and malaria certainly play an important role, and arboviral infections, bacterial zoonoses and HIV opportunistic infections may also be important[15–19].

Finally, sepsis outcomes in sSA are heterogeneous between studies[20,21] and longer term follow up sparse. Post-discharge sepsis mortality in high-income settings is significant (11 – 43% at one year) as is morbidity[22], but post-discharge outcomes in sSA are unknown.

We therefore performed a systematic review and meta-analysis of clinical studies from sSA with three aims: to describe the sepsis definitions used, the aetiology of infection, and clinical outcomes.

METHODS

The protocol of this systematic review was pre-registered on PROSPERO, number CRD42019123589, and follows PRISMA guidelines.

Searches

Searches of PubMed and Scopus were undertaken using the search terms shown in Box 1, in all fields. Identified abstracts were exported into Endnote X7.8 (Thomson Reuters, United States) and screened against inclusion and exclusion criteria.

Inclusion criteria were: prospective cohort studies, non-randomised intervention studies or randomised controlled trials recruiting adults (> 13 years) with community acquired sepsis from sSA, for which it was possible to disaggregate a total number of adults with sepsis and was possible to extract aetiology or outcome data. Any study-defined definition of sepsis was accepted. Retrospective studies were excluded due to a high risk of bias: we were concerned that, in low resource settings medical records can often be incomplete and this could introduce significant bias. Exclusion criteria were studies published before 2000 and studies recruiting preselected populations e.g. puerperal sepsis. There was no language restriction.

Sepsis AND ((Angola OR Benin OR Botswana OR Burkina Faso OR Burundi OR Cameroon OR Cape Verde OR Central African Republic OR Chad OR Comoros OR Republic of the Congo OR Congo Brazzaville OR Democratic republic of the Congo OR Cote d'Ivoire OR Djibouti OR Equatorial Guinea OR Eritrea OR Ethiopia OR Gabon OR The Gambia OR Ghana OR Guinea OR Guinea-Bissau OR Kenya OR Lesotho OR Liberia OR Madagascar OR Malawi OR Mali OR Mauritania OR Mauritius OR Mozambique OR Namibia OR Niger OR Nigeria OR Reunion OR Rwanda OR Sao Tome and Principe OR Senegal OR Seychelles OR Sierra Leone OR Somalia OR South Africa OR Sudan OR Swaziland OR Eswatini OR Tanzania OR Togo OR Uganda OR Western Sahara OR Zambia OR Zimbabwe) OR Africa)

Box 1: Systematic review search terms.

Abstracts were screened by two authors (JL and JR) and disagreements resolved by consensus. All included abstracts underwent independent full text review by the same authors and the inclusion and exclusion criteria were again applied, and disagreements resolved by consensus. Data were extracted onto an Excel spreadsheet (Microsoft, United States) for further analysis: study first author, years of recruitment, inclusion and exclusion criteria, number of patients recruited and demographics (including age, proportion of HIV infected participants and CD4 count values), in-hospital, 28- or 30-day mortality, and details of any aetiological investigations undertaken.

Quality Assessment

A modified Newcastle-Ottawa scale was used to assess risk of bias in the domains of selection, comparability and outcome (Full scale shown in Additional File 1) This was completed by two authors independently and disagreements resolved by consensus to provide a single assessment of each study incorporating all domains.

Statistical analysis

Because of concerns about meta-analysis of proportions on very heterogeneous populations, we planned meta-analysis of outcome stratified by inclusion criteria where possible: Sepsis-2 sepsis, severe sepsis, and Sepsis-3 sepsis, if available. Mortality was presented as a simple proportion with exact binomial confidence intervals and pooled mortality estimates were calculated using generalized linear mixed models (a normal-binomial model) using the packages *meta* and *lme4* in R. For interventional studies, the outcomes in the usual care arm of the study only were included in these estimates. Heterogeneity was quantified with τ^2 , I^2 and Cochran's Q test. Exploratory meta regression was undertaken to explore heterogeneity by including covariates as fixed effects (year of recruitment, proportion of patients infected with HIV and median age) and testing for improved model fit by likelihood ratio testing of nested models. A p value of <0.05 was considered a statistically significantly improved fit. Bubble plots of the meta regressions were produced with 95% confidence intervals obtained from 1000 bootstrap replicates. Summary estimates of 28- and 30-day mortality, where available, were considered together, and were presented in the same way. Pooled prevalence estimates of malaria,

bloodstream infection (BSI), and *M. tuberculosis* bloodstream infection (MTB-BSI) were calculated using random effects meta-analysis as above. For these aetiology analyses we included all studies, regardless of sepsis definition, and included both usual care and intervention arms of RCTs. All analysis was carried out in R V3.5.1 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

The abstract search yielded 3951 unique records on 17 July 2018 (Figure 1), of which 3902 were excluded. After screening 49 full-text articles, 15 were retained. One article was a secondary analysis of the aetiology of a previously presented cohort[23], meaning 15 articles were included, reporting on 14 prospective clinical studies from nine centers in six countries. These included 11 cohort studies[5,20,21,24–31], two randomized controlled trials[32,33] and one before-after interventional trial[34] (Table 1).

Data from 2800 unique participants were eligible (Table 1). Data from some participants were included in two or more separate publications, but we were able to extract aggregate data such that no participant contributed data twice. Generally, the risk of bias as assessed by the modified Newcastle-Ottawa scale was low (Additional File 2). The most commonly identified areas of concern were in ascertaining exposure (all studies ascertained HIV status but 6/14 studies lacked details of HIV testing procedures) or in comparability (5/14 studies did not test lactate and/or provide details of enrollment physiology, both components of the comparability domain of our modified Newcastle-Ottawa score). The patients recruited to the

identified studies had median age ranging from 27 to 39 years, and were predominantly HIV-infected: 2577/2800 patients had an available HIV status, with 1712/2577 (66%) being HIV-infected, though HIV rates varied between studies (median 70% [IQR 42.5-82.5%], Table 1)

The majority of studies recruited patients using a modified Sepsis-2 definition of sepsis or severe sepsis, though definitions were heterogeneous (primary study inclusion and exclusion criteria are shown in Additional File 3). SIRS was used to define sepsis or severe sepsis in 13/14 cohorts, but the definition of SIRS itself was variable. Six of the 13 studies using SIRS did not use the white cell count criterion because of resource limitations, and four different temperature thresholds were used in the 13 studies to define hypo- or hyperthermia.

Of the 13 studies using SIRS as a component of inclusion criteria, 8/13 recruited patients with a definition compatible with Sepsis-2 sepsis (SIRS plus suspected or confirmed infection, 1255 patients in total) and 5/13 studies recruited a population compatible with Sepsis-2 severe sepsis of SIRS plus organ dysfunction (1347 patients in total). A variety of organ dysfunction criteria were used including hyperlactataemia or poor Karnofsky performance score; the most common organ dysfunction criterion was low systolic blood pressure, applied in all five studies. No study specifically recruited patients with septic shock and despite the frequent use of low systolic blood pressure as a defining criterion for severe sepsis, only one study[28] explicitly defined septic shock, using a definition of hypotension (SBP < 90 mmHg or MAP < 65mmHg) refractory to 2L of intravenous fluid administered over 2 hours; 27 of 150 recruited patients fulfilled this definition.

The one study that used a non-Sepsis 2 based definition defined sepsis as fever plus one severity criterion[25], adapted from WHO severe malaria definitions.

Author and publication year	Years recruiting	Study type	Country	N centres	Centre type	Inc. criteria	n	Male sex	Age	HIV infected	Median CD4 μ L $^{-1}$
Jacob 2009[5]	2006	Cohort	Uganda	2	Referral	SII Severe sepsis	382	156/382 (41%)	35	320/382 (85%)	52
Nadjm 2012[25]	2007	Cohort	Tanzania	1	District	Fever and one severity criteria	198	67/198 (34%)	37	69/180 (38%)	NR
Jacob 2012[34] ^a and Moore 2018[23]	2008-09	Before -after	Uganda	2	Referral	SII Severe sepsis	426	207/426 (49%)	34	362/426 (85%)	63
Waitt 2015[26]	2008-09	Cohort	Malawi	1	Referral	SII Sepsis	213	87/213 (41%)	30	161/213 (76%)	NR
Ssekitoleko 2011[27] ^a	2009	Cohort	Uganda	1	Referral	SII Sepsis	96	193/418 ^b (46%)	35	331/418 ^b (83%)	NR
Ssekitoleko 2011[28]	2009	Cohort	Uganda	1	Referral	SII Sepsis	150	94/150 (63%)	35	96/150 (64%)	NR
Chimese 2012[21]	2010	Cohort	Zambia	1	Referral	SII Sepsis	161	79/161 (49%)	39	110/138 (80%)	NR
Andrews 2014[35]	2012	RCT	Zambia	1	Referral	SII Severe sepsis	112	58/109 (53%)	35	88/109 (81%)	NR
Auma 2013[29]	2012	Cohort	Uganda	1	Referral	SII Sepsis	216	106/216 (49%)	32	122/216 (56%)	NR ^e
Andrews 2017[33]	2012-13	RCT	Zambia	1	Referral	SII Severe sepsis	209	117/209 (56%)	36	187/209 (90%)	66
Huson 2014[20]	2012-13	Cohort	Gabon	1	Referral	SII Sepsis	107	NA	34	26/107 (24%)	168
Seboxa 2015[30]	2012-13	Cohort	Ethiopia	1	Referral	SII sepsis	292	151/292 (52%)	27	40/209 (19%)	NR
Rudd 2017[31]	2013	Cohort	Uganda	1	District	SII sepsis	20	11/20 (55%)	32	6/20 (30%)	NR
Amir 2016[24]	2014-15	Cohort	Uganda	1	Referral	SII Severe sepsis	218	110/218 (50%)	35	125/218 (57%)	78

Table 1: Characteristics of included studies.

RCT = randomised controlled trial. SII = sepsis-2 definition (e.g. SII sepsis is consistent with sepsis 2 definition). ^a These studies also presents data from Jacob 2009 – only new data is included in this table row. ^b disaggregated data not given for the included 96 patients; HIV prevalence for total cohort shown.

Outcomes

All studies reported either in-hospital or 28/30 day mortality, apart from one retrospective of stored samples [23], leaving 14 studies with outcome data. Of these, 11 reported only in-hospital mortality, 3 studies reported both 28/30 day mortality and in-hospital mortality, and 1 reported only 28/30 day mortality. No studies reported longer-term outcomes or estimates of morbidity. We therefore pooled the outcome data from the 13 studies with outcome data that used Sepsis-2 based inclusion criteria: in hospital mortality data were available for Sepsis-2 compatible sepsis (1159 participants from 7 studies) and severe sepsis (983 participants from 7 studies). 28/30 day mortality data were only available for severe sepsis (484 patients from 3 studies). Pooled estimates of Sepsis-2 defined sepsis and severe sepsis in-hospital mortality were 19% (95% CI 12-29%) and 39% (95% 30-47%) respectively and pooled 28- and 30-day Sepsis-2 defined severe sepsis mortality was 54% (95% 37-70%), though significant between-study heterogeneity means these summary estimates should be treated with caution (Figure 2).

Heterogeneity was explored with meta-regression. Proportion of HIV-infected participants was significantly associated with inpatient mortality in studies recruiting patients with Sepsis-2 defined sepsis when included as a fixed-effect covariate ($p=0.006$ on likelihood ratio testing of nested models), but year of recruitment was not ($p=0.06$). Neither year nor proportion of HIV-infected participants ($p=0.51$, and $p=0.83$) were associated with inpatient mortality in Sepsis-2 severe sepsis studies. Severe sepsis studies in general appeared to have higher proportion of HIV-infected participants than studies recruiting patients with sepsis (Figure 3); in only one

study[26] was it was possible to extract data to compare HIV prevalence in sepsis and severe sepsis in the same study: HIV prevalence was higher in severe sepsis in absolute terms though numbers were small and the difference was not statistically significant (HIV prevalence 26/32 [81%] severe sepsis vs 161/213 [76%] in sepsis, p = 0.89). Median age was not associated with in-hospital mortality in sepsis or severe sepsis analyses (p = 0.66 and p = 0.50 on likelihood ratio testing of nested models). In view of the small number of studies reporting 28- or 30-day mortality, meta-regression was not undertaken.

Outcome data on septic shock were largely absent. For only 27 patients from the only study which defined septic shock was it possible to extract in-hospital mortality: 16/27 (59%, 95% CI 39-77%) patients died.

Aetiology

Aetiology data could be extracted from 14 studies: 11/14 reported aerobic blood culture data, 4/14 mycobacterial blood culture and 9/14 malaria data (Figure 4) and 1/14 reported retrospective polymerase chain reaction (PCR) diagnostics on stored samples from Jacob 2012. The data shown in Figure 4 are those from the original Jacob 2012 study and not the retrospective PCR diagnostics; the latter are described below. Generally, studies performed aetiologic testing on all patients rather than restricting to a subgroup, excepting mycobacterial blood culture (carried out in 4 studies) which was restricted to HIV-infected participants in 2/4 studies and to a single study site in 1/4 studies (details of aetiologic testing availability by study are shown in Additional file 4).

Diagnosis of malaria was made by smear in all studies except the retrospective PCR study; one[29] additionally used rapid diagnostic tests and malaria-specific PCR. Pathogenic organisms isolated from aerobic blood culture are shown in Additional File 5. The commonest isolated organisms were *Staphylococcus aureus* (105/458), non-typhoidal *Salmonella* (75/458) and *Streptococcus pneumoniae* (68/458). All studies reporting blood culture results described the microbiologic methods used but only 6/11 reported external quality control procedures during the study period.

Data on antimicrobial sensitivity patterns of the aerobic blood culture isolates were available in only 2/11 studies. One study[5] gave limited details only, stating that 95% of *Salmonella* isolates were resistant to chloramphenicol and trimethoprim-sulfamethoxazole and that none of the *Staphylococcus aureus* samples were resistant to oxacillin. The second study provided more comprehensive details, but with very small numbers which limit the conclusions that can be drawn; of 20 isolated Gram-negative bacilli (16 *E. coli*, the remainder *Klebsiella*, *Salmonella* and *Citrobacter spp.*), 9/20 (45%) were resistant to third-generation cephalosporins, and this was strongly associated with mortality. All patients with third-generation cephalosporin resistant BSI died, compared with 1/11 (9%) of those with sensitive infections ($p < 0.0001$).

One study[23] reported the results of a multiplex PCR on cryopreserved blood specimens of 336/426 participants, and identified at least one potential pathogen in 245/336 samples, most commonly cytomegalovirus (139/336 [41%]), MTB (70/336

[21%]), *Plasmodium* spp. (35/336 [10%]) and *Streptococcus pneumoniae* (31/226 [9%]). Bacterial zoonoses or rickettsioses were uncommon (6/336 [2%]).

DISCUSSION

In-hospital sepsis and severe sepsis mortality in sSA is high. We found pooled in-hospital mortality of 19% (95% CI 12-29%) for Sepsis-2 defined sepsis and 39% (95% 30-47%) for Sepsis-2 defined severe sepsis. Pooled 28/30 day Sepsis-2 defined severe sepsis mortality was 54% (95% CI 37-70%). Some between-study heterogeneity in sepsis mortality is likely consequent upon heterogenous inclusion criteria, and some variability may be explained by the proportion of participants who are HIV-infected. Direct comparison of these mortality estimates to outcomes in high-income settings is difficult as estimates of sepsis mortality often derive from administrative databases and are sensitive to coding strategies. A recent meta-analysis[2] of population level estimates in post-2003 high income settings estimated pooled sepsis and severe sepsis 30-day mortality to be of 17% (95% CI 11-26%) and 26% (95% CI 20-33%), respectively. The large US patient cohort used to validate the qSOFA score[8] (n= 74,453) found considerably lower mortalities for sepsis (as defined by Sepsis-2), at 4%.

It is likely that the sepsis outcomes we describe for sSA are worse than in high-income settings. The reasons for this are likely to be multifactorial, including lack of resources. The association of HIV status with mortality highlights the likely role of HIV as a driver of poor sepsis outcomes in sSA. It is also possible that delays in presentation to hospital or delays in processes of care may contribute to high mortality. The majority of studies we have identified provide no data to address these

hypotheses, though there is some suggestion that presentation to care may be delayed. In the two Zambian RCTs[33,35], 66-74% patients with severe sepsis were unable to walk on admission, and this had been the case for a median of 5-16 days; in Uganda, patients with severe sepsis had been unwell for a median on 14 days before arrival at hospital[24]. Data on processes of care are also largely lacking, though delays are apparent in one study: in the “before” arm of the before-after intervention trial in Uganda[5, 34], 49% (117/245) of patients with severe sepsis received no antimicrobials within 6 hours of presentation, and a median of 500ml of intravenous fluids was delivered in 6 hours. However, in the usual-care arm of the two Zambian RCTs identified, mortality was high despite a prompt initiation of care: median door-to-antibiotic time was 1.3-1.5 hours and a median 1.3-2.0L of intravenous fluid was administered in the first 6 hours of hospital admission. Lack of critical care facilities may play a role; again, data are largely absent but across the two RCTs only 1% (3/318) of patients were cared for on an ICU.

Of note, the 30-day mortality of severe sepsis seemed to be higher than the in-hospital mortality, though with overlapping confidence intervals. The reasons for this are not addressed in our study, but could represent uncontrolled primary infection, secondary infection following sepsis-related immunosuppression, non-infection related mortality, or hospital practices such as palliative discharge of patients with poor prognosis. None of the studies we identified characterised longer-term (post 30-day) sepsis outcomes in sSA, and this highlights a major area for future research in resource-limited settings. Longer term follow-up of patients in resource-limited settings is logistically difficult, as is matching episodes of care in administrative or clinical databases for individuals across fragmented and under-resourced health

systems, which would be necessary to characterise sepsis outcomes from such databases; these factors may contribute to the lack of available data. No identified study characterised morbidity or economic cost either to health systems or individuals, but the combination of poor outcomes in a young adult population – often the most economically productive group any society - may have significant socio-economic implications.

In contrast to high-income settings, the populations recruited to the identified studies are young, and predominantly HIV infected. It is likely that the high HIV prevalence influences the causative pathogens: the commonest cause of bloodstream infection, when sought, was *M. tuberculosis*, and Nontyphoidal Salmonellae and *S. pneumoniae* were also commonly isolated, both of which have a strong association with HIV[37,38]. Management of HIV in the critically ill in LMICs is poorly described; this review highlights that a carefully considered strategy for HIV must be developed, and that data to inform it are required. The high prevalence of *S. aureus* is perhaps surprising, given data on incidence of causative agents of BSI in sSA[39]. Though this could represent true BSI, the possibility of poor collection technique or laboratory misidentification should be considered; though it was possible to exclude coagulase negative *Staphylococci* from our analysis as likely contaminants, determining whether *S. aureus* bacteraemia represented true BSI was not possible with the available data. Despite the fact that malaria diagnostic data is from studies carried out in malaria-endemic countries - Malawi, Uganda, Tanzania and Zambia - malaria was less common than bacterial bloodstream infection, highlighting the emerging importance of non-malarial aetiology of fever in malaria-endemic areas. In the single study employing PCR to describe sepsis aetiology beyond bloodstream infection,

CMV was extremely commonly isolated. While it may cause illness in the immunocompromised, CMV reactivation in critical illness is recognised in high income settings. Without viral testing in a non-sepsis control group, the causal role of CMV is therefore not clear[40]. Bacterial zoonoses were rare, in contrast to sparse data from fever aetiology studies from elsewhere in sSA[18], though no study carried out reference-standard serological assays. Data on antimicrobial resistance (AMR) patterns were lacking: a single study with small numbers of patients, found that Gram-negative third-generation cephalosporins resistance (3GC-R) was significantly associated with mortality. This, alongside data from across the continent suggesting that 3GC-R is an emerging problem[39,41,42], highlights the need for high quality national and sub-national surveillance for AMR across sSA to guide locally appropriate therapies.

The high prevalence of tuberculosis as a cause of sepsis has significant implications for appropriate antimicrobial therapy in sSA. Current sepsis protocols in high-income settings are based on rapid administration of broad-spectrum antimicrobials to cover common Gram-positive and Gram-negative pathogens, rather than Mycobacteria. However, it is far from clear how tuberculosis therapy should be used in this patient population in sSA, and due to lack of data it is not possible to make any recommendations from the studies we identified. Further studies are needed to guide clinicians in the best use of tuberculosis diagnostic tests or empiric therapy in sepsis, as are pharmacokinetic studies of tuberculosis therapy in the critically unwell.

We identified relatively few studies. Existing studies were geographically restricted: Uganda contributed 8/15 studies, for example, and Zambia 3/15. Several

institutions – largely tertiary referral centres – contributed multiple studies. Most studies excluded surgical or obstetric patients. The studies largely used a Sepsis-2 definition of sepsis and severe sepsis, though with a number of modifications, making between study comparisons difficult. Development of easily applicable sepsis definitions for low resource settings – or pragmatic modification of currently used screening scores such as qSOFA – could significantly increase the generalisability and utility of the evidence base in sSA.

There are weaknesses to our review, including that our searches may have missed studies. We excluded studies in preselected populations (in particular obstetric or surgical sepsis) to maximise the generalisability of our findings and minimise potential bias. These populations were also excluded *a priori* by identified studies, leading to lack of generalisability to surgical or pregnant patients. Much of sSA is unrepresented, potentially leading to bias from geographical restriction. Similarly, the results may not generalise to sepsis in district hospitals, or primary health care facilities, given the predominance of studies from tertiary facilities. The heterogeneity in mortality estimates means that the summary estimates should be regarded with caution. As with all meta-regression, there is the possibility of confounding or ecologic bias. Where it was not possible to disaggregate data from identified studies, we did not attempt to access individual participant level data. Assessing risk of bias for observational studies is difficult, with no recognised gold-standard. We chose to use the Newcastle-Ottawa score, a frequently used tool, but one which as recognised problems with reliability and validity[43,44] and as such our estimates of bias are likely to be under-estimates.

CONCLUSIONS

We have demonstrated that sepsis in sSA is dominated by the high prevalence of advanced HIV, subjecting a younger population to a high risk of sepsis of different microbial aetiology to high-income/low-HIV settings. Short term outcomes are poor despite the younger age of sepsis patients compared to high-income settings, whilst long term outcome data and morbidity data are absent. Data on sepsis aetiology beyond bloodstream infection and malaria are lacking though tuberculosis clearly plays an important role, which has implications for selection of appropriate antimicrobial therapy, though further studies are needed to develop an evidence base for the treatment of tuberculosis in the critically ill. Development and deployment of easily applicable sepsis definitions would help improve our understanding of the burden of sepsis in sSA and intervention studies aimed at the distinct population in sSA are urgently needed.

LIST OF ABBREVIATIONS

AMR: Antimicrobial Resistance

BSI: Bloodstream Infection

CD4: Cluster of Differentiation 4

HIV: Human Immunodeficiency Virus

PCR: Polymerase Chain Reaction

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses

qSOFA: Quick Sequential Organ Failure Assessment

SIRS: Systemic Inflammatory Response Score

SOFA: Sequential Organ Failure Assessment

sSA: Sub-Saharan Africa

DECLARATIONS

Ethics approval and consent to participate: not required.

Consent for publication: Not applicable.

Availability of data and materials: Not applicable.

Competing interests: The authors declare that they have no competing interests.

Funding: This article was funded by a DFID/MRC/Wellcome Joint Global Health Trials (MR/P020577/1). Additional funding was received by the National Institute for Health Research using Official Development Assistance (ODA) funding (the African Research Collaboration on Sepsis, 17/63/42). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

JL is supported by a Wellcome Trust Clinical PhD Fellowship (109105Z/15/A).

The funding bodies had no role in design of the study or collection, analysis, interpretation of data or in writing the manuscript.

Authors' contributions: All authors conceived and designed the study, JL and JR performed abstract screening and data extraction, JL analysed the data and drafted the manuscript. All authors revised and approved the final manuscript.

Acknowledgements: The study authors would like to thank the staff of the library at the Liverpool School of Tropical Medicine, particularly Cath Booth, for their assistance in locating papers for review.

FIGURES




Figure 1: Summary of studies screened and included.




A**B****C**

Figure 2: A: In-hospital sepsis mortality; B: In-hospital severe sepsis mortality C: 28- or 30- day severe sepsis mortality.

Figure 2 legend: In all cases only studies using a Sepsis-2 definition of sepsis or suspected sepsis are included. Random-effect summary estimate from generalized linear mixed model shown as well as measures of heterogeneity: τ^2 , I^2 and p value from Cochran's Q-test.







Figure 3: Meta regression showing effect of varying proportion of HIV-infected participants on inpatient mortality for sepsis and severe sepsis.

Figure 3 legend: 95% confidence intervals generated from 1000 bootstrap replicates. Likelihood ratio testing of nested models shows that including proportion of HIV-infected participants significantly improved model fit for sepsis ($p = 0.008$), but not severe sepsis ($p = 0.83$). Size of circle is proportional to number of participants with sepsis or severe sepsis.

A**B****C****Figure 4:** Pooled prevalence of A: Malaria, B: Bloodstream infection, C: *M.*

tuberculosis bloodstream infection.

Figure 4 legend: Random-effect summary estimate from generalized linear mixed model shown as well as measures of heterogeneity: τ^2 , I^2 and p value from Cochran's Q-test.

REFERENCES

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:801.
2. Fleischmann C, Scherag A, Adhikari NKJ, Hartog CS, Tsaganos T, Schlattmann P, et al. Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *Am J Respir Crit Care Med*. 2016;193:259–72.
3. Gaieski DF, Edwards JM, Kallan MJ, Carr BG. Benchmarking the Incidence and Mortality of Severe Sepsis in the United States*. *Crit Care Med*. 2013;41:1167–74.
4. Kaukonen K-M, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000-2012. *JAMA*. American Medical Association; 2014;311:1308–16.
5. Jacob ST, Moore CC, Banura P, Pinkerton R, Meya D, Opendi P, et al. Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS One*. 2009/11/13. 2009;4:e7782.
6. Maitland K, Kiguli S, Opoka RO, Engoru C, Olupot-Olupot P, Akech SO, et al. Mortality after fluid bolus in African children with severe infection. *N Engl J Med*. 2011;364:2483–95.

7. Andrews B, Semler MW, Muchemwa L, Kelly P, Lakhi S, Heimburger DC, et al. Effect of an Early Resuscitation Protocol on In-hospital Mortality Among Adults With Sepsis and Hypotension. *JAMA*. 2017;318:1233.
8. Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al. Assessment of Clinical Criteria for Sepsis. *JAMA*. American Medical Association; 2016;315:762.
9. Rudd KE, Seymour CW, Aluisio AR, Augustin ME, Bagenda DS, Beane A, et al. Association of the Quick Sequential (Sepsis-Related) Organ Failure Assessment (qSOFA) Score With Excess Hospital Mortality in Adults With Suspected Infection in Low- and Middle-Income Countries. *JAMA*. 2018;319:2202.
10. Adhikari NKJ, Rubenfeld GD. qSOFA Score for Patients With Sepsis in Low- and Middle-Income Countries. *JAMA*. 2018;319:2175.
11. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med*. 2003;31:1250–6.
12. Kaukonen K-M, Bailey M, Pilcher D, Cooper DJ, Bellomo R. Systemic Inflammatory Response Syndrome Criteria in Defining Severe Sepsis. *N Engl J Med*. 2015;372:1629–38.
13. Kumar A, Roberts D, Wood KE, Light B, Parrillo JE, Sharma S, et al. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med*. 2006;34:1589–96.
14. Seymour CW, Gesten F, Prescott HC, Friedrich ME, Iwashyna TJ, Phillips GS, et al. Time to Treatment and Mortality during Mandated Emergency Care for Sepsis. *N Engl J Med*. 2017;376:2235–44.

15. Muchemwa L, Shabir L, Andrews B, Bwalya M. High prevalence of *Mycobacterium tuberculosis* bacteraemia among a cohort of HIV-infected patients with severe sepsis in Lusaka, Zambia. *Int J STD AIDS.* 2016/03/24. 2016;
16. Jacob ST, Pavlinac PB, Nakiyingi L, Banura P, Baeten JM, Morgan K, et al. *Mycobacterium tuberculosis* bacteremia in a cohort of hiv-infected patients hospitalized with severe sepsis in uganda-high frequency, low clinical suspicion [corrected] and derivation of a clinical prediction score. *PLoS One.* 2013/08/14. 2013;8:e70305.
17. Crump JA, Ramadhani HO, Morrissey AB, Saganda W, Mwako MS, Yang LY, et al. Invasive bacterial and fungal infections among hospitalized HIV-infected and HIV-uninfected adults and adolescents in northern Tanzania. *Clin Infect Dis.* 2011/01/11. 2011;52:341–8.
18. Crump JA, Morrissey AB, Nicholson WL, Massung RF, Stoddard RA, Galloway RL, et al. Etiology of Severe Non-malaria Febrile Illness in Northern Tanzania: A Prospective Cohort Study. *PLoS Negl Trop Dis.* 2013;7:e2324.
19. Rubach MP, Maro VP, Bartlett JA, Crump JA. Etiologies of illness among patients meeting integrated management of adolescent and adult illness district clinician manual criteria for severe infections in northern Tanzania: implications for empiric antimicrobial therapy. *Am J Trop Med Hyg.* 2015;92:454–62.
20. Huson MAM, Kalkman R, Stolp SM, Janssen S, Alabi AS, Beyeme JO, et al. The impact of HIV on presentation and outcome of bacterial sepsis and other causes of acute febrile illness in Gabon. *Infection.* 2015;43:443–51.
21. Chimese SM, Andrews B, Lakhi S. The Etiology And Outcome Of Adult Patients Presenting With Sepsis To The University Teaching Hospital, Lusaka, Zambia. *Med*

J Zambia. 2012;39:19–22.

22. Winters BD, Eberlein M, Leung J, Needham DM, Pronovost PJ, Sevransky JE. Long-term mortality and quality of life in sepsis: A systematic review*. Crit Care Med. 2010;38:1276–83.
23. Moore CC, Jacob ST, Banura P, Zhang J, Stroup S, Boulware DR, et al. Etiology of Sepsis in Uganda using a Quantitative PCR-based TaqMan Array Card. Clin Infect Dis. 2019;68:266–72.
24. Amir A, Saulters KJ, Olum S, Pitts K, Parsons A, Churchill C, et al. Outcomes of patients with severe sepsis after the first 6 hours of resuscitation at a regional referral hospital in Uganda. J Crit Care. 2016;33:78–83.
25. Nadjm B, Mtobe G, Amos B, Walker NF, Diefendal H, Reyburn H, et al. Severe febrile illness in adult hospital admissions in Tanzania: a prospective study in an area of high malaria transmission. Trans R Soc Trop Med Hyg. 2012;106:688–95.
26. Waitt PI, Mukaka M, Goodson P, SimuKonda FD, Waitt CJ, Feasey N, et al. Sepsis carries a high mortality among hospitalised adults in Malawi in the era of antiretroviral therapy scale-up: a longitudinal cohort study. J Infect. 2015;70:11–9.
27. Ssekitoleko R, Jacob ST, Banura P, Pinkerton R, Meya DB, Reynolds SJ, et al. Hypoglycemia at admission is associated with inhospital mortality in Ugandan patients with severe sepsis. Crit Care Med. 2011;39:2271–6.
28. Ssekitoleko R, Pinkerton R, Muhindo R, Bhagani S, Moore CC. Aggregate evaluable organ dysfunction predicts in-hospital mortality from sepsis in Uganda. Am J Trop Med Hyg. 2011;85:697–702.
29. Auma MA, Siedner MJ, Nyehangane D, Nalusaji A, Nakaye M, Mwanga-Amumpaire J, et al. Malaria is an uncommon cause of adult sepsis in south-western

- Uganda. Malar J. 2013;12:146.
30. Seboxa T, Amogne W, Abebe W, Tsegaye T, Azazh A, Hailu W, et al. High Mortality from Blood Stream Infection in Addis Ababa, Ethiopia, Is Due to Antimicrobial Resistance. PLoS One. 2015;10:e0144944.
31. Rudd KE, Tutaryebwa LK, West TE. Presentation, management, and outcomes of sepsis in adults and children admitted to a rural Ugandan hospital: A prospective observational cohort study. PLoS One. 2017;12:e0171422.
32. Andrews B, Muchemwa L, Kelly P, Lakhi S, Heimbürger DC, Bernard GR. Simplified severe sepsis protocol: a randomized controlled trial of modified early goal-directed therapy in Zambia. Crit Care Med. 2014;42:2315–24.
33. Andrews B, Semler MW, Muchemwa L, Kelly P, Lakhi S, Heimbürger DC, et al. Effect of an Early Resuscitation Protocol on In-hospital Mortality Among Adults With Sepsis and Hypotension: A Randomized Clinical Trial. Jama. 2017/10/04. 2017;318:1233–40.
34. Jacob ST, Banura P, Baeten JM, Moore CC, Meya D, Nakiyingi L, et al. The impact of early monitored management on survival in hospitalized adult Ugandan patients with severe sepsis: a prospective intervention study*. Crit Care Med. 2012;40:2050–8.
35. Andrews B, Muchemwa L, Kelly P, Lakhi S, Heimbürger DC, Bernard GR. Simplified Severe Sepsis Protocol. Crit Care Med. NIH Public Access; 2014;42:2315–24.
36. Andrews B, Muchemwa L, Kelly P, Lakhi S, Heimbürger DC, Bernard GR. Simplified severe sepsis protocol: a randomized controlled trial of modified early goal-directed therapy in Zambia. Crit Care Med. 2014;42:2315–24.

37. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-typhoidal salmonella disease: an emerging and neglected tropical disease in Africa. *Lancet* (London, England). 2012;379:2489–99.
38. French N, Gordon SB, Mwalukomo T, White SA, Mwafulirwa G, Longwe H, et al. A trial of a 7-valent pneumococcal conjugate vaccine in HIV-infected adults. *N Engl J Med.* 2010;362:812–22.
39. Musicha P, Cornick JE, Bar-Zeev N, French N, Masesa C, Denis B, et al. Trends in antimicrobial resistance in bloodstream infection isolates at a large urban hospital in Malawi (1998–2016): a surveillance study. *Lancet Infect Dis.* 2017;17:1042–52.
40. Osawa R, Singh N. Cytomegalovirus infection in critically ill patients: a systematic review. *Crit Care.* 2009;13:R68.
41. Lai PS, Bebell LM, Meney C, Valeri L, White MC. Epidemiology of antibiotic-resistant wound infections from six countries in Africa. *BMJ Glob Heal. BMJ Specialist Journals;* 2018;2:e000475.
42. Ntirenganya C, Manzi O, Muvunyi CM, Ogbuagu O. High prevalence of antimicrobial resistance among common bacterial isolates in a tertiary healthcare facility in Rwanda. *Am J Trop Med Hyg.* 2015;92:865–70.
43. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010;25:603–5.
44. Hartling L, Milne A, Hamm MP, Vandermeer B, Ansari M, Tsertsvadze A, et al. Testing the Newcastle Ottawa Scale showed low reliability between individual reviewers. *J Clin Epidemiol.* 2013;66:982–93.

ADDITIONAL FILES

MODIFIED NEWCASTLE - OTTAWA QUALITY ASSESSMENT SCALE COHORT STUDIES

Note: A study can be awarded a maximum of one star for each numbered item within the Selection and Outcome categories. A maximum of two stars can be given for Comparability

Selection

1) Representativeness of the exposed cohort

- a) truly representative of the average adult with sepsis in the community *
- b) somewhat representative of the average adult with sepsis in the community *
- c) selected group of users eg nurses, volunteers
- d) no description of the derivation of the cohort

2) Ascertainment of exposure – HIV status

- a) reproducible description of valid testing procedures (e.g. following national guidelines)*
- b) self reported status
- c) no description

3) Ascertainment of exposure - aetiology

- a) Reproducible description of testing procedures using reference tests *
- b) Reproducible description of testing using non-reference tests (e.g Widal test for enteric fever)
- d) no or nonreproducible description of testing

Comparability

1) Comparability of cohorts on the basis of the design or analysis

- a) study reports +/- controls for lactate *
- b) study reports +/- controls for any additional physiologic parameters (BP, HR etc) *

Outcome

1) Assessment of outcome

Additional File 1: Modified Newcastle-Ottawa Quality Assessment Scale.

	Selection – representativeness	Selection – exposure (HIV)	Selection – exposure (aetiology)	Comparability	Outcome – assessment	Outcome – follow up
Waitt 2015	1	1	1	0.5	1	1
Ssekitoleko 2011 (2)	1	0	1	1	1	1
Ssekitoleko 2011 (1)	1	0	1	0	1	1
Seboxa 2015	1	0	1	0	1	1
Rudd 2017	1	0	1	1	1	1
Nadjm 2012	0	1	1	1	1	1
Jacob 2012	1	1	1	1	1	1
Jacob 2009	1	1	1	1	1	1
Huson 2015	1	1	1	0	1	1
Chimese 2012	1	0	1	1	1	1
Auma 2013	1	1	1	0.5	1	1
Andrews 2017	1	1	1	1	1	1
Andrews 2014	1	0	1	1	1	1
Amir 2016	1	1	1	1	1	1

Additional File 2: Results of quality assessment using modified Newcastle-Ottawa scale.

Additional File 2 Legend: Numbers are a normalised score for a given domain between 0 (low quality) and 1 (high quality).

Study	Inclusion criteria	Exclusion criteria
Jacob 2009	(1) 18+ years in daytime hours, susp. infection (2) 2 or more of axillary T > 37.5C or < 35.5C, HR > 90, RR > 20 (3) SBP < 100	acute CVA, GI haemorrhage, required admission to nonmedical ward, unable to consent and no guardian
Nadjm 2012	(1) "adult" wed-sun, (2) fever or history of fever, (3) any one of confusion/coma, deep breathing, severe tachypnoea or hypoxia, severe pallor, visible jaundice, unable to walk, diagnosis of "severe malaria"	none
Jacob 2012 and Moore 2018	(1) 18+ years in daytime hours, susp. infection (2) 2 or more of axillary T > 37.5C or < 35.5C, HR > 90bpm, RR > 20bpm (3)	suspected surgical or obstetric emergencies or shock without suspected infection

	SBP < 100mmHg, (4) lactate > 2.5mmol/L or KPS <= 40, (5) available guardian	
Waitt 2015	(1)16+ years Mon-Fri, susp infection, (2) 2 or more of T > 38.3C or < 35C, RR > 20bpm, HR > 90bpm, WCC > 12,000/ μ L or < 4,000/ μ L, altered mental state	hospitalised or received antibiotics in last 2 weeks, unable to consent (ie obtunded and no guardian)
Ssekitoleko 2011 (1)	(1) 18+ yrs , suspected infection, (2) 2 or more of axillary T of > 37.5°C or <35.5°C, HR > 90, RR> 20, (3) SBP < 100mmHg, (4) one of lactate of >4 mmol/L, platelet number of <100,000/ μ L, or altered mental state	acute cerebrovascular event or gastrointestinal hemorrhage or needed triage to a surgery or obstetrics and gynecology ward
Ssekitoleko 2011 (2)	18+ yrs Mon-Fri; (2) 18year+; (3) at least 2 of T < 36C or > 38C, RR > 20bpm, HR > 90bpm	Required triage to surgical or obs&gynae ward, had received any treatment or IV fluids before recruitment or suspected CVA or gastrointestinal haemorrhage
Chimese 2012	1) 16+ years, susp. infection 2) 2 of T > 38C or < 36C, HR > 90bpm, RR > 20bpm, WCC > 12,000/ μ L or < 4,000/ μ L	none
Andrews 2014	(1) 18+ year Mon-Fri, susp. infection, 2) 2+ of HR> 90bpm, RR > 20bpm, T >=38.0C or <=36.0C, or WBC > 12,000/ μ L or < 4,000/ μ L, and 3) One of SBP < 90 mmHg or MAP<65mmHg, altered mentation, creatinine > 1.85 mg/dL, platelet count<100,000/ μ L, resp>40, jaundice	gastrointestinal bleed, required immediate surgery, congestive heart failure exacerbation or end-stage renal disease or raised jugular venous pressure (JVP) > 3 cm above the sternal angle, as measured with a level and a ruler.
Auma 2013	(1) 18+ medical admission 08:00-00:00 susp. infection (2) 2+ of axillary T>37.5 or T<35.5, HR>90bpm, RR>20bpm	surgical, o&g admission
Andrews 2017	(1) 18+ year Mon-Fri, susp. infection, 2) 2+ of HR> 90bpm, RR > 20bpm, T >=38.0C or <=36.0C, or WBC > 12,000/ μ L or < 4,000/ μ L, and 3) SBP < 90 mmHg or MAP<65mmHg	RR > 40 or SpO2 < 90%, gastrointestinal bleed, required immediate surgery, congestive heart failure exacerbation or end-stage renal disease or raised jugular venous pressure (JVP) > 3 cm above the sternal angle, as measured with a level and a ruler.
Huson 2014	1) 18+ years, clinical evidence of infection, weekdays 2) tympanic T >= 38C or < 36C, 3) one of HR > 90bpm, RR > 20bpm, WBC > 12,000/ μ L or < 4,000/ μ L	none


Seboxa 2015	1) "Adult" 2) 2+ of axillary T > 38.5 or < 36.5, HR > 90bpm, RR > 20bpm	Received antimicrobials within last 72hr
Rudd 2017	1) 18+ years 2) 2+ of T > 38 or < 36, HR > 90bpm, RR > 20bpm or PCO ₂ < 32mmHg, WBC > 12,000/ μ L or < 4,000/ μ L or > 10% band forms	Surgical patients, admission to maternity ward
Amir 2016	(1) 14+ years, susp. infection; (2) 2+ of axillary T >= 38°C or < 36 36°C, HR > 90 bpm, RR > 20bpm, or WBC > 12,000/ μ L or < 4,000/ μ L, (3) signs of end-organ dysfunction including SBP <= 90 platelet count < 100,000/ μ L or GCS < 15	Triage to a surgical or o&g ward, had received any antibiotics or intravenous fluids prior to recruitment, or had a history suggestive of other diagnoses associated with lactic acidosis such as diabetic ketoacidosis, acute coronary syndrome, or chronic liver disease

Additional File 3: Inclusion and exclusion criteria for included studies.

Study	Total number of participants	Available aetiology data		
		Blood culture	Mycobacterial blood culture	Malaria
Jacob 2009	382	382/382 (100%)	249/382* (65%)	382/382 (100%)
Nadjm 2012	198	198/198 (100%)	ND	188/198 (95%)
Jacob 2012	426	426/426 (100%)	426/426 (100%)	426/426 (100%)
Waitt 2015	213	213/213 (100%)	ND	213/213 (100%)
Ssekitoleko 2011 (1)	96	NR	NR	NR
Ssekitoleko 2011 (2)	150	150/150 (100%)	ND	150/150 (100%)
Chimese 2012	161	161/161 (100%)	ND	ND
Andrews 2014	112	103/112 (92%)	82/112* (73%)	109/112 (92%)
Auma 2013	216	141/216 (65%)	ND	216/216 (100%)
Andrews 2017	209	209/209 (100%)	187/209* (89%)	47/209 (22%)
Huson 2014	107	107/107 (100%)	NR	NR
Seboxa 2015	292	292/292 (100%)	ND	ND
Rudd 2017	20	ND	ND	20/20 (100%)
Amir 2016	218	ND	ND	ND

Additional File 4: Availability of diagnostic testing by study.

Additional File 4 Legend: * = mycobacterial blood cultures were carried out at one study site only; ** = mycobacterial blood cultures were carried out in HIV positive participants only. One study (Moore 2018) which carried out retrospective testing of stored samples from Jacob 2012 is not included in this table, but is described in the text.



Additional File 5: Species of pathogenic bacteria isolated from aerobic blood culture.

Additional File 5 Legend: NTS = Non-Typhoidal Salmonellae