

RESEARCH

Open Access



Efficacy and safety of dihydroartemisinin–piperavaquine for treatment of *Plasmodium falciparum* uncomplicated malaria in adult patients on antiretroviral therapy in Malawi and Mozambique: an open label non-randomized interventional trial

Esperança Sevene^{1,2*†}, Clifford G. Banda^{3,4†}, Mavuto Mukaka^{3,5,6}, Sonia Maculuve¹, Salésio Macuacua¹, Anifa Vala¹, Mireia Piqueras⁷, Linda Kalilani-Phiri³, Jane Mallewa^{3,4}, Dianne J. Terlouw^{3,4,8}, Saye H. Khoo^{9,10}, David G. Lalloo^{8,10} and Victor Mwapasa^{3,4*}

Abstract

Background: HIV-infected individuals on antiretroviral therapy (ART) require treatment with artemisinin-based combination therapy (ACT) when infected with malaria. Dihydroartemisinin–piperavaquine (DPQ) is recommended for treatment of *Plasmodium falciparum* malaria, but its efficacy and safety has not been evaluated in HIV-infected individuals on ART, among whom drug–drug interactions are expected. Day-42 adequate clinical and parasitological response (ACPR) and incidence of adverse events were assessed in HIV-infected individuals on non-nucleoside reverse transcriptase inhibitor-based ART (efavirenz and nevirapine) with uncomplicated *P. falciparum* malaria treated with dihydroartemisinin–piperavaquine.

Methods: An open label single arm clinical trial was conducted in Malawi (Blantyre and Chikhwawa districts) and Mozambique (Manhiça district) involving patients aged 15–65 years with uncomplicated *P. falciparum* malaria who were on efavirenz-based or nevirapine-based ART. They received a directly-observed 3-day standard treatment of DPQ and were followed up until day 63 for malaria infection and adverse events. Day-42 PCR-corrected-ACPRs (95% confidence interval [CI]) were calculated for the intention-to-treat (ITT) population.

Results: The study enrolled 160 and 61 patients on efavirenz and nevirapine-based ART, with a baseline geometric mean (95% CI) parasite density of 2681 (1964–3661) and 9819 (6606–14,593) parasites/ μ L, respectively. The day-42 PCR-corrected ACPR (95% CI) was 99.4% (95.6–99.9%) in the efavirenz group and 100% in the nevirapine group. Serious adverse events occurred in 5.0% (8/160) and 3.3% (2/61) of the participants in the efavirenz and nevirapine group, respectively, but none were definitively attributable to DPQ. Cases of prolonged QT interval (> 60 ms from baseline) occurred in 31.2% (48/154) and 13.3% (8/60) of the patients on the efavirenz and nevirapine ART groups, respectively.

*Correspondence: esevene68@gmail.com; vmwapasa69@gmail.com

[†]Esperança Sevene and Clifford G. Banda contributed equally to this manuscript

¹ Centro de Investigação em Saúde de Manhiça (CISM), Maputo, Mozambique

³ University of Malawi, College of Medicine, Blantyre, Malawi

Full list of author information is available at the end of the article



These were not clinically significant and resolved spontaneously over time. As this study was not designed to compare the efficacy and safety of DPQ in the two ART groups, no formal statistical comparisons were made between the two ART groups.

Conclusions: DPQ was highly efficacious and safe for the treatment of malaria in HIV-infected patients concurrently taking efavirenz- or nevirapine-based ART, despite known pharmacokinetic interactions between dihydroartemisinin-piperazine and efavirenz- or nevirapine-based ART regimens.

Trial registration Pan African Clinical Trials Registry (PACTR): PACTR201311000659400. Registered on 4 October 2013, <https://pactr.samrc.ac.za/Search.aspx>

Keywords: Human immunodeficiency virus, Antiretroviral drugs, Dihydroartemisinin-piperazine, Malaria, Drug-drug interactions

Background

Malaria and human immunodeficiency virus (HIV) infections co-exist in most parts of sub-Saharan Africa [1]. Antiretroviral treatment (ART) naïve HIV-infected individuals are more susceptible to *Plasmodium falciparum* malaria infection than the HIV-uninfected population [2, 3]. The World Health Organization (WHO) recommends the use of artemisinin-based combination therapy (ACT) to treat *P. falciparum* malaria infections [4]. Dihydroartemisinin-piperazine (DPQ) is one of the most commonly used WHO-recommended ACT, but there are few data about its efficacy and safety in those taking ART despite the fact that large numbers of HIV-infected patients on ART are likely to be treated with DPQ for malaria.

Most ART regimens in sub-Saharan Africa contain non-nucleoside reverse transcriptase inhibitors, efavirenz and nevirapine. These drugs are metabolized by the cytochrome (CYP) P450 enzymes, particularly CYP3A4 and CYP2B6, which also metabolize artemisinin-derivatives and piperazine [5–7]. Thus, co-administration of ART and DPQ may result in drug–drug interactions [5]. Indeed, a recent pharmacokinetic study found that in malaria uninfected HIV-infected individuals on efavirenz-based ART, piperazine concentrations were 43% lower than ART naïve controls [8]. This drug–drug interaction may compromise DPQ's efficacy. In addition, malaria uninfected HIV-infected individuals on nevirapine-based ART treated with DPQ had higher piperazine concentrations than ART naïve controls who received DPQ only [8]. This drug–drug interaction may increase piperazine-related adverse events.

In view of the limited data on the efficacy and safety of DPQ in HIV-malaria co-infected patients, a single arm clinical trial was conducted to estimate the efficacy of DPQ when used to treat parasitologically-confirmed uncomplicated clinical *P. falciparum* malaria in HIV-infected people on standard ART regimens (efavirenz- or nevirapine-based ART). Specifically, the trial assessed the day-42 PCR-corrected adequate clinical and

parasitological response [ACPR] to examine whether it exceeds 90%, the WHO recommended benchmark for an efficacious anti-malarial drugs [9]. In addition, the trial assessed the safety of DPQ by determining the incidence of adverse events.

Methods

Study sites and study population

This study was part of a multi-country single arm clinical trial aimed at assessing the efficacy and safety of two artemisinin-based combinations (DPQ and artemether-lumefantrine) when used to treat malaria in HIV-infected adults on standard ART. One of the single arm trial assessing the efficacy and safety of AL was conducted in Zambia and findings have been reported elsewhere [10]. In this paper, findings from another single arm trial assessing the efficacy and safety of DPQ which was conducted from October 2013 to June 2015 at Queen Elizabeth Central Hospital and Chikhwawa District Hospital in Malawi as well as Manhica District Hospital in Mozambique, are reported. These are settings of moderate-high transmission of malaria [11, 12] and high HIV prevalence [13, 14]. Blantyre is an urban district in Southern Malawi with an estimated population 1,239,647, while Chikhwawa is a rural district located 34 km south of Blantyre with an estimated population of 518,284. In 2014, the malaria parasite prevalence in under-five children in Malawi was 33% and higher in rural (37%) than urban areas (11%) [11]. HIV prevalence in Malawi was estimated at 10.6% in 2015 [13]. Manhica is a rural district in Southern Mozambique located 80 km north of Maputo city, with an estimated population of 178,000 in 2014. Malaria parasite prevalence in under-five children was estimated at 51% in 2013 [12] while HIV prevalence in the district was estimated at 39.9% in 2010 [14].

During the study period, the criteria for initiating ART in the two countries were WHO HIV disease stages 3 or 4, CD4 cell count <350, pregnancy or lactation [15]. In October–December 2013, >87% of ART individuals in Malawi were on a fixed dose combination of tenofovir/

lamivudine/efavirenz while 6% were on stavudine/lamivudine/nevirapine or zidovudine/lamivudine/nevirapine [16]. In Mozambique, the majority of the ART individuals were on fixed dose zidovudine/lamivudine/nevirapine but the use of fixed dose tenofovir/lamivudine/efavirenz increased steadily over the study period. In both countries, artemether–lumefantrine was the first-line treatment for uncomplicated malaria. DPQ was registered but not routinely available within the public health system in the two countries. Nevertheless, it was evaluated in this study because it is one of the WHO recommended artemisinin-based combinations and has a more convenient dosing schedule than AL (once daily for 3 days). The efficacy and safety of AL was assessed in a separate trial [10].

Study design and clinical procedures

This was a single arm clinical trial (Registration number: PACTR201311000659400). HIV-infected patients on nevirapine- or efavirenz-based ART suspected of having malaria were pre-screened through history taking and clinical examination to determine their eligibility for the study. The study inclusion criteria were as follows: age ≥ 15 to ≤ 65 years; weight ≥ 35 kg; documented fever (axillary ≥ 37.5 °C) or history of fever 24 h prior enrolment; smear positive *P. falciparum* malaria monoinfection with asexual malaria parasite densities $< 200,000/\mu\text{L}$; ability to swallow oral medications and willingness and ability to comply with scheduled visits, supervised treatment administration, laboratory tests, and other study procedures. The following were the exclusion criteria: severe malaria as per WHO criteria [17]; mixed infection with another *Plasmodium* species; haemoglobin (Hb) concentration < 7 g/dL; severe sickle cell disease or sickle-haemoglobin C anaemia; evidence of pregnancy or lactation; use of any anti-malarial treatment or drug with anti-malarial activity within the past 1 month, except cotrimoxazole; history of DPQ hypersensitivity reactions; gastrointestinal diseases that could alter gut absorption or motility; history of splenectomy; history of epilepsy or convulsions; pre-existing clinically-significant cardiac, liver, renal, neurological or psychiatric abnormalities; alternative clinical cause of fever other than malaria and participation in any investigational drug study in the past 30 days.

Finger-prick blood samples were taken from those who satisfied the preliminary eligibility criteria and tested for malaria using Rapid Diagnostic Test (RDT) (SD BIO-LINE Malaria Ag P.f/Pan test produced by Alere) and for haemoglobin concentration using Hemocue Haemoglobinometer. Thick blood smear microscopy examinations were performed on patients with RDT positive malaria while clinical examinations were performed in those with confirmed malaria parasitaemia. Consenting

participants were enrolled and scheduled for a 3-day hospital admission.

The participants received dihydroartemisinin–piperaquine (Eurartesim[®], Sigma Tau): 3 tablets for study participants < 60 kg or 4 tablets study for participants ≥ 60 kg. Each tablet contained dihydroartemisinin/piperaquine 40 mg/320 mg, respectively, administered at 0 h, 24 (+ 4) and 48 (+ 4) h after the first dose. Participants' vital signs were measured at 6-hourly intervals and adverse events were monitored. A 12-Lead electrocardiogram (ECG) was performed before the first dose of DPQ and within 2 h after administration of the third dose DPQ. Any patient with Fridericia-corrected QT (QTcF) interval of ≥ 450 ms or QTc increase of > 60 ms from the baseline underwent follow-up ECGs until resolution of the abnormality. Participants were discharged at least 24 h after taking the third (last) dose of DPQ (post-treatment day 3) and advised to come for follow up visits on post-treatment days 7 (± 1), 14 (± 1), 21 (± 2), 28 (± 2), 35 (± 2), 42 (± 2) and 63 (± 2). Participants were encouraged to return to the health facility any time they felt unwell (unscheduled visits). All adverse events were graded using the DAIDS criteria [18]. Adverse events with onset or increased severity after the first dose of DPQ were counted as treatment-emergent adverse events (TEAEs). During follow up visits, participant's time and any incurred expenses when attending the study clinic were appropriately compensated, as approved by the ethics committees.

Laboratory procedures

During the admission period, thick blood slides were collected pre-dosing and at 6-hourly intervals until after obtaining two consecutive malaria negative smears. The slides were also collected at scheduled and unscheduled follow-up visits. The slides were Giemsa-stained and read by an experienced microscopist using standard protocols [19]. For quality control, all slides were re-read by a second microscopist; a third microscopist settled any discrepant readings. Dry blood spot (DBS) samples were collected on filter paper (Whatman 3MM[®]) at baseline and during recurrent malaria episodes. Parasite DNA was extracted from the DBS samples, amplified using polymerase chain reaction (PCR) and genotyped for merozoite specific protein (MSP) 1 and 2 to distinguish malaria recrudescence from re-infection, using methods previously described [20]. Samples that did not produce results were classified as indeterminate.

Venous blood samples were collected on days 0, 3, 28, 42 and 63 for biochemistry tests using a Beckman CX5[®] Chemistry analyzer, on days 0, 3, 7, 28, 42 and 63 for haematological tests using a Beckman Coulter[®] HMX Analyzer and on days 0, 28, and 63 for CD4 cell count

measurement using a BD FACSCount™ machine. Plasma samples collected on days 0, 28, and 63 were stored for future HIV viral load assays.

Blood samples for sparse pharmacokinetic (PK) assays were collected in sampling windows of 0–6, 6–48, 48–60 h, and on days 7, 21, 28 or 35 from first dose, as previously recommended [21]. In this paper, the relationship between day-7 concentrations and ACPR was explored since day-7 concentrations of the slowly eliminated partner drug of ACT have been shown to be a better determinant of therapeutic response than the area under the concentration–time curve [22]. The PK samples were analysed using a previously described HPLC–UV assay method [8]. The lower limit of quantitation (LLOQ) of the piperazine was 25 ng/mL, with a coefficient of variation of <10%. The PK laboratory at the Malawi-Liverpool Wellcome Trust Clinical Research Programme in Blantyre, Malawi, participated in the World Wide Antimalarial Resistance Network's external quality assurance programme [23].

Study endpoints

The primary study endpoint was proportion of patients with PCR-corrected day 42 ACPR, defined as patients who did not have parasitaemia on day 42 that exhibited identical *P. falciparum* malaria PCR markers (merozoite surface protein 1 and 2) with those at baseline, irrespective of axillary temperature, and who had not previously met any of the criteria of early treatment failure (ETF), late clinical failure (LCF) or late parasitological failure (LPF). Standard WHO definitions of ETF and LCF were used [9].

The other primary study end points were grade 3 or 4 TEAEs of special interest (Fridericia-corrected QT interval prolongation, dizziness, palpitations, urticaria or itchiness) and serious adverse events (SAEs) as per standard definitions [24]. Local study physicians determined the relationships between DPQ and the adverse events (AEs). A Data Safety and Monitoring Board reviewed serious AEs and adverse events of special interest (AESIs) and assessed the validity of study physicians' decisions. Secondary endpoints, included day 42 PCR-uncorrected ACPR, time for parasite to decline by 50% (PC₅₀) and 90% (PC₉₀), fever clearance time and trends in haemoglobin concentrations and CD4 cell counts from baseline to day 28.

Samples size

Sample size calculation was based on estimates of total treatment failure rate (TTFR). The estimated day-42 PCR corrected TTFR was ≤10% [25]. A precision of 5%, around this point estimate, allowed the upper limit of the 95% Wald binomial confidence interval to be 15%.

Using the formula for estimating a single study population sample size [26], our effective sample size was estimated at 138 for each ART type. The final sample size, for each ART group, was 163 after adjusting for an anticipated loss-to-follow-up rate of 15%. This sample size was achieved for the efavirenz-ART group but not for nevirapine-ART group, as the ART programs in both countries had successfully transitioned from nevirapine-to efavirenz-based ART first-line regimens. Sample size calculations including all subsequent statistical analysis were performed in STATA 13.1

Statistical analyses

For the primary end-point, three analysis populations were used. Firstly, the Intention-to-treat (ITT) population included patients who received at least 1 dose of study medication. Secondly, the per-protocol (PP) population included all participants who had the primary endpoint data at day 42, received a full course of DPQ and adhered to the follow-up visit schedule. Thirdly, the safety population included all patients who received any amount of study medication and had at least 1 assessment after dosing. ACPR plus 95% CI in PP and ITT populations were calculated. Sensitivity analyses were performed using the ITT and PP populations that first considered all participants with missing data as having parasitological failure and then considered the same participants as having treatment success.

Statistical analyses for secondary ACPR endpoints were similar to the primary endpoints. In addition, the Kaplan–Meier survival plots were used to summarize the time to PCR-corrected and uncorrected treatment failure. Parameters assessing post-treatment parasite clearance (PC₅₀, PC₉₀ and parasite clearance half-life) in the two ART groups were estimated using the WorldWide Antimalarial Resistance Network parasite clearance estimator, as described elsewhere [27].

Descriptive statistics were computed for baseline variables in the two ART groups. However, as this study was designed to estimate and not to compare efficacy and safety of DPQ between the two ART groups, no formal statistical comparisons of baseline characteristics efficacy or safety endpoints were made between the two ART groups. As part of a priori exploratory analysis, and where appropriate, Wilcoxon rank-sum/Mann–Whitney U test was used to compare distributions of the day-7 piperazine concentrations in those who attained or did not attain ACPR by day 42. Piperazine concentrations below the LLOQ were imputed to half the lower limit of quantification and included in the estimation of median piperazine exposure if the imputed values were <10% of the data. Additionally, Wilcoxon matched paired

signed-rank test was used to compare baseline and day 28 CD4 cell and haemoglobin values in each ART group.

Results

Study profile

A total of 1864 patients presenting at the health facilities with symptoms suggestive of malaria were screened for trial eligibility (Fig. 1). Two hundred and twenty-one patients with positive *P. falciparum* malaria blood met the eligibility criteria and were enrolled in the trial. None of the participants on nevirapine-ART and 5.6% (9/160) on efavirenz-ART were lost-to-follow-up or withdrew consent.

Baseline characteristics

The baseline characteristics of enrolled study participants are summarized in Table 1. Most of the

participants in the nevirapine-ART group (83.6%, N= 61) were from Mozambique while those on efavirenz-ART were predominantly from Malawi (75%, N= 160). Participants in the nevirapine-ART group were older and had been on ART for a longer period than those in the efavirenz-ART group. The geometric mean parasite density and median CD4 cell count were higher in the nevirapine-ART group than in the efavirenz-ART arm. A lower proportion of participants on nevirapine-ART were on cotrimoxazole prophylaxis compared with those on efavirenz-ART. There were no major differences between the two groups in body mass index and hemoglobin concentrations. The median Fridericia-corrected QT (QTcF) interval at recruitment was longer in the nevirapine-ART group than the efavirenz-ART group.




Fig. 1 Trial profile and flow chart of participants. Trial profile and flow chart of participants enrolled into the efavirenz (EFV) and nevirapine (NVP) antiretroviral groups in Malawi and Mozambique and receiving dihydroartemisinin–piperaquine (DPQ)

Table 1 Baseline characteristics of enrolled participants in Malawi and Mozambique, stratified by group of antiretroviral therapy

Variable	Efavirenz group (N = 160)	Nevirapine group (N = 61)
Age in years, median (IQR)	38.6 (32.3–44.7)	47.0 (37.9–51.8)
Female, %	107 (66.9)	46 (75.4)
Body mass index, kg/m ² , median (IQR)	20.8 (19.3–22.7)	21.1 (18.5–23.9)
WHO BMI classification, n (%)		
Underweight	26 (16.3)	15 (24.6)
Normal	123 (76.8)	35 (57.4)
Overweight	5 (3.1)	6 (9.8)
Obese	6 (3.8)	5 (8.2)
Duration on ART in months, median (IQR)	11 (5–32)	55 (33–70)
Presenting symptoms, n (%)		
Fever	146 (91.3)	57 (93.4)
Headache	144 (90.0)	57 (93.4)
Body aches	89 (55.6)	41 (67.2)
Nausea	40 (25.0)	33 (54.1)
Axillary temperature at enrolment, median (IQR)	37.7 (37.0–37.9)	37.1 (36.6–37.8)
Geometric Mean Parasite density, 95% CI.	2681 (1964–3661)	9819 (6606–14,593)
Parasite density > 2000 parasites/μL, n (%)	84 (52.5)	53 (86.9)
Pre-treatment hemoglobin concentration, g/dL, median (IQR)	12.2 (10.9–13.5)	11.9 (10.7–12.7)
Pre-treatment CD4 cell count, median (IQR) cells/mm ³	256 (140–360)	390 (237–500)
CD4 cell count < 350, n (%)	106 (66.3)	27 (44.3)
Median (IQR) QTcF interval (ms)	395 (360–418)	418 (389–438)
Reported use of any insecticide-treated bed net before presentation, % (n)	44 (27.5)	16 (26.2)
Current use of cotrimoxazole prophylaxis, % (n)	109 (68.1)	12 (19.6)

Treatment dosage and tolerability

The total median (range) dosages of dihydroartemisinin and piperazine administered to participants were 7.0 mg/kg (5.1–9.9) and 56.3 mg/kg (40.9–79.1), respectively in the efavirenz-ART group and 7.0 mg/kg (5.6–8.6) and 55.7 mg/kg (45.0–68.6), respectively in the nevirapine-ART group. The dosages were well tolerated: 3 participants in the efavirenz-ART group and none in the nevirapine-ART group vomited following intake of DPQ. Participants who vomited were re-dosed and none was withdrawn from the study due to persistent vomiting.

Treatment efficacy

As shown in Table 2, only one participant in the efavirenz-ART group, with baseline CD4 count of 26 cells/μL, had ETF on treatment day 2. Also, seven cases in the efavirenz-ART group had LPF. Parasite genotyping results were available in 6 of the 7 LPF cases which were all classified as re-infections. No treatment failures occurred in the nevirapine-ART group.

In the ITT analyses, the PCR-corrected day 42 ACPR was 99.4% (95% confidence interval [CI] 95.6–99.9%)

in the efavirenz-ART group and 100% in the nevirapine-ART group (see Additional file 1). The day 42 PCR-uncorrected ACPR was 95.6% (95% CI 90.3–97.5%) in the efavirenz-ART group and 100% in the nevirapine-ART group (see Additional file 2).

In the PP analyses, the day 42 PCR-corrected ACPR was 99.3% (95% CI 95.4–99.9%) in the efavirenz-ART group and 100% in the nevirapine-ART group. The day 42 PCR-uncorrected ACPR was 94.7% (95% CI 89.7–97.3%) in the efavirenz-ART group and 100% in the nevirapine-ART group.

In sensitivity analyses which considered missed visits or samples, unavailable PCR results and loss to follow up as treatment failures, the day-42 PCR-corrected ACPR was 93.1% (95% CI 88.0–96.2%) in the ITT population and 98.8% (95% CI 95.1–99.7%) in the PP population. Details of the different scenarios accounting for the missing results are shown in Table 2.

Parasite clearance time

Parasite clearance parameters were calculated in 46 and 84 participants in the nevirapine- and efavirenz-ART groups, respectively, who had detectable parasitemia at

Table 2 Efficacy outcomes and adequate clinical and parasitological response rates by day 42 among the enrolled participants stratified by group of antiretroviral therapy

Variable	DPQ + efavirenz N = 160	DPQ + nevirapine N = 61
Early treatment failure-no. (%)	1 (0.6)	0
Development of danger signs or severe malaria	0	0
Parasitaemia on day 2 greater than day 0	0	0
Parasitemia on day 3 \geq 25% of count on day 0	1 (0.6)	0
Parasitemia on day 3 with axillary temperature \geq 37.5 °C	0	0
Late clinical failure-no. (%)	0	0
Late parasitological failure-no. (%)	7 (4.4)	0
Recrudescence	0	0
Reinfection	6 (3.8)	0
Indeterminate or sample unavailable	1 (0.6)	0
Adequate clinical and parasitological response rates by different scenarios		
Intention-to-treat analysis		
PCR-corrected cure rate-% (95% CI)		
Scenario 1 ^a	99.4 (95.6–99.9)	100
Scenario 2 ^b	93.1 (88.0–96.2)	100
PCR-uncorrected cure rate-% (95% CI)		
Scenario 1 ^a	95.6 (90.3–97.5)	100
Scenario 2 ^b	89.4 (83.5–93.3)	100
Scenario 3 ^c	89.9 (82.3–94.4)	100
Scenario 4 ^d	89.8 (77.4–95.8)	100
Per-protocol analysis		
Number of patients		
Number of patients	151	61
PCR-corrected cure rate-% (95% CI)		
Scenario 5 ^e	99.3 (95.4–99.9)	100
Scenario 6 ^f	98.8 (95.1–99.7)	100
PCR-uncorrected cure rate-% (95% CI)		
Scenario 5 ^e	95.3 (90.5–97.8)	100
Scenario 6 ^f	94.7 (89.7–97.3)	100

^a Scenario 1: Indeterminate or unavailable PCR samples or loss to follow up by day 42 (n = 9, in efavirenz-ART group) in intention-to-treat population treated as treatment success

^b Scenario 2: Indeterminate or unavailable PCR samples or loss to follow up by day 42 (n = 9, in efavirenz-ART group) in the intention-to-treat population treated as treatment failures

^c Scenario 3: On cotrimoxazole prophylaxis in the intention-to treat analysis and combined with scenario 2 above

^d Scenario 4: Not on cotrimoxazole prophylaxis in the intention-to treat and combined with scenario 2 above

^e Scenario 5: Indeterminate or unavailable PCR samples in per-protocol population treated as treatment success


^f Scenario 6: Indeterminate or unavailable PCR samples in per-protocol population treated as treatment failure

two or more post-dosing time points. The median (range) PC₅₀ and the median (range) PC₉₀ in the ITT population were 3.1 (0.2–10.3) h and 8.2 (2.4–19.7) h, respectively, in the nevirapine-group and 4.2 (0.6–40.3) h and 10.1 (3.2–63.1) h, respectively, in the efavirenz-group. The median parasite clearance half-life (range) were 2.1 (1.1–6.8) and 2.2 (1.2–9.8) h in the nevirapine- and efavirenz-ART groups, respectively (Fig. 2). One participant (2.2% [95% CI 0.3–14.8]) in the nevirapine-ART group and five participants (6.0% [2.5–13.7]) in the efavirenz-ART group, had parasite clearance half-life of >5.5 h. All six

participants had a baseline parasite density of > 3500 parasites/ μ L and CD4 cell count of < 250 cells/mm³. None of these six participants experienced malaria recurrence during the follow up period.

Day-7 plasma piperazine concentrations

On day 7 of follow-up, blood samples for plasma piperazine quantification were available from 179 of the retained 216 participants. Of these, 62.0% (111/179) had day-7 piperazine concentrations that were below the LLOQ (< 25 ng/mL): 60.0% (81/135) in the efavirenz-ART



group and 68.2% (30/44) in the nevirapine-ART group. The piperazine values for these individuals were imputed to half the lower limit of quantification. However, since these values were more than >10% of the data, they were excluded in calculation of median piperazine exposure. Overall, the median (range) day-7 piperazine concentration in participants with concentrations >LLOQ was 57 (25.5–592.8) ng/mL in the efavirenz-ART group and 79.1 (25.4–211.1) ng/mL in the nevirapine-ART group. After excluding <LLOQ values, participants who experienced malaria recurrence by day 42 ($n=3$) had a median (range) day-7 piperazine concentration of 63.9 (49.7–78.9) ng/mL and this value was 66.7 (25.4–592.8) ng/mL in participants who did not experience malaria recurrence ($n=65$, $p=0.965$).

Fever clearance

At baseline, 58.1% (93/160) of participants in the efavirenz-ART group and 31.1% (19/61) in the nevirapine-ART group were febrile (axillary temperature ≥ 37.5 °C). The median fever clearance time (IQR) was 6 h (6–60) in the efavirenz-ART group and 6 h (6–12) in the nevirapine-ART group. By treatment day 2, only 3.8% (6/160) participants in the efavirenz-ART group and 1.6% (1/61) in the nevirapine-ART group were febrile.

Safety

From enrolment to follow-up day 63, there were 69 adverse events of special interest (AESIs) (54 and 15 in the efavirenz-ART group and nevirapine-ART group, respectively) and 12 serious adverse events (SAEs)

(10 and 2 in the efavirenz-ART group and nevirapine-ART group, respectively) (Table 3). None of the SAEs and AESIs were judged to be definitely related to DPQ. Forty-eight (30.0%) of the AESIs in the efavirenz-ART group and 8 (13.1%) in the nevirapine-ART group were judged to be probably related to DPQ (Table 3). These were mostly cases of prolonged QTcF interval. One death occurred on day 59 in a participant on efavirenz-ART who had been treated for deep vein thrombosis from day 33 and had CD4 count persistently below 50 since enrolment. Additional file 3 provides details on the SAEs that occurred during follow up. Except for the death, all participants with AESIs and SAEs recovered without sequelae.

Haematological parameters

The mean (SD) haemoglobin concentration decreased from baseline to day 7 (efavirenz-ART group: 12.2 to 11.7 g/dL, $p<0.001$ and nevirapine-ART group: 11.7 to 10.7 g/dL, $p<0.001$), but increased thereafter up to day 42, from 11.7 to 12.4 g/dL in the efavirenz-ART group ($p<0.001$) and from 10.7 to 11.6 g/dL in the nevirapine-ART group ($p<0.001$). Following DPQ treatment, the median (IQR) CD4 cell count increased from baseline to day 28, from 257 (140–357) to 320 (216–521) in the efavirenz-ART group ($p<0.001$) and from 390 (204–500) to 429 (204–580) in the nevirapine-ART group ($p=0.133$).

QT interval abnormalities

Data on baseline QTcF interval were available for 218 participants; 158 and 60 in the efavirenz-ART and nevirapine-ART group, respectively. The proportions of participants with predose QTcF ≥ 450 ms were 3.8% (6/158) in the efavirenz-ART group and 11.7% (7/60) in the nevirapine-ART group. Only 4 participants in the efavirenz-ART group had missing QTcF values on day 2 (last day of treatment). A change in QTcF interval of >60 ms from baseline to day 2 occurred in 31.2% (48/154) and 13.3% (8/60) in the efavirenz and nevirapine groups, respectively (Table 4). No participant had an absolute day 2 QTcF interval ≥ 500 ms. Observed QTcF interval abnormalities resolved by day 7 or 14 of follow up. No cardiovascular abnormalities were detected in these individuals.

Discussion

The WHO recommends the use of first-line anti-malarial drugs with PCR-corrected ACPR of >90%. In this study, the day 42 PCR-corrected ACPR for DPQ exceeded 99% among adult HIV-infected patients with uncomplicated malaria concurrently taking efavirenz- or nevirapine-based ART; the lower limit of the 95% CI for the DPQ ACPR exceeded 95%. Rapid parasite clearance, similar to that in HIV-uninfected individuals [28–31] was achieved,

Table 3 Summary of serious adverse events and adverse events of special interest stratified by antiretroviral group

Type of adverse event	DPQ and efavirenz-based ART regimen N = 160	DPQ and nevirapine- based ART regimen N = 61
	n ^a (%) [#] ^b	n ^a (%) [#] ^b
Serious adverse events (SAEs)	8 (5.0), [10]	2 (3.3), [2]
Adverse event(s) of special interest (AESI)	54 (33.8), [54]	15 (24.6), [15]
Fridericia corrected QT prolongation ^c	48 (30.0)	8 (13.1)
Palpitations	4 (2.5)	2 (3.3)
Dizziness	2 (1.3)	2 (3.3)
Urticaria	0	1 (1.6)
Itchiness	0	1 (1.6)
Chest pain	0	1 (1.6)
Drug-related adverse events (SAEs)		
Not related	7 (4.4), [9]	2 (3.3), [2]
Possibly related	1 (0.6), [1]	0
Probably related	0	0
Definitely related	0	0
Drug-related adverse events (AESI)		
Not related	1 (0.6), [1]	4 (6.6) [4]
Possibly related	5 (3.1), [5]	3 (4.9), [3]
Probably related	48 (30.0), [48]	8 (13.1), [8]
Definitely related	0	0

^a n The number of participants that experienced the event; % is percentage of participants experiencing that event in the intention-to-treat population

^b # The total number of events that occurred

^c Fridericia corrected QT prolongation defined as change in QTc interval of > 60 ms from baseline to day 2 (last day) of antimalarial treatment, also reported separately (in detail) in Table 4

Table 4 Median Fridericia corrected QT interval (QTcF), and proportion with abnormal ECG findings from baseline to last day of dosing among participants in the efavirenz- (N = 158) and nevirapine- (N = 60) based antiretroviral therapy groups

Time of ECG test	Median (IQR) QTcF in ms		Proportion with QTcF ≥ 450 ms		Proportion with QTcF change > 60 ms from baseline to last day of dosing on day 2 (taken within 2 h of dosing) ^a	
	Efavirenz-based ART group	Nevirapine-based ART group	Efavirenz-based ART group	Nevirapine-based ART group	Efavirenz ^b -based ART group	Nevirapine ^b -based ART group
	NA	NA	n (%)	n (%)	n (%)	n (%)
Day 0	395 (360–418)	418 (390–438)	6 (3.8)	7 (11.8)	NA	NA
Day 1	403 (383–423)	418 (388–437)	10 (6.3)	6 (10.0)	19 (12.3)	1 (1.7)
Day 2	424 (408–442)	434 (414–459)	29 (18.4)	21 (35.0)	48 (31.2)	8 (13.3)
Follow up visit after day 2 (day 7 or 14 of follow up)	410 (384–436)	424 (402–437)	0	0	NA	NA

IQR interquartile range, ECG electrocardiogram, ART antiretroviral drug, NA not applicable

^a Adverse event of special interest

^b N = 154 in Efavirenz group and 60 in the Nevirapine group

irrespective of the type of ART taken by the study participants. Malaria reinfections occurred in the efavirenz-ART group but, overall, day 42-PCR uncorrected ACPR

remained high in both ART groups, suggesting that DPQ had a good prophylactic effect against malaria reinfections. DPQ treatment also resulted in rapid resolution of

fever and marked improvement in haemoglobin concentrations irrespective of ART as well as improvement in CD4 cell count, especially in those taking efavirenz-based ART.

An anti-malarial treatment response depends on host immunity and anti-malarial drug blood concentrations. Efavirenz has been associated with induction of CYP3A4 and CYP2B6 enzymes while nevirapine has been associated with inhibition or sometimes induction of the enzymes [5]. Previous pharmacokinetic studies found that, compared with antiretroviral naïve HIV-infected individuals, those taking DPQ plus efavirenz-ART had lower piperazine area under the concentration–time curve (AUC) [8, 32] while those taking DPQ plus nevirapine-ART had higher piperazine AUC [8]. Similarly, other studies found that when co-administered with efavirenz or nevirapine, ACT resulted in altered pharmacokinetics of artemisinin and its metabolites [33–37]. The finding of high efficacy of DPQ suggests that any alterations in the pharmacokinetics of dihydroartemisinin or piperazine due to efavirenz or nevirapine had limited clinical significance in this study population. However, this needs further confirmation in future studies which would correlate PK data with observed efficacy endpoints.

To date, no previous study has assessed the efficacy and safety of DPQ in malaria-HIV coinfecting non-pregnant adults on ART. A previous Tanzanian study [38] found that HIV-infected adults with uncomplicated malaria on efavirenz-ART who were treated with artemether–lumefantrine had a lower PCR-uncorrected day 28 ACPR (82.5%) than HIV-infected antiretroviral naïve individuals (94.5%), while those on nevirapine-based ART had a higher day 28 PCR-uncorrected ACPR (97.6%). No cases of ETF were detected in a study population in which almost half of the participants had CD4 cell counts < 350. Thus, the study showed high efficacy of artemether and its metabolites (responsible for the initial clearance of malaria parasite [39, 40]) despite ART co-administration, which is consistent with our study findings. However, in contrast to our findings, the Tanzanian study found high rates of recurrent malaria in those on efavirenz-ART. This is likely to reflect the longer half life and hence superior prophylactic effect of the ACT-partner drug, piperazine, in this study compared with lumefantrine in the previous study.

Day-7 plasma ACT concentrations have been proposed as a marker of overall exposure of the longer acting partner drug of ACT and have been shown to be predictive of treatment efficacy by day 28 after malaria treatment [22]. In the present malaria-HIV co-infected sub-population, most participants had day-7 piperazine concentrations that were below the LLOQ. However, in participants

who had piperazine concentration values > LLOQ, the piperazine concentrations were not significantly different between those who experienced malaria recurrence by day 42 and those who did not. The observed < LLOQ concentrations could be due to increased metabolism of piperazine, as a result of efavirenz induction of CYP3A4 enzymes, or a limitation in the HPLC assay to detect smaller concentrations of piperazine. Nevertheless, the observed high efficacy of DPQ highlights that any PK interaction between efavirenz or nevirapine and piperazine did not predict the clinical outcomes in this study.

Delayed parasite clearance (parasite clearance half-life of > 5.5 h) has been shown to be associated with resistance of the parasites to artemisinins [41]. Although a smaller proportion of participants in both ART groups had parasite clearance half-life of > 5.5 h, they did not experience malaria recurrence in the follow up period, despite having most participants with piperazine concentrations below the lower limit of quantification. Notably, they all had a low baseline CD4 cell count with a relatively higher malaria parasite load compared to the rest of the participants. This higher parasite load coupled with low immunity at presentation could explain the delayed clearance of malaria parasites that was observed in these participants.

Treatment-emergent AESIs occurred in nearly a third and one-quarter of the participants on efavirenz- and nevirapine-based ART, respectively. However, only a few cases of AESIs in the efavirenz-group and none in the nevirapine-group were judged to be possibly associated with DPQ (Table 3). QTcF prolongation of at least 60 ms from baseline to the last treatment day did occur in a sizable proportion of participants but none had an absolute QTcF interval of 500 ms and there were no clinically detectable events. The QTcF prolongation resolved spontaneously by day 14. The observed QTcF prolongation may have been due to fever resolution [42, 43] rather than piperazine, but whatever the mechanism, we can conclude that DPQ can be safely used in this group of patients. This observation is also in line with WHO's recommendation that there is no evidence of increased risk of cardiotoxicity following exposure to current doses of DPQ for treatment of uncomplicated malaria [44]. Additionally, a similar phenomenon, of prolonged QTc interval which resolves following recovery from malaria, has been previously observed in 17.1% (n = 152) of adults living with HIV and on efavirenz-based ART who were treated for uncomplicated malaria with artemether–lumefantrine in Zambia [10]. Nevertheless, this needs to be confirmed in future studies of DPQ use in malaria-HIV coinfecting adults on efavirenz- or nevirapine based ART.

The major strengths of this study were the directly-supervised DPQ dosing, large sample size of participants on efavirenz-based ART and minimal loss to follow-up (<5%). However, the required sample size was not achieved in the nevirapine-based ART group because the national ART programme phased out this regimen during the course of the study. Nevertheless, participants in this group had high baseline parasite densities which permitted accurate assessment of DPQ's parasite clearance and prophylactic efficacy.

Nearly one-half of enrolled study participants in the efavirenz group had parasite densities of <2000/mm³ and nearly two-thirds were on cotrimoxazole prophylaxis which has some antimalarial effects [45, 46]. Thus, it can be argued that the high ACPR found in the efavirenz-group could have partly been due to the anti-malarial effects of cotrimoxazole and immunity-mediated clearance of low density parasitaemias. However, day-42 PCR-corrected and uncorrected ACPR did not significantly vary according to the use of cotrimoxazole prophylaxis or baseline parasite density. In addition, at least two-thirds of participants in this group had CD4 cell count <350/mm³ which is likely to have compromised their ability to clear parasites and prevent reinfections [2, 3, 25]. In spite of this, DPQ achieved high ACPR in this group suggesting its high therapeutic and prophylactic effectiveness.

In this study, the day-42 PCR corrected and uncorrected ACPR appeared to be higher among individuals on nevirapine-based ART than among those on efavirenz-based ART. However, this study was not designed to compare the efficacy and safety of DPQ between the two groups and the full sample size was not achieved in the nevirapine arm. It is therefore inappropriate to make direct comparison of DPQ efficacy between individuals taking the two ART regimens.

Conclusion

This study found that dihydroartemisinin–piperazine was highly efficacious and safe when used to treat uncomplicated *P. falciparum* malaria in individuals taking efavirenz or nevirapine-based ART. A higher than expected observed cases of QTc interval prolongation (>60 ms from baseline to day 2) following treatment with dihydroartemisinin–piperazine were observed and thought to likely be due to resolution of fever as part of the malaria recovery process. Under the “HIV Test and Treat” approach, many HIV-infected individuals in high burden countries will initiate ART early before they are severely immunosuppressed and this study supports the use of dihydroartemisinin–piperazine for the treatment of uncomplicated malaria in such patients.

Additional files

Additional file 1. Day 42 PCR-corrected efficacy plot. Kaplan–Meier survival plot of participants who were treated with dihydroartemisinin–piperazine (DHA PPQ) in the efavirenz (EFV)- and nevirapine (NVP) based antiretroviral therapy (ART) groups according to polymerase chain reaction (PCR) corrected adequate clinical and parasitological response (ACPR) by day 42 in the intention-to-treat population with loss to follow up and indeterminate or unavailable PCR samples treated as treatment success.

Additional file 2. Day 42 PCR-uncorrected efficacy plot. Kaplan–Meier survival plot of participants who were treated with dihydroartemisinin–piperazine (DHA PPQ) in the efavirenz (EFV)- and nevirapine (NVP) based antiretroviral therapy (ART) groups according to polymerase chain reaction (PCR) uncorrected adequate clinical and parasitological response (ACPR) by day 42 in the intention-to-treat population with loss to follow up and indeterminate or unavailable PCR samples treated as treatment success.

Additional file 3. Details of serious adverse events that occurred during follow up.

Abbreviations

ACT: artemisinin-based combination therapy; ACPR: adequate clinical and parasitological response; ART: antiretroviral therapy; DPQ: dihydroartemisinin–piperazine; ETF: Early treatment failure; LCF: Late clinical failure; LPF: Late parasitological failure.

Acknowledgements

We thank all trial participants for their participation in this study and the study team for their unwavering dedication.

Authors' contributions

ES, CGB, SMaculuve, SMacuacua, AV, MP and JM collected the data and supervised the day to day running of the study with support from VM. DJT, SHK, DGL and VM designed the experiment while VM conceived the idea of the study. CGB, LKP, MM and VM conducted the analyses and interpretation of the data. CGB compiled the first draft of the manuscript including generating the manuscript figures with support from ES and VM. All authors read and approved the final manuscript.

Funding

This work was supported by the European & Developing Countries Clinical Trials Partnership (EDCTP) (Grant Number IP07.31060.003 to VM). The funder had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Availability of data and materials

Data from this trial are held at the Malawi Liverpool Wellcome Trust Clinical Research Programme (MLW) which encourages optimal use of data by employing controlled access approach to data sharing with a robust system to review requests for data use and provide secure data access that is consistent with relevant ethics committee approvals. The datasets are therefore not publicly available but are available on reasonable request from the corresponding author and requests can also be initiated by contacting MLW: data@mlw.mw.

Ethics approval and consent to participate

The trial received ethics approval from the College of Medicine Research and Ethics Committee, Malawi; Comité Nacional de Bioética para Saúde, Mozambique and Liverpool School of Tropical Medicine Research Committee, UK and regulatory approval from the Pharmacy Medicines and Poisons Board, Malawi and Ministério da Saúde, Departamento Farmacêutico, Mozambique. Written informed consent was sought from all participants aged ≥ 18 years. Assent was sought from those who were aged < 18 years.

Consent for publication

Written informed consent to publish the data was sought from all participants aged ≥ 18 years. Assent was sought from those who were aged < 18 years, except for “mature minors”.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Centro de Investigação em Saúde de Manhiça (CISM), Maputo, Mozambique. ² Eduardo Mondlane University, Maputo, Mozambique. ³ University of Malawi, College of Medicine, Blantyre, Malawi. ⁴ Malawi Liverpool Wellcome Trust Clinical Research Programme, Blantyre, Malawi. ⁵ Oxford Centre for Tropical Medicine and Global Health, Oxford, UK. ⁶ Mahidol-Oxford Tropical Medicine Research Unit, Bangkok, Thailand. ⁷ Institute for Global Health, Universitat de Barcelona, Barcelona, Spain. ⁸ Liverpool School of Tropical Medicine, Liverpool, UK. ⁹ University of Liverpool, Liverpool, UK. ¹⁰ Tropical and Infectious Diseases Unit, Royal Liverpool University Hospital, Liverpool, UK.

Received: 14 February 2019 Accepted: 12 August 2019

Published online: 20 August 2019

References

- Kwenti TE. Malaria and HIV coinfection in sub-Saharan Africa: prevalence, impact, and treatment strategies. *Res Rep Trop Med*. 2018;9:123–36.
- Hewitt K, Steketee R, Mwapasa V, Whitworth J, French N. Interactions between HIV and malaria in non-pregnant adults: evidence and implications. *AIDS*. 2006;20:1993–2004.
- Laufer MK, van Oosterhout JGG, Thesing PC, Thumba F, Zijlstra EE, Graham SM, et al. Impact of HIV-associated immunosuppression on malaria infection and disease in Malawi. *J Infect Dis*. 2006;193:872–8.
- WHO. Guidelines for the treatment of malaria. 3rd ed. Geneva: World Health Organization; 2015.
- Khoo S, Back D, Winstanley P. The potential for interactions between antimalarial and antiretroviral drugs. *AIDS*. 2005;19:995–1005.
- Lee TM-N, Huang L, Johnson MK, Lizak P, Kroetz D, Aweeka F, et al. In vitro metabolism of piperazine is primarily mediated by CYP3A4. *Xenobiotica*. 2012;42:1088–95.
- Seden K, Gibbons S, Marzolini C, Schapiro JM, Burger DM, Back DJ, et al. Development of an evidence evaluation and synthesis system for drug–drug interactions, and its application to a systematic review of HIV and malaria co-infection. *PLoS ONE*. 2017;12:e0173509.
- Banda CG, Dzinjalimala F, Mukaka M, Mallewa J, Maiden V, Terlouw DJ, et al. Pharmacokinetics of piperazine and safety profile of dihydroartemisinin–piperazine coadministered with antiretroviral therapy in malaria-uninfected HIV-positive Malawian adults. *Antimicrob Agents Chemother*. 2018;62:e00634–18.
- WHO. Methods for surveillance of antimalarial drug efficacy. Geneva: World Health Organization; 2009.
- Banda CG, Chaponda M, Mukaka M, Mulenga M, Hachizovu S, Kabuya JB, et al. Efficacy and safety of artemether–lumefantrine as treatment for *Plasmodium falciparum* uncomplicated malaria in adult patients on efavirenz-based antiretroviral therapy in Zambia: an open label non-randomized interventional trial. *Malar J*. 2019;18:180.
- Malawi malaria indicator survey 2014. Ministry of Health National Malaria Control Programme, Malawi. 2015.
- Galatas B, Guinovart C, Bassat Q, Aponte JJ, Nhamússua L, Macete E, et al. A prospective cohort study to assess the micro-epidemiology of *Plasmodium falciparum* clinical malaria in Ilha Josina Machel (Manhiça, Mozambique). *Malar J*. 2016;15:444.
- National Statistical Office. Malawi demographic and health survey 2015–16; 2017.
- Ministerio da Saúde. National survey on prevalence, behavioral risks and information about HIV and AIDS in Mozambique (INSIDE). Maputo: Ministerio da Saúde; 2009.
- WHO. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection. Geneva: World Health Organization; 2013.
- Ministry of Health. Malawi integrated HIV program report. 2013.
- WHO. Management of severe malaria—a practical handbook. 3rd ed. Geneva: World Health Organization; 2013.
- DAIDS. Division of AIDS table for grading the severity of adult and pediatric adverse events. 2004. p. 1–20. <https://rsc.niaid.nih.gov/sites/default/files/daids-ae-grading-table-v2-nov2014.pdf>. Accessed 21 Jan 2019.
- WPRO. Malaria microscopy standard operating procedures. WHO Western Pacific Region; 2018.
- Reeder JC, Marshall VM. A simple method for typing *Plasmodium falciparum* merozoite surface antigens 1 and 2 (MSA-1 and MSA-2) using a dimorphic-form specific polymerase chain reaction. *Mol Biochem Parasitol*. 1994;68:329–32.
- WHO. Methods and techniques for assessing exposure to antimalarial drugs in clinical field studies. Geneva: World Health Organization; 2016.
- White NJ, Stepniewska K, Barnes K, Price RN, Simpson J. Simplified antimalarial therapeutic monitoring: using the day-7 drug level? *Trends Parasitol*. 2008;24:159–63.
- Lourens C, Watkins WM, Barnes KI, Sibley CH, Guerin PJ, White NJ, et al. Implementation of a reference standard and proficiency testing programme by the World Wide Antimalarial Resistance Network (WWARN). *Malar J*. 2010;9:375.
- ICH. International conference on harmonisation of technical requirements for registration of pharmaceuticals for human use: Clinical safety data management: E2A; 1994.
- Van Geertruyden J-P, Mulenga M, Mwananyanda L, Chalwe V, Moerman F, Chilengi R, et al. HIV-1 immune suppression and antimalarial treatment outcome in Zambian adults with uncomplicated malaria. *J Infect Dis*. 2006;194:917–25.
- Kirkwood BR, Sterne JAC, Kirkwood BR. Essential medical statistics. Hoboken: Blackwell Science; 2003.
- Flegg JA, Guerin PJ, White NJ, Stepniewska K. Standardizing the measurement of parasite clearance in falciparum malaria: the parasite clearance estimator. *Malar J*. 2011;10:339.
- Gargano N, Ubben D, Tommasini S, Bacchieri A, Corsi M, Bhattacharyya PC, et al. Therapeutic efficacy and safety of dihydroartemisinin–piperazine versus artesunate–mefloquine in uncomplicated *Plasmodium falciparum* malaria in India. *Malar J*. 2012;11:233.
- Sylla K, Abiola A, Tine RCK, Faye B, Sow D, Ndiaye JL, et al. Monitoring the efficacy and safety of three artemisinin based-combinations therapies in Senegal: results from two years surveillance. *BMC Infect Dis*. 2013;13:598.
- Sow D, Ndiaye J-L, Sylla K, Ba MS, Tine RCK, Faye B, et al. Evaluation of the efficacy and safety of three 2-drug combinations for the treatment of uncomplicated *Plasmodium falciparum* malaria in Senegal: artesunate–amodiaquine, dihydroartemisinin–piperazine, and artemether–lumefantrine. *Med Sante Trop*. 2015;26:45–50.
- Grandesso F, Guindo O, Woi Messe L, Makarimi R, Traore A, Dama S, et al. Efficacy of artesunate–amodiaquine, dihydroartemisinin–piperazine and artemether–lumefantrine for the treatment of uncomplicated *Plasmodium falciparum* malaria in Maradi, Niger. *Malar J*. 2018;17:52.
- Kajubi R, Huang L, Jagannathan P, Chamankhah N, Were M, Ruel T, et al. Antiretroviral therapy with efavirenz accentuates pregnancy-associated reduction of dihydroartemisinin–piperazine exposure during malaria chemoprevention. *Clin Pharmacol Ther*. 2017;102:520–8.
- Byakika-Kibwika P, Lamorde M, Mayito J, Nabukeera L, Namakula R, Mayanja-Kizza H, et al. Significant pharmacokinetic interactions between artemether/lumefantrine and efavirenz or nevirapine in HIV-infected Ugandan adults. *J Antimicrob Chemother*. 2012;67:2213–21.
- Hoglund RM, Byakika-Kibwika P, Lamorde M, Merry C, Ashton M, Han-pithakpong W, et al. Artemether–lumefantrine co-administration with antiretrovirals: population pharmacokinetics and dosing implications. *Br J Clin Pharmacol*. 2015;79:636–49.
- Kredo T, Mauff K, Van Der Walt JS, Wiesner L, Maartens G, Cohen K, et al. Interaction between artemether–lumefantrine and nevirapine-based antiretroviral therapy in HIV-1-infected patients. *Antimicrob Agents Chemother*. 2011;55:5616–23.
- Fehintola FA, Scarsi KK, Ma Q, Parikh S, Morse GD, Taiwo B, et al. Nevirapine-based antiretroviral therapy impacts artesunate and dihydroartemisinin disposition in HIV-infected Nigerian adults. *AIDS Res Treat*. 2012;2012:703604.
- Scarsi KK, Fehintola FA, Ma Q, Aweeka FT, Darin KM, Morse GD, et al. Disposition of amodiaquine and desethylamodiaquine in HIV-infected Nigerian subjects on nevirapine-containing antiretroviral therapy. *J Antimicrob Chemother*. 2014;69:1370–6.
- Maganda BA, Minzi OMS, Kamuhabwa AAR, Ngasala B, Sasi PG. Outcome of artemether–lumefantrine treatment for uncomplicated malaria in HIV-infected adult patients on anti-retroviral therapy. *Malar J*. 2014;13:205.
- White N. The parasite clearance curve. *Malar J*. 2011;10:278.

40. WWARN Parasite Clearance Study Group WPCS, Abdulla S, Ashley EA, Bassat Q, Bethell D, Björkman A, et al. Baseline data of parasite clearance in patients with falciparum malaria treated with an artemisinin derivative: an individual patient data meta-analysis. *Malar J*. 2015;14:359.
41. WWARN K13 Genotype-Phenotype Study Group. Association of mutations in the *Plasmodium falciparum* Kelch13 gene (Pf3D7_1343700) with parasite clearance rates after artemisinin-based treatments—a WWARN individual patient data meta-analysis. *BMC Med*. 2019;17:1.
42. Mytton OT, Ashley EA, Peto L, Price RN, La Y, Hae R, et al. Electrocardiographic safety evaluation of dihydroartemisinin piperazine in the treatment of uncomplicated falciparum malaria. *Am J Trop Med Hyg*. 2007;77:447–50.
43. Price RN, Nosten F, White NJ. Prolongation of the QTc interval in African children treated for falciparum malaria. *Am J Trop Med Hyg*. 1998;59:503.
44. WHO Evidence Review Group. The cardiotoxicity of antimalarials. 2017. <http://www.who.int/malaria/mpac/mpac-mar2017-erg-cardiotoxicity-report-session2.pdf>. Accessed 21 Jan 2019.
45. Manyando C, Njunju EM, D'Alessandro U, van Geertruyden J-P. Safety and efficacy of co-trimoxazole for treatment and prevention of *Plasmodium falciparum* malaria: a systematic review. *PLoS ONE*. 2013;8:e56916.
46. Mbeye NM, ter Kuile FO, Davies M-A, Phiri KS, Egger M, Wandeler G, et al. Cotrimoxazole prophylactic treatment prevents malaria in children in sub-Saharan Africa: systematic review and meta-analysis. *Trop Med Int Health*. 2014;19:1057–67.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

