

The persistent high burden of child and maternal malnutrition in most states of India:

the Global Burden of Disease Study 2016

Soumya Swaminathan et al, for the India State-level Disease Burden Initiative Collaborators

Web Appendix

Correspondence to: Lalit Dandona, lalit.dandona@phfi.org

Table 1: GBD 2016 India data inputs for malnutrition indicators and covariates	3
Table 2: Percent contribution of under-five and all-age DALYs attributable to child and maternal malnutrition and its components in India by sex in 2016	21
Table 3: Prevalence of low birth weight for India and states grouped by epidemiological transition level, 1990-2016.....	22
Table 4: Prevalence of wasting in under-five children for India and states grouped by epidemiological transition level, 1990-2016	23
Table 5: Prevalence of stunting in under-five children for India and states grouped by epidemiological transition level, 1990-2016	24
Table 6: Prevalence of exclusive breastfeeding in the first six months for India and states grouped by epidemiological transition level, 1990-2016	25
Table 7: Prevalence of anaemia in reproductive age women for India and states grouped by epidemiological transition level, 1990-2016	26
Table 8: Prevalence of overweight in children aged 2-4 years for India and states grouped by epidemiological transition level, 1990-2016	27

Table 1: GBD 2016 India data inputs for malnutrition indicators and covariates

Abel R, Rajaratnam J, Gnanasekaran VJ, Jayaraman P. Prevalence of anaemia and iron deficiency in three trimesters in Rural Vellore district, South India. <i>Trop Doct.</i> 2001; 31(2): 86-9.
Abel R, Rajaratnam J, Kalaimani A, Kirubakaran S. Can iron status be improved in each of the three trimesters? A community-based study. <i>Eur J Clin Nutr.</i> 2000; 54(6): 490-3.
Abel R, Sampathkumar V. Tamil Nadu nutritional survey comparing children aged 0-3 years with the NCHS/CDC reference population. <i>Indian J Pediatr.</i> 1998; 65(4): 565-72.
Agarwal AK, Sen AK, Kalra NK, Gupta N. Prevalence of anaemia during pregnancy in district Burdwan, West Bengal. <i>Indian J Public Health.</i> 1999; 43(1): 26-31.
Agarwal D, Pandey C, Agarwal K. Vitamin A administration and preschool child mortality. <i>Nutr Res.</i> 1995; 15(5): 669-80. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. <i>Cochrane Database Syst Rev.</i> 2010; CD008524.
Agarwal DK, Agarwal KN, Tripathi AM. Nutritional status in rural pregnant women of Bihar and Uttar Pradesh. <i>Indian Pediatr.</i> 1987; 24(2): 119-25.
Agarwal DK, Bhardwaj B, Singla PN, Tripathi AM, Agarwal KN. Etiology of maternal and early childhood deficiency anemia. <i>Indian J Pediatr.</i> 1986; 53(3): 389-95.
Agarwal KN, Agarwal DK, Mishra KP. Impact of anaemia prophylaxis in pregnancy on maternal haemoglobin, serum ferritin & birth weight. <i>Indian J Med Res.</i> 1991; 277-80.
Agarwal KN, Gomber S, Bisht H, Som M. Anemia prophylaxis in adolescent school girls by weekly or daily iron-folate supplementation. <i>Indian Pediatr.</i> 2003; 40(4): 296-301.
Arya S, Chellani H, Pandey J. Evaluation of safety of oral vitamin "A" megadose co-administered with measles vaccination. <i>Indian Pediatr.</i> 2000; 37(12): 1341-7. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. <i>Cochrane Database Syst Rev.</i> 2010; CD008524.
Awasthi S, Das R, Verma T, Vir S. Anemia and undernutrition among preschool children in Uttar Pradesh, India. <i>Indian Pediatr.</i> 2003; 40(10): 985-90.
Awasthi S, Pande VK. Cause-specific mortality in under fives in the urban slums of Lucknow, north India. <i>J Trop Pediatr.</i> 1998; 44(6): 358-61.
Awasthi S, Peto R, Read S, Clark S, Pande V, Bundy D, DEVTA (Deworming and Enhanced Vitamin A) team. Vitamin A supplementation every 6 months with retinol in 1 million pre-school children in north India: DEVTA, a cluster-randomised trial. <i>Lancet.</i> 2013; 381(9876): 1469-77.
Bains K, Mann SK. Physical Fitness in Relation to Energy and Iron Status of Female College Students. <i>Food Nutr Bull.</i> 2000; 21(3):305-10.
Bamji MS, V S Murthy PV, Williams L, Vardhana Rao MV. Maternal nutritional status & practices & perinatal, neonatal mortality in rural Andhra Pradesh, India. <i>Indian J Med Res.</i> 2008; 127(1): 44-51.
Bang AT, Bang RA, Baitule SB, Reddy MH, Deshmukh MD. Effect of home-based neonatal care and management of sepsis on neonatal mortality: field trial in rural India. <i>Lancet.</i> 1999; 354(9194): 1955-61.
Bang AT, Bang RA, Reddy HM, Deshmukh MD, Baitule SB. Reduced incidence of neonatal morbidities: effect of home-based neonatal care in rural Gadchiroli, India. <i>J Perinatol.</i> 2005; 25 (Suppl 1): S51-61.
Baruah J, Kusre G, Bora R. Pattern of Gross Congenital Malformations in a Tertiary Referral Hospital in Northeast India. <i>Indian J Pediatr.</i> 2015; 82(10): 917-22.

Bhandari N, Mazumder S, Taneja S, Sommerfelt H, Strand TA. Effect of implementation of Integrated Management of Neonatal and Childhood Illness (IMNCI) programme on neonatal and infant mortality: cluster randomised controlled trial. *BMJ*. 2012; 344: e1634.

Bharati P, Ghosh R, Gupta R. Socioeconomic Condition and Anaemia among the Mahishya Population of Southern West Bengal, India. *Malays J Nutr*. 2004; 10(1): 23-30.

Bhargava M, Iyer PU, Kumar R, Ramji S, Kapani V, Bhargava SK. Relationship of maternal serum ferritin with foetal serum ferritin, birth weight and gestation. *J Trop Pediatr*. 1991; 37(4): 149-52.

Bhargava SK, Singh KK, Saxena BN, India Council of Medical Research (ICMR). ICMR Task Force National Collaborative Study on Identification of High Risk Families, Mothers and Outcome of their Off-springs with particular reference to the problem of maternal nutrition, low birth weight, perinatal and infant morbidity and mortality in rural and urban slum communities. Summary, conclusions and recommendations. 1991; 28(12): 1473-80.

Bisai S, Mahalanabis D, Sen A, Bose K, Datta N. Maternal early second trimester pregnancy weight in relation to birth outcome among Bengalee Hindus of Kolkata, India. *Ann Hum Biol*. 2007; 34(1): 91-101.

Biswas R, Biswas AB, Manna B, Bhattacharya SK, Dey R, Sarkar S. Effect of vitamin A supplementation on diarrhoea and acute respiratory tract infection in children. A double blind placebo controlled trial in a Calcutta slum community. *Eur J Epidemiol*. 1994; 10(1): 57–61. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. *Cochrane Database Syst Rev*. 2010; CD008524.

Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, Hogan D, Shiekh S, Qureshi ZU, You D, Lawn JE; Lancet Stillbirth Epidemiology Investigator Group. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health*. 2016; 4(2): e98-108.

Brabin L, Nicholas S, Gogate A, Gogate S, Karande A. High prevalence of anaemia among women in Mumbai, India. *Food Nutr Bull*. 1998; 19(3): 205-9.

Brahmam G, Madhavan NK, Laxmaiah A, Reddy C, Ranganathan S, Vishnuvardhana Rao M, Nadamuni Naidu A, Vijayaraghavan K, Sivakumar B, Kamala Krishnaswamy, Gowarinti Sastry J, Mohan Ram M, Praldad Rao N, Vinodini R. Community Trials with Iron and Iodine Fortified Salt (Double Fortified Salt). In: Geertman RM, ed. *Proceedings of the 8th World Salt Symposium*. Amsterdam, The Netherlands: Elsevier; 2000. 955-60.

Center for Vaccine Development, University of Maryland, Centers for Disease Control and Prevention, Department of Medical Microbiology and Immunology, Göteborg University, International Vaccine Institute, National Institute of Cholera and Enteric Diseases, Perry Point Cooperative Studies Program Coordinating Center, U.S. Department of Veterans Affairs, School of Medicine, University of Virginia, University of Chile. India - Kolkata Global Enteric Multicenter Study 2007-2011. Baltimore, MD, United States: Center for Vaccine Development, University of Maryland.

Center for Vaccine Development, University of Maryland, Centers for Disease Control and Prevention, Department of Medical Microbiology and Immunology, Göteborg University, International Vaccine Institute, National Institute of Cholera and Enteric Diseases, Perry Point Cooperative Studies Program Coordinating Center, U.S. Department of Veterans Affairs, School of Medicine, University of Virginia, University of Chile. India - Kolkata Global Enteric Multicenter Study 2011-2013. Baltimore, United States: Center for Vaccine Development, University of Maryland.

Centre for Research on the Epidemiology of Disasters (CRED), Catholic University of Leuven. EM-DAT: The OFDA/CRED International Disaster Database. Brussels, Belgium: Catholic University of Leuven.

Chakraborty B, Bhattacharya AR. Xerophthalmia in under-five children in West Bengal. Indian Pediatr. 1987; 24(7): 600-2.

Chaturvedi S, Kapil U, Gnanasekaran N, Sachdev HP, Pandey RM, Bhanti T. Nutrient intake amongst adolescent girls belonging to poor socioeconomic group of rural area of Rajasthan. Indian Pediatr. 1996; 33(3): 197-201.

Chiplonkar S, Agte V, Mengale S. Relative importance of micronutrient deficiencies in iron deficiency anemia. Nutr Res. 2003; 23(10): 1355-67.

Chowdhury S, Kumar R, Ganguly NK, Kumar L, Walia BNS. Effect of vitamin A supplementation on childhood morbidity and mortality. Indian J Med Sci. 2002; 56(6): 259-64. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. Cochrane Database Syst Rev. 2010; CD008524.

Christian Medical College - Vellore, Fogarty International Center, National Institutes of Health, Foundation for the National Institutes of Health. India - Vellore Malnutrition and Enteric Disease (MAL-ED) Study 2009-2014. [Data shared for this analysis]

Dahiya S, Kapoor AC. Anaemia in women in a selected rural area of Haryana: effect of dietary intake level. Indian J Nutr Diet. 1995; 224-8.

Das S, Bapat U, More NS, Alcock G, Fernandez A, Osrin D. Nutritional status of young children in Mumbai slums: a follow-up anthropometric study. Open Nutr J. 2012; 11(1): 100.

Department of Women and Child Development, Ministry of Human Resource Development, Government of India, United Nations Children's Fund (UNICEF) - India Country Office. India Summary Report on the Multiple Indicator Cluster Survey 2000. New Delhi, India: UNICEF India Country Office; 2000.

Department of Women and Child Development, Ministry of Human Resource Development, Government of India, United Nations Children's Fund (UNICEF) - India Country Office. India Summary Report on the Multiple Indicator Cluster Survey 2000. New Delhi, India: UNICEF India Country Office; 2000.

Department of Women and Child Development, Ministry of Human Resource Development, Government of India. India Nutrition Profile Survey 1995-1996. New Delhi, India: Ministry of Human Resource Development.

Desai, Sonalde, Reeve Vanneman, National Council of Applied Economic Research, University of Michigan. India Human Development Survey 2005. Ann Arbor, Michigan: Inter-University Consortium for Political and Social Research.

Deshmukh V, Lahariya C, Krishnamurthy S, Das MK, Pandey RM, Arora NK. Taken to Health Care Provider or Not, Under-Five Children Die of Preventable Causes: Findings from Cross-Sectional Survey and Social Autopsy in Rural India. Indian J Community Med. 2016; 41(2): 108-19.

Directorate of Economics & Statistics and Office of Chief Registrar (Births & Deaths), Government of National Capital Territory of Delhi. Report on Medical Certification of Cause of Deaths in Delhi-2013. New Delhi, India: Directorate of Economics & Statistics and Office of Chief Registrar (Births & Deaths); 2014.

Fakhri S, Srivastava I, Ahmad P, Hasan SS. Prevalence of xerophthalmia in pre-school children in an urban slum. Indian Pediatr. 1993; 30(5): 668-70.

Fall CHD, Sachdev HS, Osmond C, Lakshmy R, Biswas SD, Prabhakaran D, Tandon N, Ramji S, Reddy KS, Barker DJP, Bhargava SK. Adult metabolic syndrome and impaired glucose tolerance are associated with different patterns of BMI gain during infancy: Data from the New Delhi Birth Cohort. *Diabetes Care*. 2008; 31(12): 2349-56.

FAO Supply Utilization Accounts 1961-2013. Personal Correspondence with Dr. Josef Schmidhuber, 2016. [Data shared for this analysis]

Feldon K, Bahl S, Bhatnagar P, Wenger J. Severe vitamin A deficiency in India during pulse polio immunization. *Indian J Med Res*. 2005; 122(3): 265-7.

Food and Agriculture Organization of the United Nations (FAO). FAOSTAT Food Balance Sheets, October 2014. Rome, Italy: FAO.

Ford Foundation, Nutrition Foundation of India. India Profiles of Undernutrition and Underdevelopment: Studies of Poor Communities in Seven Regions of the Country. New Delhi, India: Nutrition Foundation of India. (Scientific Report Series No. 8).

George K, Prasad J, Singh D, Minz S, Albert DS, Mulyil J, Joseph KS, Jayaraman J, Kramer MS. Perinatal outcomes in a South Asian setting with high rates of low birth weight. *BMC Pregnancy Childbirth*. 2009 Feb; 9:5.

Ghosh R, Bharati P. Haemoglobin status of adult women of two ethnic groups living in a peri-urban area of Kolkata city, India: a micro-level study. *Asia Pac J Clin Nutr*. 2003; 12(4): 451-9.

Ghosh SK, Yadav RS, Das BS, Sharma VP. Influence of nutritional and haemoglobin status on malaria infection in children. *Indian J Pediatr*. 1995; 62(3): 321-6.

Global Burden of Disease Health Financing Collaborator Network, Institute for Health Metrics and Evaluation (IHME). Global Development Assistance for Health, Government, Prepaid Private, and Out-of-Pocket Health Spending 1995-2014. Seattle, United States: IHME; 2017.

Gomber S, Kumar S, Rusia U, Gupta P, Agarwal KN, Sharma S. Prevalence & etiology of nutritional anaemias in early childhood in an urban slum. *Indian J Med Res*. 1998; 107: 269-73.

Goudar SS, Goco N, Somannavar MS, Vernekar SS, Mallapur AA, Moore JL, Wallace DD, Sloan NL, Patel A, Hibberd PL, Koso-Thomas M, McClure EM, Goldenberg RL. Institutional deliveries and perinatal and neonatal mortality in Southern and Central India. *Reprod Health*. 2015; 12(Suppl 2): S13.

Government of Orissa, Micronutrient Initiative, United Nations Children's Fund, World Health Organization. Orissa Impact of Vitamin A Supplementation Delivered with Oral Polio Vaccine as Part of Immunization Campaign [Draft final report]. Bhubheshwar, India: Government of Orissa; 2001.

Gupta PC, Subramoney S, Sreevidya S. Smokeless tobacco use, birth weight, and gestational age: population based, prospective cohort study of 1217 women in Mumbai, India. *BMJ*. 2004; 328(7455): 1538.

Gupta V, Shukla K. Epidemiology of anaemia in pre-school children from a rural and a slum community of Varanasi. *Indian J Prev Soc Med*. 1985; 85-9.

Gupte SC, Shaw AN, Shah KC. Hematological findings and severity of G6PD deficiency in Vataliya Prajapati subjects. *J Assoc Physicians India*. 2005; 53: 1027-30.

Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2007-2009. New Delhi, India: Ministry of Health and Family Welfare.

Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2008-2010. New Delhi, India: Ministry of Health and Family Welfare.

Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2009-2011. New Delhi, India: Ministry of Health and Family Welfare.
Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2010-2012. New Delhi, India: Ministry of Health and Family Welfare.
Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2011-2013. New Delhi, India: Ministry of Health and Family Welfare.
Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2012-2014. New Delhi, India: Ministry of Health and Family Welfare.
Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2013-2015. New Delhi, India: Ministry of Health and Family Welfare.
Health Management Information System (HMIS), National Health Mission, Ministry of Health and Family Welfare, Government of India. India Performance Related to Immunisation 2014-2016. New Delhi, India: Ministry of Health and Family Welfare.
Hebert JR, Gupta PC, Bhonsle RB, Sinor PN, Mehta H, Mehta FS. Development and testing of a quantitative food frequency questionnaire for use in Gujarat, India. <i>Public Health Nutr.</i> 1999; 2(1): 39–50.
Hirve S, Ganatra B. A prospective cohort study on the survival experience of under five children in rural western India. <i>Indian Pediatr.</i> 1997; 34(11): 995-1001.
India Neonatal Preterm Birth Complications Data 1995 as provided by the Global Burden of Disease 2010 Neonatal Conditions Expert Group. [Data shared for this analysis]
Indian Council of Medical Research (ICMR). India Study on Causes of Death by Verbal Autopsy 2003. New Delhi, India: ICMR. [Data shared for this analysis]
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, Macro International. India National Family Health Survey (NFHS-1) 1992-1993: National Report. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, Macro International. India National Family Health Survey Data (NFHS-1) 1992-1993. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, ORC Macro. India National Family Health Survey (NFHS-2) 1998-1999: National and State Reports. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, ORC Macro. India National Family Health Survey Data (NFHS-2) 1998-1999. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, Macro International. India National Family Health Survey (NFHS-3) 2005-2006: National and State Reports. Mumbai, India: IIPS.

International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, Macro International. India National Family Health Survey Data (NFHS-3) 2005-2006. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India, ICF. India National Family Health Survey (NFHS-4) 2015-2016: National and State-level Factsheets. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household Survey (DLHS-1) 1998-1999: National Report. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household Survey Data (DLHS-1) 1998-1999. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household Survey (DLHS-2) 2002-2004: National and State Reports. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household Survey Data (DLHS-2) 2002-2004. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household and Facility Survey (DLHS-3) 2007-2008: National and State Reports. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household and Facility Survey Data (DLHS-3) 2007-2008. Mumbai, India: IIPS.
International Institute for Population Sciences (IIPS), Ministry of Health and Family Welfare, Government of India. India District Level Household and Facility Survey Data (DLHS-4) 2012-2013: State Reports. Mumbai, India: IIPS.
International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. India District Level Household and Facility Survey Data (DLHS-4) 2012-2013. Mumbai, India: IIPS.
International Institute for Population Sciences, World Health Organization. India World Health Survey 2003.
International Social Survey Programme (ISSP), GESIS - Leibniz Institute of Social Sciences. ISSP: Family and Changing Gender Roles IV, ZA5900 (Version 4.0.0). Mannheim, Germany: GESIS - Leibniz Institute of Social Sciences; 2012.
Jain DL, Sarathi V, Upadhye D, Gulhane R, Nadkarni AH, Ghosh K, Colah RB. Newborn screening shows a high incidence of sickle cell anemia in Central India. <i>Hemoglobin</i> . 2012; 36(4): 316–22.
Jain S, Chopra H, Garg SK, Bhatnagar M, Singh JV. Anemia in children: early iron supplementation. <i>Indian J Pediatr</i> . 2000; 67(1): 19-21.
Kameswaran C, Bhatia BD, Bhat BV, Oumachigui A. Perinatal mortality: a hospital based study. <i>Indian Pediatr</i> . 1993; 30(8): 997-1001.
Kapil U, Pathak P, Tandon M, Singh C, Pradhan R, Dwivedi SN. Micronutrient deficiency disorders amongst pregnant women in three urban slum communities of Delhi. <i>Indian Pediatr</i> . 1999; 36(10): 983-9.
Kapoor D, Agarwal KN, Sharma S, Kela K, Kaur I. Iron status of children aged 9-36 months in an urban slum Integrated Child Development Services project in Delhi. <i>Indian Pediatr</i> . 2002; 39(2): 136-44.

Kapoor G, Aneja S. Nutritional disorders in adolescent girls. Indian Pediatr. 1992; 29(8): 969-73.
Kaushik SL, Parmar VR, Grover N, Kaushik R. Neonatal mortality rate: relationship to birth weight and gestational age. Indian J Pediatr. 1998; 65(3): 429-33.
Khandait DW, Ambadekar NN, Zodpey SP, Vasudeo ND. Maternal age as a risk factor for stillbirth. Indian J Public Health. 2000; 44(1): 28-30.
Kodkany BS, Derman RJ, Honnungar NV, Tyagi NK, Goudar SS, Mastiholi SC, Moore JL, McClure EM, Sloan N, Goldenberg RL. Establishment of a Maternal Newborn Health Registry in the Belgaum District of Karnataka, India. Reprod Health. 2015; 12 Suppl 2: S3.
Koen M, Lemson M, Sampathkumar V, Abel R. Prevalence of anaemia among pregnant mothers in a rural South Indian population. J Obstet Gynaecol India. 1992; 42(6): 283-7.
Kumar CSV, Anand Kumar H, Sunita V, Kapur I. Prevalence of anemia and worm infestation in school going girls at Gulbargha, Karnataka. Indian Pediatr. 2003; 40(1): 70-2.
Kumar V, Datta N, Wadhwa SS, Singh S. Morbidity and mortality in diarrhea in rural Haryana Indian. Indian J Pediatr. 1985; 52(418): 455-61.
Luwang N, Gupta V. Anaemia in pregnancy in a rural community-influence of dietary intake in the multifactorial aetiology. Indian J Nutr Diet. 1980; 17(11): 414-7.
Mahapatra A, Geddam JJ, Marai N, Murmu B, Mallick G, Bulliyya G, Acharya AS, Satyanarayana K. Nutritional status of preschool children in the drought affected Kalahandi district of Orissa. Indian J Med Res. 2000; 111: 90-4.
Mahavarkar SH, Madhu CK, Mule VD. A comparative study of teenage pregnancy. J Obstet Gynaecol. 2008; 28(6): 604-7.
Mathad V, Metgud C, Mallapur MD. Nutritional status of under-fives in rural area of South India. Indian J Med Sci. 2011; 65(4): 151-6.
Max Planck Institute for Demographic Research, Vienna Institute of Demography. Human Fertility Collection - Total Fertility Rate and Mean Age at Birth. Rostock, Germany: Max Planck Institute for Demographic Research.
McClure EM, Pasha O, Goudar SS, Chomba E, Garces A, Tshefu A, Althabe F, Esamai F, Patel A, Wright LL, Moore J, Kodkany BS, Belizan JM, Saleem S, Derman RJ, Carlo WA, Hambidge KM, Buekens P, Liechty EA, Bose C, Koso-Thomas M, Jobe AH, Goldenberg RL; Global Network Investigators. Epidemiology of stillbirth in low-middle income countries: A Global Network Study. Acta Obstet Gynecol Scand. 2011; 90(12): 1379-85.
Mehrotra SK, Mathur JS, Maheshwari BB. Epidemiological aspects of nutritional anaemia in children below five years. Indian J Pediatr. 1976; 43(340): 132-5.
Meshram II, Arlappa N, Balakrishna N, Laxmaiah A, Mallikarjun Rao K, Gal Reddy C, Ravindranath M, Sharad Kumar S, Brahmam GNV. Prevalence and determinants of undernutrition and its trends among pre-school tribal children of Maharashtra State, India. J Trop Pediatr. 2012; 58(2): 125-32.
Meshram II, Arlappa N, Balakrishna N, Mallikharjuna Rao K, Laxmaiah A, Brahmam GN. Trends in the prevalence of undernutrition, nutrient & food intake and predictors of undernutrition among under five year tribal children in India. Asia Pac J Clin Nutr. 2012; 21(4): 568-76.
Meshram II, Balakrishna N, Arlappa N, Rao KM, Laxmaiah A, Brahmam GNV. Prevalence of Undernutrition, Its Determinants, and Seasonal Variation Among Tribal Preschool Children of Odisha State, India. Asia Pac J Public Health. 2014; 26(5): 470-80.

Meshram II, Kodavanti MR, Chitty GR, Manchala R, Kumar S, Kakani SK, Kodavalla V, Avula L, Ginnela Narsimhachary Veera B. Influence of Feeding Practices and Associated Factors on the Nutritional Status of Infants in Rural Areas of Madhya Pradesh State, India. Asia Pac J Public Health. 2015; 27(2): NP1345-61.
Meshram II, Laxmaiah A, Venkaiah K, Brahmam GNV. Impact of feeding and breastfeeding practices on the nutritional status of infants in a district of Andhra Pradesh, India. Natl Med J India. 2012; 25(4): 2016.
Ministry of Health and Family Welfare, Government of India, United Nations Children's Fund. India Coverage Evaluation Survey 2005.
Ministry of Health and Family Welfare, Government of India, United Nations Children's Fund. India Coverage Evaluation Survey 2007.
Ministry of Health and Family Welfare, Government of India, United Nations Children's Fund. India Coverage Evaluation Survey Data 2009-2010. [Data shared for this analysis]
Ministry of Statistics and Programme Implementation, Government of India. Employment and Unemployment Survey, National Sample Survey Round 38, January-December 1983. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Survey on Participation in Education, National Sample Survey Round 42, July 1986- June 1987. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Employment and Unemployment Survey, National Sample Survey Round 43, July 1987- June 1988. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Survey on Literacy and Culture, National Sample Survey Round 47, July - December 1991. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Employment and Unemployment Survey, National Sample Survey Round 50, July 1993- June 1994. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Survey on Healthcare, National Sample Survey Round 52, July 1995-June 1996. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Common Property Resources, Sanitation & Hygiene Services, National Sample Survey Round 54, January - June 1998. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Employment and Unemployment Survey, India National Sample Survey Round 55, July 1999-June 2000. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Consumer Expenditure Survey, National Sample Survey Round 60, January-June 2004. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Employment and Unemployment Survey, National Sample Survey Round 61, July 2004- June 2005. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. Consumer Expenditure Survey, National Sample Survey Round 64, July 2007- June 2008. New Delhi, India: Ministry of Statistics and Programme Implementation.

Ministry of Statistics and Programme Implementation, Government of India. Household expenditure on services and durable goods survey, National Sample Survey Round 72, July 2014 - June 2015. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. India State Series: Gross State Domestic Product at Factor Cost by Industry of Origin Tables 1980-81 to 1993-94. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. India State Series: Gross State Domestic Product at Factor Cost by Industry of Origin Tables 1993-94 to 2004-05. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Statistics and Programme Implementation, Government of India. India State Series: Gross State Domestic Product at Factor Cost by Industry of Origin Tables 2004-05 to 2013-14. New Delhi, India: Ministry of Statistics and Programme Implementation.
Ministry of Women and Child Development, Government of India, United Nations Children's Fund. India Rapid Survey on Children 2013-2014. New Delhi, India: Ministry of Women and Child Development.
Mony PK, Varghese B, Thomas T. Estimation of perinatal mortality rate for institutional births in Rajasthan state, India, using capture-recapture technique. <i>BMJ Open</i> . 2015; 5(3): e005966.
Nadiger HA, Krishnamachari KA, Naidu AN, Rao BS, Srikantia SG. The use of common salt (sodium chloride) fortified with iron to control anaemia: results of a preliminary study. <i>Br J Nutr</i> . 1980; 43(1): 45-51.
Nallam NR, Paul I, Gnanamani G. Anemia and hypoalbuminuria as an adjunct to soil-transmitted helminthiasis among slum school children in Visakhapatnam, South India. <i>Asia Pac J Clin Nutr</i> . 1998; 7(2): 164-9.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural First Repeat Survey of Diet and Nutritional Status Report 1988-1990. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural First Repeat Survey of Diet and Nutritional Status Data 1988-1990. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Survey of Diet and Nutritional Status Report 1990-1992. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Survey of Diet and Nutritional Status Data 1990-1992. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Survey of Diet and Nutritional Status Report 1991-1992. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Survey of Diet and Nutritional Status Data 1991-1992. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition, Indian Council of Medical Research. India National Nutrition Monitoring Bureau Eight States Pooled Data 1991-1992. [Data shared for this analysis]

National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Urban Slums Survey of Diet and Nutritional Status Report 1993-1994. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Urban Slums Survey of Diet and Nutritional Status Data 1993-1994. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Report 1994-1995. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Data 1994-1995. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Second Repeat Survey of Diet and Nutritional Status Report 1996-1997. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Second Repeat Survey of Diet and Nutritional Status Data 1996-1997. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Tribal First Repeat Survey of Diet and Nutritional Status Report 1998-1999. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Tribal First Repeat Survey of Diet and Nutritional Status Data 1998-1999. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Report 2000-2001. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Data 2000-2001. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Micronutrient Deficiency Survey Report 2003-2006. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Micronutrient Deficiency Survey Data 2003-2006. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Report 2004-2006. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Survey of Diet and Nutritional Status Data 2004-2006. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Tribal Second Repeat Survey of Diet and Nutritional Status Report 2007-2008. Hyderabad, India: NIN.

National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Tribal Second Repeat Survey of Diet and Nutritional Status Data 2007-2008. [Data shared for this analysis]
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Third Repeat Survey of Diet and Nutritional Status Report 2011-2012. Hyderabad, India: NIN.
National Nutrition Monitoring Bureau, National Institute of Nutrition (NIN), Indian Council of Medical Research. India Rural Third Repeat Survey of Diet and Nutritional Status Data 2011-2012. [Data shared for this analysis]
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Report 2010-2011. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Data 2010-2011. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Report 2011-2012. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Data 2011-2012. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Report 2012-2013. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Data 2012-2013. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Data 2010-2013. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Annual Health Survey Report 2010-2013. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Census 1981. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Medical Certification of Cause of Death State-Level Tabulations 2000. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Medical Certification of Cause of Death State-Level Tabulations 2001. New Delhi, India: Office of the Registrar General & Census Commissioner.
Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India Medical Certification of Cause of Death State-Level Tabulations 2002. New Delhi, India: Office of the Registrar General & Census Commissioner.

Office of the Registrar General & Census Commissioner, Ministry of Home Affairs, Government of India. India SRS Verbal Autopsy 2010-2013. New Delhi, India: Office of the Registrar General and Census Commissioner. [Data shared for this analysis]

Office of the Registrar General and Census Commissioner, Ministry of Health and Family Welfare, Government of India, National Institute of Health and Family Welfare, Nutrition Foundation of India, National Institute of Nutrition, Indian Council of Medical Research. India Clinical, Anthropometric and Bio-chemical (CAB) Survey Data 2014 [Biomarker Component of Annual Health Survey]. New Delhi, India: Office of the Registrar General and Census Commissioner.

Office of the Registrar General and Census Commissioner, Ministry of Health and Family Welfare, Government of India, National Institute of Health and Family Welfare, Nutrition Foundation of India, National Institute of Nutrition, Indian Council of Medical Research. India Clinical, Anthropometric and Bio-chemical (CAB) Survey Report 2014 [Biomarker Component of Annual Health Survey]. New Delhi, India: Office of the Registrar General and Census Commissioner.

Olofin I, McDonald CM, Ezzati M, Flaxman S, Black RE, Fawzi WW, Caulfield LE, Danaei G; Nutrition Impact Model Study (anthropometry cohort pooling). Associations of suboptimal growth with all-cause and cause-specific mortality in children under five years: a pooled analysis of ten prospective studies. *PLoS One*. 2013; 8(5): e64636.

Panigrahi S, Patra PK, Khodiar PK. Neonatal screening of sickle cell anemia: a preliminary report. *Indian J Pediatr*. 2012; 79(6): 747–50.

Pasricha S-R, Black J, Muthayya S, Shet A, Bhat V, Nagaraj S, Prashanth NS, Sudarshan H, Biggs B-A, Shet AS. Determinants of anemia among young children in rural India. *Pediatrics*. 2010; 126(1): e140-9.

Pathak P, Singh P, Kapil U, Raghuvanshi RS. Prevalence of iron, vitamin A, and iodine deficiencies amongst adolescent pregnant mothers. *Indian J Pediatr*. 2003; 70(4): 299-301.

Pathak P, Tandon M, Kapil U, Singh C. Prevalence of iron deficiency anemia amongst pregnant women in urban slum communities of Delhi. *Indian Pediatr*. 1999; 36(3): 322-3.

Prinjha S, Thakur JS, Bhatia SS. Pilot testing of WHO child growth standards in Chandigarh: implications for India's child health programmes. *Bull World Health Organ*. 2009; 87(2): 116-22.

Purwar M, Ughade S, Bhagat B, Agarwal V, Kulkarni H. Bacterial vaginosis in early pregnancy and adverse pregnancy outcome. *J Obstet Gynaecol Res*. 2001; 27(4): 175-81.

Rahmathullah L, Tielsch JM, Thulasiraj RD, Katz J, Coles C, Devi S, John R, Prakash K, Sadanand AV, Edwin N, Kamaraj C. Impact of supplementing newborn infants with vitamin A on early infant mortality: community based randomised trial in southern India. *BMJ*. 2003; 327(7409): 254.

Rahmathullah L, Underwood BA, Thulasiraj RD, Milton RC, Ramaswamy K, Rahmathullah R, Babu G. Reduced mortality among children in southern India receiving a small weekly dose of vitamin A. *N Engl J Med*. 1990; 323(14): 929-35. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. *Cochrane Database Syst Rev*. 2010; CD008524.

Rajaratnam J, Abel R, Asokan JS, Jonathan P. Prevalence of anemia among adolescent girls of rural Tamilnadu. *Indian Pediatr*. 2000; 37(5): 532-6.

Rajaratnam J, Abel R, Ganesan C, Jayaseelan SA. Maternal anaemia: a persistent problem in rural Tamil Nadu. *Natl Med J India*. 2000; 13(5): 242-5.

Ramakrishnan U, Latham MC, Abel R, Frongillo EA Jr. Vitamin A supplementation and morbidity among preschool children in south India. *Am J Clin Nutr*. 1995; 61(6): 1295-303.

Raman L, Subbalaxmi P, Vasumathi N, Rawal A, Vasanthi G, Parvathi C, Adinarayana K, Pawashe A, Rao K. Iron and folic acid nutritional status of women in slum. *Nutr Report Int*. 1989; 73-81.

<p>Randomised trial to assess benefits and safety of vitamin A supplementation linked to immunisation in early infancy. WHO/CHD Immunisation-Linked Vitamin A Supplementation Study Group. Lancet. 1998; 352(9136): 1257-63.</p>
<p>Rao KV, Radhaiah G, Raju SV. Association of growth status and the prevalence of anaemia in preschool children. Indian J Med Res. 1980; 71: 237-46.</p>
<p>Rawat CM, Garg SK, Singh JV, Bhatnagar M, Chopra H, Bajpai SK. Prevalence of anaemia among adolescent girls in rural area of District Meerut, U.P. Indian J Public Health. 2001; 45(1): 24-6.</p>
<p>Ray SK, Mallick S, Kumar S, Biswas B. A rapid assessment of anaemia in pregnancy in West Bengal with special reference to care seeking behaviour of mothers. Indian J Public Health. 2000; 44(2): 58-64.</p>
<p>Reddaiah VP, Kapoor SK. Socio-biological factors in underfive deaths in a rural area. Indian J Pediatr. 1992; 59(5): 567-71.</p>
<p>Reddaiah VP, Raj PP, Ramachandran K, Nath LM, Sood SK, Madan N, Rusia U. Supplementary iron dose in pregnancy anemia prophylaxis. Indian J Pediatr. 1989; 56(1): 109-14.</p>
<p>Rusia U, Gupta S, Agarwal N, Singh K, Sikka M, Madan N. Efficacy of the new program of iron supplementation in pregnancy in India. Indian J Hematol Blood Transfus. 1999; 17(4): 87-91.</p>
<p>Rusia U, Madan N, Agarwal N, Sikka M, Sood SK. Effect of maternal iron deficiency anaemia on foetal outcome. Indian J Pathol Microbiol. 1995; 38(3): 273-9.</p>
<p>Sachdeva S, Alam S, Beig FK, Khan Z, Khalique N. Determinants of vitamin A deficiency amongst children in Aligarh District, Uttar Pradesh. Indian Pediatr. 2011; 48(11): 861-6.</p>
<p>Sachdev HPS, Osmond C, Fall CHD, Lakshmy R, Ramji S, Dey Biswas SK, Prabhakaran D, Tandon N, Reddy KS, Barker DJP, Bhargava SK. Predicting adult metabolic syndrome from childhood body mass index: follow-up of the New Delhi birth cohort. Arch Dis Child. 2009; 94(10): 768-74.</p>
<p>Saleem S, McClure EM, Goudar SS, Patel A, Esamai F, Garces A, Chomba E, Althabe F, Moore J, Kodkany B, Pasha O, Belizan J, Mayansyan A, Derman RJ, Hibberd PL, Liechty EA, Krebs NF, Hambidge KM, Buekens P, Carlo WA, Wright LL, Koso-Thomas M, Jobe AH, Goldenberg RL, on behalf of the Global Network Maternal Newborn Health Registry Study Investigators. A prospective study of maternal, fetal and neonatal deaths in low- and middle-income countries. Bull World Health Organ. 2014; 92(8): 605-12.</p>
<p>Sarin A. Severe anemia of pregnancy, recent experience. Int J Gynaecol Obstet. 1995; 50(Suppl 2): 45-49.</p>
<p>Saxena N, Nayar D, Kapil U. Prevalence of underweight, stunting and wasting. Indian Pediatr. 1997; 34(7): 627-31.</p>
<p>Scott W, Mathew NT. A Development Monitoring Service at the Local Level: Monitoring Change in Kerala: The First Five Years. Geneva, Switzerland: United Nations Research Institute for Social Development; 1985.</p>
<p>Sehgal A, Telang S, Passah SM, Jyothi MC. Maternal and Neonatal Profile and Immediate Outcome in Extremely Low Birth Weight Babies in Delhi. Trop Doct. 2004; 34(3): 165-8.</p>
<p>Sethi V, Goindi G, Kapil U. Prevalence of anemia amongst primary school age children (6-11 years) in National Capital Territory of Delhi. Indian J Pediatr. 2003; 70(6): 519-20.</p>
<p>Shah SN, Bakash A, Rauf A, Muzzafar A, Zuthshi ML. Incidence of iron deficiency anaemia in rural population of Kashmir. Indian J Public Health. 1982; 26(3): 144-54.</p>
<p>Sharma A, Prasad K, Rao KV. Identification of an appropriate strategy to control anemia in adolescent girls of poor communities. Indian Pediatr. 2000; 37(3): 261-7.</p>

Sharma JB, Soni D, Murthy NS, Malhotra M. Effect of dietary habits on prevalence of anemia in pregnant women of Delhi. <i>J Obstet Gynaecol Res.</i> 2003; 29(2): 73-8.
Shobha S, Sharada D. Efficacy of twice weekly iron supplementation in anemic adolescent girls. <i>Indian Pediatr.</i> 2003; 40(12): 1186-90.
Sidhu S, Kumari K, Uppal M. Prevalence of anemia in Schedule Caste preschool children of Punjab. <i>Indian J Med Sci.</i> 2002; 56(5): 218-21.
Sidhu S. Incidence of anaemia among scheduled caste pre-school children of Punjab. <i>Indian J Matern Child Health.</i> 1996; 7(3): 76-7.
Singhal PK, Mathur GP, Mathur S, Singh YD. Mortality patterns in under six children in I.C.D.S. urban slum. <i>Indian Pediatr.</i> 1986; 23(8): 617-22.
Singla PN, Agarwal KN. Studies on normal hemoglobin and hematocrit values in healthy children based on hematinic supplementation. <i>Indian Pediatr.</i> 1981; 18(11): 821-5.
Sivakumar B, Brahmam GN, Madhavan Nair K, Ranganathan S, Vishnuvardhan Rao M, Vijayaraghavan K, Krishnaswamy K. Prospects of fortification of salt with iron and iodine. <i>Br J Nutr.</i> 2001; 167-173.
Smith JL, Brooker S. Impact of hookworm infection and deworming on anaemia in non-pregnant populations: a systematic review. <i>Trop Med Int Health.</i> 2010; 15(7): 776-95.
Sood SK, Ramachandran K, Rani K, Ramalingaswami V, Mathan VI, Ponniah J, Baker SJ. WHO sponsored collaborative studies on nutritional anaemia in India. The effect of parenteral iron administration in the control of anaemia of pregnancy. <i>Br J Nutr.</i> 1979; 42(3): 399-406.
Spector JM, Agrawal P, Kodkany B, Lipsitz S, Lashoher A, Dziekan G, Bahl R, Merialdi M, Mathai M, Lemer C, Gawande A. Improving Quality of Care for Maternal and Newborn Health: Prospective Pilot Study of the WHO Safe Childbirth Checklist Program. <i>PLoS One.</i> 2012; 7(5): e35151.
State Bureau of Health Intelligence & Vital Statistics, Directorate of Health Services, Department of Health & Family Welfare, Government of Odisha . India - Odisha Medical Certification of Cause of Death Data 2009. Odisha, India: Department of Health & Family Welfare, Government of Odisha.
State Bureau of Health Intelligence & Vital Statistics, Directorate of Health Services, Department of Health & Family Welfare, Government of Odisha . India - Odisha Medical Certification of Cause of Death Data 2010. Odisha, India: Department of Health & Family Welfare, Government of Odisha
State Bureau of Health Intelligence & Vital Statistics, Directorate of Health Services, Department of Health & Family Welfare, Government of Odisha . India - Odisha Medical Certification of Cause of Death Data 2011. Odisha, India: Department of Health & Family Welfare, Government of Odisha.
State Bureau of Health Intelligence & Vital Statistics, Directorate of Health Services, Department of Health & Family Welfare, Government of Odisha . India - Odisha Medical Certification of Cause of Death Data 2012. Odisha, India: Department of Health & Family Welfare, Government of Odisha.
State Bureau of Health Intelligence & Vital Statistics, Directorate of Health Services, Department of Health & Family Welfare, Government of Odisha . India - Odisha Medical Certification of Cause of Death Data 2013. Odisha, India: Department of Health & Family Welfare, Government of Odisha.
Straughn HK, Goldenberg RL, Tolosa JE, Daly S, de Codes J, Festin MR, Limpongsanurak S, Lumbiganon P, Paul VK, Peedicayil A, Purwar M, Sabogal JC, Shenoy S. Birthweight-specific neonatal mortality in developing countries and obstetric practices. <i>Int J Gynaecol Obstet.</i> 2003; 80(1): 71-8.
Swain S, Singh S, Bhatia BD, Pandey S, Krishna M. Maternal hemoglobin and serum albumin and fetal growth. <i>Indian Pediatr.</i> 1994; 31(7): 777-82.
Swaminathan M, Apte S, Someswara Rao K. Nutrition of the people of Ankola Taluk N. Kanara. <i>Indian J Med Res.</i> 1960; 762-74.

Talsania NJ, Lala MK. Evaluation of antenatal risk scoring in a preterm birth prevention and perinatal loss. Indian J Matern Child Health. 1994; 5(1): 5-9.
Thangaleela T, Vijayalakshmi P. Prevalence of Anaemia in Pregnancy. Indian J Nutr Diet. 1994 ; 31(2):26-9.
Thilothammal N, Sujaritha R, Banu K, Ratnam S, Ezhlarasi. The nutritional status of south Indian women in child bearing age and its influence on birth weight and maturity of offspring. J Obstet Gynaecol India. 1993; 871-7.
Thomas SV, Sindhu K, Ajaykumar B, Sulekha Devi PB, Sujamol J. Maternal and obstetric outcome of women with epilepsy. Seizure. 2009; 18(3): 163-6.
Tielsch JM, Rahmathullah L, Thulasiraj RD, Katz J, Coles C, Sheeladevi S, John R, Prakash K. Newborn vitamin A dosing reduces the case fatality but not incidence of common childhood morbidities in South India. J Nutr. 2007; 137(11): 2470-4.
Trivedi SS, Pasrija S. Teenage pregnancies and their obstetric outcomes. Trop Doct. 2007; 37(2): 85-8.
Uberoi IS, DeSweemer C, Taylor CE. A study of anemia among rural Punjabi children. Indian J Med Res. 1972; 60(5): 793-9.
Upadhyay RP, Rai SK, Krishnan A. Using three delays model to understand the social factors responsible for neonatal deaths in rural Haryana, India. J Trop Pediatr. 2013; 59(2): 100-5.
Venkatarao T, Ramakrishnan R, Nair NG, Radhakrishnan S, Sundaramoorthy L, Koya PK, Kumar SK. Effect of vitamin A supplementation to mother and infant on morbidity in infancy. Indian Pediatr. 1996; 33(4): 279-86. as it appears in Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. Cochrane Database Syst Rev. 2010; CD008524.
Verma M, Chhatwal J, Kaur G. Prevalence of anemia among urban school children of Punjab. Indian Pediatr. 1998; 35(12): 1181-6.
Vijayaraghavan K, Rao DH. Diet & nutrition situation in rural India. Indian J Med Res. 1998; 108:243-53.
Vogel JP, Souza JP, Mori R, Morisaki N, Lumbiganon P, Laopaiboon M, Ortiz-Panozo E, Hernandez B, Pérez-Cuevas R, Roy M, Mittal S, Cecatti JG, Tunçalp Ö, Gülmezoglu AM; WHO Multicountry Survey on Maternal and Newborn Health Research Network. Maternal complications and perinatal mortality: findings of the World Health Organization Multicountry Survey on Maternal and Newborn Health. BJOG. 2014; 121(Supp 1): 76-88.
Working Group on Fortification of Salt with Iron. Use of common salt fortified with iron in the control and prevention of anemia--a collaborative study. Report of the Working Group on Fortification of Salt with Iron. Am J Clin Nutr. 1982; 35(6): 1442-51.
World Bank. Survey of Living Conditions 1997-1998, Uttar Pradesh and Bihar. Washington D.C., United States: World Bank.
World Bank. World Development Indicators - Vitamin A Supplementation Coverage Rate. Washington D.C., United States: World Bank.
World Health Organization (WHO), United Nations Children's Fund (UNICEF). WHO and UNICEF Reported Administrative Data for Immunization Coverage Time Series. Available from: http://www.who.int/immunization/monitoring_surveillance/data/en/
World Health Organization (WHO). Global Database on Child Growth and Malnutrition - Historical. Geneva, Switzerland: WHO.
World Health Organization (WHO). WHO Global Database on Anemia, Nutrition Landscape Information System. Geneva, Switzerland: WHO.

World Health Organization (WHO). WHO Global Database on Vitamin A Deficiency. Geneva, Switzerland: WHO.

World Health Organization (WHO). WHO Global Survey on Maternal and Perinatal Health 2004-2008. Geneva, Switzerland: WHO.

World Health Organization. Study on Global Ageing and Adult Health (SAGE) Pilot Study 2005 Data from the Data Archive of Social Research on Aging. Los Altos, United States: Sociometrics Corporation. Available from: <http://home.socio.com/age2728.php>

Yusufji, Mathan VI, Baker SJ. Iron, folate, and vitamin B 12 nutrition in pregnancy: a study of 1000 women from southern India. Bull World Health Organ. 1973; 48(1): 15-22.

CONFIDENTIAL

Table 2: Percent contribution of under-five and all-age DALYs attributable to child and maternal malnutrition and its components in India by sex in 2016

Risk factors	Percent of under-five DALYs (95% uncertainty interval)			Percent of all-age DALYs (95% uncertainty interval)		
	Boys	Girls	Both sexes	Men	Women	Both sexes
Child and maternal malnutrition	66.8% (64.4 to 69.5)	63.5% (60.5 to 67.2)	65.2% (62.7 to 68.0)	12.9% (12.1 to 13.8)	16.6% (15.5 to 17.7)	14.6% (13.7 to 15.5)
Low birth weight and short gestation	44.1% (42.4 to 45.9)	38.3% (36.7 to 40.1)	41.3% (39.7 to 42.8)	7.4% (6.7 to 8.2)	7.2% (6.3 to 8.1)	7.3% (6.6 to 8.1)
Child growth failure	19.2% (17.4 to 21.2)	22.0% (19.4 to 24.7)	20.6% (18.7 to 22.6)	3.2% (2.9 to 3.6)	4.1% (3.5 to 4.8)	3.6% (3.2 to 4.1)
Suboptimal breastfeeding	2.7% (2.2 to 3.4)	3.0% (2.4 to 3.8)	2.9% (2.3 to 3.6)	0.5% (0.4 to 0.6)	0.6% (0.4 to 0.7)	0.5% (0.4 to 0.6)
Iron deficiency	2.9% (1.9 to 4.1)	2.5% (1.6 to 3.6)	2.7% (1.8 to 3.8)	2.2% (1.6 to 3.0)	5.1% (3.8 to 6.6)	3.6% (2.6 to 4.7)
Vitamin A deficiency	1.0% (0.6 to 1.7)	1.2% (0.7 to 1.8)	1.1% (0.6 to 1.7)	0.2% (0.1 to 0.3)	0.2% (0.1 to 0.4)	0.2% (0.1 to 0.3)
Zinc deficiency	0.4% (0.0 to 1.0)	0.5% (0.0 to 1.4)	0.4% (0.0 to 1.2)	0.1% (0.0 to 0.2)	0.1% (0.0 to 0.3)	0.1% (0.0 to 0.2)

The percent contribution of the components of child and maternal malnutrition add up to more than the overall child and maternal malnutrition risk factor due to overlap of the components. DALY=disability-adjusted life-year.

Table 3: Prevalence of low birth weight for India and states grouped by epidemiological transition level, 1990-2016

States of India grouped by ETL (population in 2016)	Prevalence						Percent change						Annualised percent change
	1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	1990-2000	2000-2016	1990-2016	1990-2000	2000-2016	1990-2016	
India (1316 million)	19.1 (18.1 to 20.1)	19.0 (17.9 to 20.1)	19.6 (18.4 to 20.8)	-0.5 (-2.5 to 1.6)	3.0 (0.6 to 5.3)	2.5 (-0.1 to 5.2)	-0.05	0.18	0.09				
Low ETL (626 million)	18.0 (16.7 to 19.3)	17.9 (16.5 to 19.2)	18.7 (17.2 to 20.2)	-0.8 (-4.2 to 2.7)	4.3 (0.5 to 8.0)	3.5 (0.0 to 7.5)	-0.08	0.26	0.13				
Bihar	16.3 (14.6 to 17.9)	15.7 (14.0 to 17.4)	16.2 (14.6 to 18.0)	-3.7 (-11.8 to 5.0)	3.7 (-5.5 to 13.5)	-0.2 (-8.5 to 8.8)	-0.38	0.23	-0.01				
Jharkhand	16.7 (14.7 to 18.5)	16.6 (14.7 to 18.4)	18.1 (16.2 to 20.0)	-0.4 (-8.3 to 7.7)	9.0 (0.1 to 18.8)	8.5 (0.3 to 17.6)	-0.04	0.54	0.31				
Uttar Pradesh	16.9 (15.2 to 18.6)	17.1 (15.3 to 18.8)	18 (16.2 to 19.9)	1.3 (-6.3 to 9.8)	5.5 (-2.7 to 14.3)	6.9 (-1.6 to 15.7)	0.13	0.34	0.26				
Rajasthan	17.6 (16.1 to 19.1)	17.7 (16.2 to 19.1)	18.7 (17.1 to 20.3)	0.4 (-6.8 to 7.4)	6.0 (-1.0 to 13.7)	6.4 (-0.8 to 13.5)	0.04	0.37	0.24				
Meghalaya	13.5 (11.8 to 15.3)	15.0 (13.3 to 16.9)	17.2 (15.3 to 19.3)	10.9 (2.3 to 21.1)	14.9 (5.7 to 24.5)	27.4 (17.4 to 39.3)	1.04	0.87	0.93				
Assam	16.3 (14.6 to 18.0)	16.5 (14.7 to 18.1)	17.8 (16.0 to 19.8)	0.8 (-7.9 to 9.6)	8.2 (-0.8 to 18.3)	9.1 (0.2 to 18.8)	0.08	0.50	0.33				
Chhattisgarh	18.3 (16.6 to 19.8)	18.3 (16.6 to 19.7)	20.5 (18.8 to 22.1)	-0.1 (-7.8 to 8.2)	12.4 (4.6 to 20.7)	12.3 (4.3 to 21.2)	-0.01	0.74	0.45				
Madhya Pradesh	23.3 (21.7 to 24.9)	22.9 (21.4 to 24.6)	23.0 (21.5 to 24.4)	-1.4 (-8.2 to 6.1)	0.2 (-6.5 to 7.5)	-1.2 (-7.9 to 5.8)	-0.14	0.01	-0.05				
Odisha	20.6 (19.2 to 22.2)	20.5 (19.1 to 22.0)	21.7 (20.2 to 23.3)	-0.5 (-7.7 to 7.6)	6.0 (-1.0 to 13.8)	5.5 (-1.9 to 14.0)	-0.05	0.36	0.21				
Lower-middle ETL (92 million)	19.1 (18.0 to 20.1)	19.4 (18.2 to 20.6)	21.3 (20.0 to 22.7)	1.6 (-2.7 to 6.3)	9.8 (5.0 to 14.8)	11.6 (7.1 to 16.6)	0.16	0.59	0.42				
Arunachal Pradesh	15.0 (13.1 to 16.8)	15.1 (13.4 to 17.0)	15.8 (13.9 to 17.8)	1.3 (-7.0 to 10.8)	4.5 (-4.0 to 13.6)	5.8 (-2.4 to 15.0)	0.13	0.27	0.22				
Mizoram	11.9 (10.8 to 13.2)	13.1 (11.8 to 14.4)	14.5 (13.0 to 16.1)	9.7 (2.4 to 18.8)	11 (1.9 to 19.2)	21.8 (11.9 to 31.0)	0.93	0.65	0.76				
Nagaland	18.0 (16.7 to 19.2)	19.1 (17.6 to 20.5)	20.8 (19.4 to 22.3)	6.6 (-0.1 to 13.9)	8.8 (2.1 to 16.6)	16.1 (9.1 to 23.8)	0.64	0.53	0.57				
Uttarakhand	16.3 (14.8 to 17.7)	16.7 (15.4 to 18.1)	19.7 (18.3 to 21.1)	2.8 (-4.6 to 10.9)	17.4 (9.7 to 25.5)	20.7 (13.3 to 29.8)	0.28	1.01	0.73				
Gujarat	20.5 (19.2 to 21.7)	20.7 (19.5 to 22.1)	22.3 (20.9 to 23.7)	1.4 (-4.4 to 7.6)	7.5 (1.7 to 13.8)	9.0 (3.5 to 15.3)	0.13	0.46	0.33				
Tripura	15.5 (14.2 to 17.0)	16.2 (14.8 to 17.7)	18.7 (17.2 to 20.3)	4.4 (-2.8 to 12.5)	15.3 (8.1 to 23.7)	20.3 (11.6 to 29.3)	0.43	0.89	0.71				
Sikkim	13.5 (12.1 to 15.2)	14.6 (13.2 to 16.3)	19.3 (17.7 to 21.1)	8 (-1.1 to 18.0)	32 (22.9 to 43.1)	42.6 (31.4 to 55.0)	0.77	1.75	1.37				
Manipur	12.9 (11.6 to 14.5)	13.1 (11.8 to 14.7)	14.2 (12.6 to 15.9)	1.5 (-6.6 to 9.2)	8.4 (-0.6 to 17.4)	10.0 (1.0 to 19.0)	0.15	0.50	0.37				
Higher-middle ETL (446 million)	21.1 (20.2 to 22.1)	21.3 (20.3 to 22.3)	21.6 (20.4 to 22.8)	0.7 (-2.2 to 3.6)	1.4 (-1.3 to 4.3)	2.1 (-1.0 to 5.0)	0.07	0.09	0.08				
Haryana	15.9 (14.3 to 17.4)	16.6 (15 to 18.1)	19.1 (17.4 to 20.9)	4.7 (-2.2 to 12.4)	15 (7.1 to 23.6)	20.4 (12.2 to 28.9)	0.46	0.88	0.72				
Delhi	17.6 (16.1 to 18.8)	18.3 (16.8 to 19.7)	18.6 (16.9 to 20.3)	4.3 (-2.9 to 12.2)	1.7 (-6.7 to 10.4)	6.1 (-2.0 to 15.1)	0.42	0.11	0.23				
Telangana	18.7 (17.3 to 20.3)	19.8 (18.3 to 21.6)	22.6 (21.2 to 24.1)	6.1 (-1.9 to 14.9)	14.1 (6.6 to 22.1)	21.0 (12.8 to 29.2)	0.59	0.83	0.74				
Andhra Pradesh	20.6 (19.3 to 22.1)	20.8 (19.5 to 22.3)	21.0 (19.6 to 22.4)	0.9 (-5.1 to 7.4)	0.7 (-5.6 to 6.9)	1.6 (-4.9 to 8.3)	0.09	0.05	0.06				
Jammu and Kashmir	12.9 (11.2 to 14.7)	13.6 (12.0 to 15.6)	15.6 (13.4 to 17.6)	5.6 (-3.5 to 15.9)	14.5 (4.0 to 25.6)	21 (10.6 to 32.5)	0.55	0.85	0.74				
Karnataka	20.3 (19.1 to 21.6)	20.9 (19.7 to 22.2)	21.4 (20.1 to 22.7)	2.9 (-3.7 to 9.5)	2.2 (-3.4 to 8.6)	5.2 (-0.8 to 11.6)	0.29	0.14	0.19				
West Bengal	18.4 (16.9 to 19.8)	18.1 (16.6 to 19.5)	18.5 (16.9 to 20.0)	-1.5 (-7.7 to 5.5)	1.9 (-4.9 to 9.0)	0.4 (-6.2 to 8.0)	-0.15	0.12	0.02				
Maharashtra	27.6 (26.2 to 29.0)	27.4 (26.0 to 28.8)	25.9 (24.6 to 27.3)	-0.9 (-6.8 to 4.9)	-5.3 (-10.5 to -0.1)	-6.2 (-11.7 to -1.0)	-0.09	-0.34	-0.24				
Union territories other than Delhi	16.8 (15.5 to 18.0)	17.5 (16.2 to 18.8)	20.0 (18.5 to 21.5)	4.4 (-1.5 to 11.1)	13.8 (6.8 to 20.9)	18.9 (11.9 to 26.1)	0.43	0.81	0.67				
High ETL (152 million)	17.5 (16.5 to 18.5)	17.5 (16.6 to 18.6)	17.8 (16.8 to 19.0)	0.0 (-3.8 to 4.0)	1.9 (-2.0 to 5.5)	1.9 (-1.8 to 5.9)	0.00	0.12	0.07				
Himachal Pradesh	14.7 (13.4 to 16.2)	15.2 (13.8 to 16.7)	17.3 (15.8 to 19.0)	3.3 (-5.2 to 11.8)	14.1 (5.2 to 23.9)	17.9 (8.8 to 27.4)	0.32	0.83	0.63				
Punjab	15.0 (13.7 to 16.5)	15.2 (13.9 to 16.7)	15.4 (14.1 to 17.0)	1.2 (-6.3 to 8.9)	1.5 (-5.4 to 8.8)	2.7 (-4.1 to 9.7)	0.12	0.09	0.10				
Tamil Nadu	20.7 (19.5 to 21.9)	20.4 (19.1 to 21.6)	20.4 (19.2 to 21.7)	-1.6 (-6.6 to 4.4)	0.3 (-5.2 to 5.6)	-1.3 (-6.8 to 4.7)	-0.16	0.02	-0.05				
Goa	20.4 (19.3 to 21.5)	20.7 (19.6 to 21.9)	21.1 (19.8 to 22.3)	1.6 (-3.4 to 6.9)	1.7 (-3.5 to 7.4)	3.3 (-2.2 to 9.6)	0.16	0.10	0.13				
Kerala	13.2 (12.2 to 14.2)	13.6 (12.7 to 14.7)	14.1 (13.1 to 15.2)	3.4 (-3.6 to 11.0)	3.3 (-3.3 to 10.3)	6.9 (-0.2 to 13.9)	0.34	0.20	0.26				

ETL=epidemiological transition level.

Table 4: Prevalence of wasting in under-five children for India and states grouped by epidemiological transition level, 1990-2016

States of India grouped by ETL	Prevalence						Percent change						Annualised percent change	
	1990 (95% uncertainty interval)		2000 (95% uncertainty interval)		2016 (95% uncertainty interval)		1990-2000 (95% uncertainty interval)		2000-2016 (95% uncertainty interval)		1990-2016 (95% uncertainty interval)			
	1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	1990-2000	2000-2016	1990-2000	2000-2016	1990-2016	1990-2016		
India (1316 million)	21.6 (20.7 to 22.5)	19.9 (19.4 to 20.3)	15.9 (15.6 to 16.2)	-8.2 (-11.7 to -4.3)	-19.9 (-22.2 to -17.5)	-26.5 (-29.8 to -22.9)	-0.85	-1.38	-1.18	-1.38	-1.18	-1.18		
Low ETL (626 million)	21.7 (20.2 to 23.3)	20.2 (19.5 to 20.9)	15.1 (14.6 to 15.6)	-6.7 (-13.0 to 0.2)	-25.3 (-29 to 21.6)	-30.3 (-35.5 to -24.3)	-0.69	-1.81	-1.38	-1.38	-1.38	-1.38		
Bihar	19.7 (15.9 to 23.6)	20.7 (18.5 to 22.9)	14.1 (13.0 to 15.2)	4.9 (-14.0 to 27.2)	-31.7 (-40.2 to -21.9)	-28.4 (-41.7 to -10.7)	0.48	-2.36	-1.28	-1.28	-1.28	-1.28		
Jharkhand	39.5 (34.5 to 44.6)	37.7 (34.4 to 41.5)	16.2 (15.0 to 17.5)	-4.7 (-15.0 to 8.2)	-57 (-62.1 to -51.5)	-59 (-64.7 to -52.2)	-0.48	-5.14	-3.37	-3.37	-3.37	-3.37		
Uttar Pradesh	17.8 (14.4 to 21.6)	15.8 (14.5 to 17.2)	13.5 (12.2 to 14.6)	-11.4 (-26.7 to 9.2)	-14.7 (-25.1 to -3.5)	-24.4 (-38.8 to -6.2)	-1.21	-0.99	-1.07	-1.07	-1.07	-1.07		
Rajasthan	19.5 (15.9 to 23.2)	17.5 (16.0 to 19.0)	17.2 (15.9 to 18.4)	-10.6 (-25.1 to 7.9)	-1.5 (-11.6 to 9.4)	-12.0 (-26.5 to 9.4)	-1.12	-0.10	-0.49	-0.49	-0.49	-0.49		
Meghalaya	17.0 (13.4 to 21.3)	18.3 (16.1 to 20.8)	12.8 (11.9 to 13.8)	7.6 (-14.1 to 33.5)	-29.7 (-39.8 to -18.2)	-24.3 (-39.8 to -2.7)	0.74	-2.18	-1.07	-1.07	-1.07	-1.07		
Assam	21.3 (18.2 to 24.4)	17.2 (15.6 to 18.7)	15.5 (14.0 to 16.9)	-19.2 (-32.8 to -1.2)	-9.8 (-21.8 to 3.5)	-27.1 (-38.7 to -12.9)	-2.11	-0.64	-1.21	-1.21	-1.21	-1.21		
Chhattisgarh	24.7 (19.9 to 29.3)	21.1 (18.2 to 23.8)	17.0 (15.7 to 18.4)	-14.4 (-27.4 to 1.3)	-19.5 (-30.2 to 4.5)	-31.1 (-43.0 to -14.0)	-1.55	-1.35	-1.42	-1.42	-1.42	-1.42		
Madhya Pradesh	23.9 (20.3 to 27.7)	25.4 (23.8 to 27.0)	16.9 (15.8 to 18.0)	6.5 (-8.2 to 25.1)	-33.5 (-39.4 to -27.5)	-29.2 (-39.8 to -15.8)	0.63	-2.52	-1.32	-1.32	-1.32	-1.32		
Odisha	24.8 (20.9 to 28.7)	23.7 (22.1 to 25.2)	17.9 (16.7 to 19.0)	-4.4 (-17.5 to 12.9)	-24.5 (-31.4 to -17.1)	-27.9 (-38.4 to -13.5)	-0.45	-1.74	-1.25	-1.25	-1.25	-1.25		
Lower-middle ETL (92 million)	21.5 (18.9 to 24.2)	19.4 (18.2 to 20.6)	17.1 (16.2 to 18.0)	-9.7 (-19.7 to 1.9)	-11.8 (-18.7 to -4.4)	-20.3 (-29.7 to -8.6)	-1.01	-0.78	-0.87	-0.87	-0.87	-0.87		
Arunachal Pradesh	10.0 (7.4 to 13.1)	10.7 (9.0 to 12.6)	16.9 (15.8 to 17.9)	7.0 (-17.4 to 39.8)	58.2 (32.8 to 92.3)	69.3 (27.9 to 130.5)	0.688	2.91	2.05	2.05	2.05	2.05		
Mizoram	12.7 (10.4 to 15.4)	11.8 (10.2 to 13.5)	14.8 (13.7 to 15.7)	-7.3 (-23.6 to 11.8)	25.6 (7.7 to 46.8)	16.4 (-5.6 to 44.3)	-0.75	1.43	0.59	0.59	0.59	0.59		
Nagaland	11.3 (8.7 to 14.3)	11.7 (10.1 to 13.6)	11.7 (10.8 to 12.5)	4.0 (-18.4 to 32.0)	0.1 (-14.4 to 17.6)	4.1 (-18.1 to 36.0)	0.40	0.01	0.16	0.16	0.16	0.16		
Uttarakhand	23.3 (18.6 to 27.8)	21.0 (18.1 to 24.0)	11.3 (10.3 to 12.2)	-9.6 (-24.3 to 8.8)	-46.5 (-54.4 to -36.7)	-51.6 (-60.1 to -38.8)	-1.01	-3.83	-2.75	-2.75	-2.75	-2.75		
Gujarat	22.6 (19.2 to 26.0)	20.5 (19.0 to 22.1)	18.8 (17.5 to 19.9)	-9.1 (-21.1 to 5.5)	-8.4 (-17.1 to 1.1)	-16.8 (-28.7 to -0.4)	-0.95	-0.55	-0.70	-0.70	-0.70	-0.70		
Tripura	21.4 (17.0 to 26.0)	20.8 (18.3 to 23.5)	16.9 (15.4 to 18.3)	-2.6 (-18.6 to 19.8)	-19.1 (-30.9 to -4.8)	-21.2 (-36.6 to 0.1)	-0.26	-1.31	-0.91	-0.91	-0.91	-0.91		
Sikkim	8.3 (6.3 to 10.9)	7.1 (5.8 to 8.6)	5.5 (4.9 to 6.1)	-15.5 (-34.3 to 9.7)	-22.2 (-38.7 to -1.8)	-34.2 (-50.8 to -12.5)	-1.66	-1.55	-1.60	-1.60	-1.60	-1.60		
Manipur	10.0 (7.8 to 12.7)	9.4 (8.0 to 10.9)	7.8 (7.0 to 8.6)	-6.0 (-23.9 to 17.4)	-17.3 (-31.6 to 0.8)	-22.3 (-39.5 to 1.1)	-0.62	-1.18	-0.96	-0.96	-0.96	-0.96		
Higher-middle ETL (446 million)	22.6 (21.2 to 24.1)	20.6 (19.9 to 21.2)	17.1 (16.6 to 17.6)	-9.1 (-14.9 to -2.9)	-16.9 (-20.1 to -13.4)	-24.5 (-29.1 to -18.8)	-0.95	-1.15	-1.07	-1.07	-1.07	-1.07		
Haryana	17.9 (15.4 to 20.9)	13.5 (12.3 to 14.8)	15.3 (14.0 to 16.6)	-24.9 (-38.1 to -9.0)	13.6 (-0.1 to 29.7)	-14.7 (-28.3 to 1.8)	-2.82	0.80	-0.61	-0.61	-0.61	-0.61		
Delhi	25.3 (22.3 to 28.3)	19.1 (17.1 to 21.0)	14.3 (12.8 to 15.8)	-24.7 (-36.9 to -10.5)	-25.1 (-34.8 to -13.1)	-43.6 (-51.2 to -34.4)	-2.79	-1.79	-2.18	-2.18	-2.18	-2.18		
Telangana	22.3 (18.9 to 25.9)	21.7 (19.8 to 23.6)	17.9 (16.6 to 19)	-3.0 (-16.1 to 13.5)	-17.4 (-26.8 to -7.4)	-19.9 (-32.5 to -4.6)	-0.30	-1.19	-0.85	-0.85	-0.85	-0.85		
Andhra Pradesh	18.9 (16.0 to 21.9)	17.0 (15.7 to 18.5)	17.9 (16.9 to 18.9)	-10.0 (-23.5 to 6.8)	4.9 (-4.8 to 16.1)	-5.7 (-19.6 to 11.7)	-1.05	0.30	-0.22	-0.22	-0.22	-0.22		
Jammu and Kashmir	17.5 (14.1 to 21.1)	14.3 (12.8 to 16.1)	6.9 (6.3 to 7.5)	-18.0 (-32.3 to -0.5)	-51.6 (-58.6 to -44.1)	-60.3 (-67.8 to 50.1)	-1.97	-4.44	-3.50	-3.50	-3.50	-3.50		
Karnataka	23.2 (19.8 to 27.1)	20.8 (19.2 to 22.5)	17.6 (16.6 to 18.7)	-10.3 (-23.3 to 4.3)	-15.3 (-23.4 to -6.4)	-24.1 (-35.1 to -9.7)	-1.09	-1.03	-1.05	-1.05	-1.05	-1.05		
West Bengal	21.9 (18.2 to 25.7)	20.3 (18.7 to 21.8)	15.5 (14.4 to 16.6)	-7.3 (-21.2 to 10.0)	-23.6 (-31.1 to -14.8)	-29.2 (-40.5 to -13.9)	-0.75	-1.67	-1.32	-1.32	-1.32	-1.32		
Maharashtra	26.2 (22.8 to 29.4)	24.6 (23.2 to 26)	19.5 (18.4 to 20.6)	-5.9 (-16.2 to 7.6)	4.9 (-4.8 to 16.1)	-5.7 (-19.6 to 11.7)	-1.05	-1.44	-1.12	-1.12	-1.12	-1.12		
Union territories other than Delhi	14.2 (11.6 to 17.3)	12.6 (10.2 to 15.2)	12.2 (9.5 to 15.8)	-11.2 (-27.5 to 9.3)	-1.5 (-25.4 to 30.1)	-12.6 (-35.8 to 15.9)	-1.18	-0.22	-0.59	-0.59	-0.59	-0.59		
High ETL (152 million)	19.1 (17.4 to 20.8)	16.6 (15.7 to 17.4)	16.2 (15.6 to 16.8)	-13.3 (-21.2 to -4.1)	-1.8 (-7.7 to 4.4)	-14.9 (-23.3 to -5.4)	-1.41	-0.11	-0.62	-0.62	-0.62	-0.62		
Himachal Pradesh	15.1 (12.3 to 18.0)	14.7 (13.0 to 16.5)	11.0 (9.5 to 12.6)	-2.8 (-21.2 to 21.1)	-25.6 (-38.3 to -10.3)	-27.6 (-33.0 to -8.0)	-0.28	-1.83	-1.24	-1.24	-1.24	-1.24		
Punjab	13.7 (11.6 to 16.0)	10.0 (8.8 to 11.0)	11.4 (10.4 to 12.4)	-27.1 (-40.4 to -9.2)	14.4 (-6.6 to 31.5)	-16.6 (-30.4 to -0.3)	-3.11	0.84	-0.70	-0.70	-0.70	-0.70		
Tamil Nadu	23.9 (20.8 to 27.3)	21.0 (19.7 to 22.5)	18.8 (17.8 to 19.8)	-12.1 (-23.0 to 1.0)	-10.5 (-18.0 to -2.8)	-21.3 (-31.8 to -9.0)	-1.28	-0.69	-0.92	-0.92	-0.92	-0.92		
Goa	18.5 (15.5 to 21.6)	15.0 (13.1 to 16.9)	15.3 (14.1 to 16.3)	-19.1 (-30.5 to -3.9)	1.9 (-12.2 to 18.8)	-17.6 (-30.0 to 0.0)	-2.10	0.12	-0.74	-0.74	-0.74	-0.74		
Kerala	14.0 (11.3 to 16.9)	12.8 (11.4 to 14.3)	15.7 (14.7 to 16.8)	-8.5 (-24.5 to 12.9)	22.2 (7.5 to 39.8)	11.8 (-8.4 to 40.0)	-0.88	1.26	0.43	0.43	0.43	0.43		

ETL=epidemiological transition level.

Table 5: Prevalence of stunting in under-five children for India and states grouped by epidemiological transition level, 1990-2016

States of India grouped by ETI	Prevalence			Percent change			Annualised percent change
	1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	
India (1316 million)	54.1 (52.5 to 55.9)	51.8 (50.4 to 53.2)	42.1 (40.5 to 43.7)	-4.4 (-8.4 to -0.8)	-18.6 (-22.2 to -14.7)	-22.2 (-26.1 to -18.1)	-0.45
Low ETI (626 million)	57.9 (55.3 to 60.9)	56.8 (54.8 to 59.2)	49.4 (46.8 to 52.0)	-1.9 (-7.5 to 4.5)	-13.1 (-18.6 to 7.4)	-14.7 (-21.1 to -8.5)	-0.19
Bihar	58.0 (52.4 to 66.5)	58.0 (52.6 to 63.9)	55.3 (49.4 to 62.5)	-1.0 (-14.3 to 13.3)	-4.7 (-18.2 to 12.4)	-5.7 (-20.0 to 12.4)	-0.10
Jharkhand	54.1 (47.7 to 61.7)	52.2 (45.8 to 59.9)	49.8 (44.7 to 55.9)	-3.5 (-19.4 to 15.6)	-4.7 (-20.2 to 13.8)	-8.0 (-22.1 to 8.6)	-0.35
Uttar Pradesh	59.3 (53.2 to 66.9)	58.5 (53.9 to 63.9)	55.2 (50.2 to 60.7)	-1.4 (-14.4 to 13.1)	-5.5 (-17.0 to 7.3)	-6.9 (-19.5 to 8.8)	-0.14
Rajasthan	56.3 (50.4 to 63.4)	54.5 (50.6 to 59.2)	38.2 (33.5 to 42.6)	-3.2 (-15.9 to 10.6)	-29.9 (-39.6 to -20.1)	-32.1 (-42.9 to -21)	-0.32
Meghalaya	50.3 (44.1 to 57.9)	50.3 (45.5 to 55.9)	46.1 (41.5 to 51.3)	-0.4 (-15.9 to 18.0)	-8.4 (-21.7 to 4.9)	-8.8 (-23.3 to 7.8)	-0.04
Assam	56.4 (50.4 to 64)	54.6 (49.7 to 60.1)	40.2 (33.6 to 46.3)	-3.2 (-16.9 to 11.7)	-26.5 (-39.7 to -13.3)	-28.8 (-42.4 to 14.6)	-0.33
Chhattisgarh	56.6 (50.7 to 64.8)	54.4 (47.8 to 62.7)	39.7 (33.8 to 47.1)	-3.9 (-19.1 to 14.0)	-27.1 (-41.7 to 9.5)	-29.8 (-43.4 to -13.7)	-0.40
Madhya Pradesh	59.0 (52.7 to 66.1)	58.5 (54.3 to 63.4)	46.0 (40.6 to 51.9)	-0.9 (-13.8 to 13.3)	-21.3 (-32.1 to 9.3)	-22.0 (-33.6 to -7.4)	-0.09
Odisha	57.1 (50.5 to 64.5)	54.4 (49.3 to 60.9)	39.5 (34.8 to 45.0)	-4.7 (-18.9 to 11.3)	-27.4 (-38.5 to -13.6)	-30.8 (-41.8 to -17.2)	-0.48
Lower-middle ETI (92 million)	55.1 (51.1 to 60.2)	52.1 (48.9 to 55.7)	39.1 (35.2 to 43.4)	-5.4 (-14.5 to 4.8)	-25.0 (-33.7 to -15.5)	-29.1 (-38.1 to -19.4)	-0.56
Arunachal Pradesh	40.4 (32.7 to 48.1)	36.4 (31.7 to 43.2)	31.0 (27.3 to 34.9)	-9.9 (-28.1 to 15.0)	-14.9 (-30.8 to -14.9)	-23.3 (-37.1 to -2.6)	-1.03
Mizoram	40.7 (34.7 to 47.0)	40.4 (37.1 to 44.8)	29.9 (26.1 to 33.9)	-0.8 (-16.8 to 17.7)	-25.8 (-37.5 to -14.1)	-26.4 (-39.8 to -10.9)	-0.08
Nagaland	43.8 (37.9 to 50.0)	41.4 (37.5 to 47.0)	30.4 (26.0 to 34.5)	-5.4 (-19.4 to 12.2)	-26.7 (-40.3 to -13.4)	-30.6 (-42.4 to -15.7)	-0.55
Uttarakhand	54.2 (48.7 to 60.8)	50.5 (44.6 to 57.4)	34.8 (30.4 to 39.4)	-6.8 (-22.5 to 9.5)	-31.1 (-42.8 to -17.0)	-35.8 (-46.2 to -24.1)	-0.71
Gujarat	57.1 (51.6 to 63.9)	54.4 (50.0 to 59.4)	41.3 (36.2 to 47.1)	-4.7 (-16.7 to 9.0)	-24.0 (-35.0 to -11.0)	-27.6 (-39.0 to -15.0)	-0.48
Tripura	51.0 (43.4 to 59.8)	47.8 (41.7 to 55.3)	31.0 (26.5 to 35.2)	-6.2 (-23.9 to 17.0)	-35.2 (-47.5 to -21.9)	-39.3 (-51.9 to -25.1)	-0.64
Sikkim	46.8 (40.2 to 54)	40.8 (36.3 to 47.0)	26.2 (22.2 to 30.0)	-12.8 (-27.4 to 6.4)	-35.8 (-47.9 to -21.6)	-44.0 (-54.8 to -31.0)	-1.36
Manipur	42.3 (34.4 to 50.6)	42.6 (38.4 to 49.3)	32.4 (27.9 to 36.4)	0.7 (-17.6 to 27.4)	-24.0 (-37.6 to -10.5)	-23.5 (-39.6 to -3.4)	0.07
Higher-middle ETI (446 million)	52.4 (49.6 to 55.5)	48.3 (46.1 to 50.7)	33.6 (31.9 to 35.5)	-7.8 (-14.2 to -1.3)	-30.5 (-35.2 to -25.4)	-35.9 (-40.7 to -30.8)	-0.81
Haryana	55.7 (50.0 to 62.8)	53.2 (49.4 to 57.9)	36.9 (32.4 to 41.1)	-4.5 (-17.0 to 10.8)	-30.7 (-41.0 to -20.9)	-33.8 (-44.9 to -22.8)	-0.46
Delhi	46.2 (38.3 to 54.7)	46.9 (42.1 to 53.5)	28.6 (23.4 to 33.9)	1.4 (-18.0 to 28.0)	-39.1 (-52.2 to -23.9)	-38.2 (-52.7 to -21.3)	0.14
Telangana	53.3 (47.6 to 59.9)	44.5 (37.8 to 52.4)	32.5 (28.2 to 37.0)	-16.6 (-30.9 to 2.0)	-26.9 (-40.7 to -9.4)	-39.0 (-49.0 to -26.5)	-1.80
Andhra Pradesh	52.7 (46.5 to 59.6)	46.1 (41.5 to 52.1)	33.9 (29.6 to 38.6)	-12.5 (-25.6 to 3.9)	-26.4 (-38.0 to -12.6)	-35.6 (-47.0 to -23.1)	-1.33
Jammu and Kashmir	50.2 (45.2 to 56.2)	46.2 (42.5 to 51.4)	30.4 (25.9 to 34.7)	-8.0 (-19.6 to 6.2)	-34.3 (-45.3 to -22.5)	-39.6 (-49.9 to -28)	-0.83
Karnataka	51.9 (46.1 to 58.9)	48.3 (44.2 to 54.0)	36.6 (32.1 to 41.0)	-6.9 (-19.9 to 8.6)	-24.4 (-35.6 to -12.1)	-29.6 (-40.8 to -17.6)	-0.71
West Bengal	52.7 (46.1 to 60.3)	50.9 (45.9 to 57.5)	33.5 (29.1 to 38.4)	-3.5 (-18.8 to 15.3)	-34.1 (-45.5 to -21.8)	-36.4 (-47.8 to -21.8)	-0.36
Maharashtra	52.3 (46.2 to 59.4)	48.2 (43.9 to 54.0)	32.6 (28.4 to 37.2)	-7.7 (-21.5 to 8.8)	-32.5 (-42.8 to -21.1)	-37.7 (-48.3 to -25.4)	-0.80
Union territories other than Delhi	45.9 (38.4 to 53.8)	38.1 (30.6 to 46.3)	23.1 (17.1 to 31.8)	-17.1 (-35.5 to 7.1)	-37.0 (-56.0 to -11.5)	-47.8 (-64.4 to -26.2)	-1.86
High ETI (152 million)	46.3 (42.5 to 49.9)	40.5 (38.4 to 43.2)	24.6 (22.5 to 26.7)	-12.5 (-20.1 to -2.5)	-39.3 (-45.8 to -32.9)	-46.9 (-52.4 to -40.6)	-1.33
Himachal Pradesh	54.9 (48.5 to 63.3)	51.3 (46.0 to 58.3)	34.1 (28.3 to 39.6)	-6.6 (-22.4 to 11.1)	-33.5 (-46.6 to -20.5)	-37.9 (-50.9 to -24.4)	-0.68
Punjab	51.1 (45.2 to 57.6)	47.7 (43.3 to 53.0)	32.1 (28.4 to 36.2)	-6.8 (-20.5 to 9.8)	-32.7 (-43.4 to -21.3)	-37.2 (-47.7 to -25.4)	-0.70
Tamil Nadu	45.3 (38.8 to 50.7)	38.8 (35.7 to 43.3)	23.4 (20.1 to 26.9)	-14.3 (-25.9 to 3.3)	-39.6 (-50.1 to -28.4)	-48.3 (-57.2 to -36.9)	-1.54
Goa	27.6 (21.0 to 36.0)	23.7 (19.8 to 28.5)	19.1 (15.6 to 22.6)	-14.1 (-37.5 to 16.3)	-31.1 (-49.5 to -5.7)	-1.51	-1.42
Kerala	44.3 (37.3 to 51.5)	37.3 (33.5 to 42.8)	19.3 (16.3 to 22.5)	-15.7 (-29.7 to 4.7)	-48.2 (-59.2 to -37.6)	-56.4 (-65.5 to -45.3)	-1.69

ETI =epidemiological transition level.

Table 6: Prevalence of exclusive breastfeeding in the first six months for India and states grouped by epidemiological transition level, 1990-2016

	Prevalence						Percent change			Annualised percent change		
	1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	1990-2000	2000-2016	1990-2016	1990-2000	2000-2016	1990-2016
States of India grouped by ETL												
India (1316 million)	37.7 (25.6 to 50.2)	38.9 (29.4 to 48.7)	39.9 (22.3 to 58.3)	3.7 (-8.3 to 16.3)	1.5 (-13.4 to 19.7)	5.3 (-11.2 to 26.6)	0.31	0.04	0.14			
Low ETL (626 million)	38.3 (31.2 to 46.1)	40.5 (31.9 to 48.8)	41.2 (33.7 to 49.5)	5.7 (-12.9 to 27.2)	1.8 (-19.5 to 29.3)	7.6 (-17.7 to 41.9)	0.56	0.11	0.28			
Bihar	39.1 (21.8 to 57.8)	43.1 (25.9 to 62.6)	43.4 (25.2 to 64.0)	10.2 (-30.6 to 76.3)	0.6 (-42.2 to 78.7)	10.8 (-43.6 to 116.4)	0.97	0.03	0.39			
Jharkhand	37.4 (20.9 to 55.3)	39.4 (21.5 to 58.1)	40.9 (22.5 to 59.1)	5.4 (-33.6 to 71)	3.7 (-43.5 to 88.0)	9.4 (-45.4 to 115.4)	0.53	0.23	0.34			
Uttar Pradesh	39.4 (22.6 to 58.2)	40.9 (24.0 to 60.5)	42.2 (23.4 to 59.4)	4.0 (-34.1 to 67.9)	3.1 (-41.8 to 86.6)	7.2 (-43.6 to 101.0)	0.40	0.19	0.27			
Rajasthan	38.5 (21.8 to 57.7)	38.9 (22.5 to 58.2)	38.8 (23.0 to 57.9)	1.1 (-36.5 to 62.7)	-0.1 (-42.2 to 79.0)	1.0 (-46.3 to 104.0)	0.11	-0.01	0.04			
Meghalaya	36.1 (19.1 to 54.6)	36.2 (20.5 to 55.0)	36.7 (19.3 to 55.8)	0.2 (-37 to 62.9)	1.4 (-46.4 to 88.5)	1.6 (-51.2 to 105.3)	0.02	0.09	0.06			
Assam	38.3 (22.5 to 57.1)	39.9 (23.5 to 58.5)	39.7 (21.9 to 58.7)	4.0 (-33.7 to 64.2)	-0.5 (-44.5 to 75.5)	3.4 (-48 to 101.7)	0.39	-0.03	0.13			
Chhattisgarh	37.7 (21.9 to 56.4)	40.9 (25.2 to 60.3)	41.1 (24.7 to 58.3)	8.5 (-29.1 to 70.9)	0.5 (-40.3 to 69.9)	9.1 (-41.5 to 104.8)	0.82	0.03	0.34			
Madhya Pradesh	35.7 (19.2 to 53.8)	38.1 (22.1 to 55.5)	39 (21.7 to 57.4)	6.6 (-33.4 to 84.8)	2.4 (-43.1 to 89.0)	9.2 (-43.6 to 126.7)	0.64	0.15	0.34			
Odisha	37.8 (21.7 to 56.7)	40.1 (23.0 to 58.5)	39.6 (22.5 to 58.6)	6.1 (-33.1 to 68.7)	-1.3 (-43.3 to 70.7)	4.7 (-45.2 to 106.8)	0.59	-0.08	0.18			
Lower-middle ETL (92 million)												
Arunachal Pradesh	37.0 (24.3 to 51.0)	37.8 (25.7 to 51.5)	37.1 (25.3 to 51.1)	2.1 (-25.9 to 44.1)	-1.9 (-37.6 to 54.6)	0.1 (-39.1 to 67.7)	0.21	-0.12	0.01			
Mizoram	36.1 (20.2 to 55.5)	37.9 (21.4 to 56.8)	37.9 (20.4 to 56.4)	5.1 (-36.5 to 69.3)	0 (-44.9 to 90.1)	5.1 (-47.5 to 107.5)	0.50	0.00	0.19			
Nagaland	37.4 (20.6 to 55.2)	38.4 (22.6 to 56.7)	37.3 (20.4 to 56.1)	2.9 (-34 to 66.6)	-3.0 (-46.9 to 74.2)	-0.2 (-49.4 to 98.1)	0.28	-0.19	-0.01			
Uttarakhand	40.1 (23.1 to 58.9)	40.1 (23.2 to 58.0)	38.6 (22.0 to 56.7)	0.1 (-38.5 to 61.7)	-3.7 (-48.0 to 72.3)	-3.6 (-51.9 to 74.8)	0.01	-0.24	-0.14			
Gujarat	37.0 (21.2 to 55.6)	39.4 (22.7 to 57.5)	38.3 (21.6 to 57.0)	6.5 (-32.8 to 67.8)	-2.9 (-47.1 to 81.4)	3.4 (-49.5 to 105.4)	0.63	-0.18	0.13			
Tripura	36.4 (19.8 to 55.3)	36.7 (20.5 to 55.5)	36.5 (21.3 to 55.5)	0.9 (-36.6 to 66.0)	-0.7 (-46.6 to 85.7)	0.2 (-50.5 to 102.4)	0.09	-0.05	0.01			
Sikkim	39.1 (21.7 to 57.9)	39.2 (21.7 to 59.5)	37.5 (22.2 to 55.6)	0.3 (-38.0 to 65.8)	-4.2 (-46.3 to 74.4)	-3.9 (-47.5 to 88.7)	0.03	-0.27	-0.15			
Manipur	35.3 (19.7 to 54.2)	36.8 (20.1 to 55.9)	32.2 (17.9 to 49.6)	4.4 (-35.6 to 64.5)	-12.5 (-52.1 to 59.9)	-8.6 (-53.2 to 70.8)	0.43	-0.83	-0.35			
Higher-middle ETL (446 million)												
Haryana	44.7 (28.0 to 64.2)	46.3 (29.2 to 65.3)	46.4 (28.9 to 65.8)	3.6 (-30.6 to 56.8)	0.3 (-39.8 to 62.9)	3.9 (-41.4 to 81.2)	0.35	0.02	0.15			
Delhi	37.1 (29.6 to 45.6)	37.9 (31.0 to 45.1)	38.1 (31.2 to 45.9)	2.1 (-15.7 to 23.2)	0.6 (-20.6 to 26.7)	2.8 (-21.6 to 36.4)	0.21	0.03	0.10			
Telangana	37.4 (20.1 to 54.6)	37.7 (20.4 to 56.6)	37.0 (21.5 to 55.0)	0.8 (-37.9 to 66.7)	-1.8 (-45.3 to 85.3)	-1.1 (-47.7 to 100.4)	0.08	-0.12	-0.04			
Andhra Pradesh	38.4 (22.4 to 56.9)	38.0 (21.6 to 56.4)	36.3 (20.8 to 55.2)	-1.1 (-38.3 to 59.6)	-4.5 (-45.7 to 66.0)	-5.5 (-51.4 to 78.3)	-0.11	-0.28	-0.22			
Jammu and Kashmir	30.9 (16.4 to 48.6)	33.1 (17.1 to 50.1)	36.8 (20.6 to 55.3)	7.1 (-34.0 to 75.6)	11.2 (-42.1 to 114.3)	19.1 (-42.4 to 150.9)	0.69	0.66	0.67			
Karnataka	40.7 (24.0 to 59.1)	41.6 (24.4 to 60.4)	40.1 (22.8 to 58.5)	2.4 (-35.7 to 62.4)	-3.8 (-43.2 to 65.9)	-1.7 (-48.4 to 86.2)	0.24	-0.24	-0.06			
West Bengal	40.6 (22.7 to 58.7)	41.5 (24.0 to 60.0)	41.5 (24.7 to 59.9)	2.4 (-36.0 to 63.1)	-0.1 (-42.2 to 71.5)	2.2 (-45.2 to 97.6)	0.23	-0.01	0.08			
Madhya Pradesh	37.9 (21.2 to 57.3)	38.4 (21.5 to 57.6)	39.0 (21.1 to 57.1)	1.5 (-37.7 to 62.0)	1.5 (-42.7 to 81.4)	3.0 (-45.4 to 93.2)	0.15	0.09	0.11			
Punjab	37.3 (20.3 to 55.1)	37.5 (21.7 to 55.2)	37.8 (19.8 to 57.4)	0.7 (-36.5 to 64.4)	0.7 (-46.3 to 80.4)	1.4 (-51.1 to 118.7)	0.07	0.04	0.05			
Union territories other than Delhi	36.4 (19.4 to 54.5)	37.4 (21.2 to 56.1)	37.7 (21.9 to 54.8)	2.8 (-35.3 to 66.2)	0.9 (-42.1 to 81.5)	3.8 (-46.7 to 109.7)	0.28	0.06	0.14			
High ETL (152 million)												
37.9 (20.7 to 56.0)	38.4 (20.7 to 56.6)	37.0 (19.2 to 55.8)	1.3 (-37.0 to 68.5)	-3.5 (-48.6 to 78.9)	-2.3 (-54.1 to 95.2)	0.13	-0.22	-0.09				
38.6 (28.3 to 50.0)	38.5 (28.6 to 49.1)	38.6 (28.4 to 48.8)	-0.3 (-23.7 to 30.2)	0.3 (-29.4 to 42.3)	0 (-33.2 to 42.7)	0.02	0.02	0.00				
Himachal Pradesh	39.3 (21.9 to 57.3)	39.6 (22.6 to 58.1)	39.2 (21.6 to 58.6)	0.6 (-37.7 to 58.3)	-1.1 (-44.2 to 81.4)	-0.5 (-48.7 to 90.5)	0.06	-0.07	-0.02			
Tamil Nadu	36.1 (20.0 to 56.2)	37.2 (21.3 to 56.2)	37.9 (21.3 to 55.7)	2.9 (-38.1 to 66.8)	2.1 (-43.8 to 81.4)	5.1 (-48.2 to 116.1)	0.29	0.13	0.19			
Goa	37.0 (21.4 to 55.5)	36.8 (20.5 to 55.1)	33.8 (18.4 to 51.5)	-0.5 (-40.3 to 62.8)	-8.0 (-50.1 to 68.5)	-8.5 (-53.5 to 83.6)	-0.05	-0.52	-0.34			
Kerala	40.5 (23.3 to 60.3)	39.7 (23.0 to 58.0)	39.7 (23.2 to 57.2)	-2.1 (-37.5 to 60.3)	0.0 (-44.3 to 83.7)	-2.1 (-49.1 to 89.9)	-0.21	0.00	-0.08			

ETL=epidemiological transition level.

Table 7: Prevalence of anaemia in reproductive age women for India and states grouped by epidemiological transition level, 1990-2016

	States of India grouped by ETI	Prevalence			Percent change			Annualised percent change		
		1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	1990-2000	2000-2016	1990-2016
India (1316 million)	55.9 (54.7 to 57.3)	54.3 (53.1 to 55.5)	56.2 (54.9 to 57.5)	-3.0 (-4.1 to -1.9)	3.5 (2.3 to 4.8)	0.4 (-0.5 to 1.3)	-0.29	0.22	0.02	
Low ETI (626 million)	59.4 (58 to 60.6)	57.5 (56.2 to 58.8)	59.3 (57.9 to 60.7)	-3.2 (-4.8 to -1.5)	3.2 (1.4 to 4.9)	-0.1 (-1.4 to 1.3)	-0.32	0.19	-0.01	
Bihar	66.0 (64.0 to 68.0)	63.8 (62.0 to 65.6)	66.0 (64.3 to 67.9)	-3.4 (-6.4 to -0.2)	3.5 (0.2 to 6.8)	0.0 (-2.5 to 3.0)	-0.33	0.21	0.00	
Jharkhand	64.2 (62.3 to 66.1)	62.7 (60.8 to 64.6)	64.8 (62.8 to 66.6)	-2.2 (-4.9 to 0.9)	3.3 (0.2 to 6.9)	1.0 (-1.5 to 3.8)	-0.23	0.21	0.04	
Uttar Pradesh	55.9 (54.0 to 58.0)	53.7 (52.2 to 55.4)	55.8 (54.1 to 57.6)	-3.9 (-7.2 to -0.8)	3.9 (0.7 to 7.2)	-0.2 (-3.0 to 2.8)	-0.40	0.24	-0.01	
Rajasthan	56.6 (55.1 to 58.1)	54.7 (53.3 to 56.3)	56.6 (55.0 to 58.0)	-3.3 (-5.6 to -0.7)	3.4 (0.7 to 5.9)	0.0 (-2.3 to 2.4)	-0.34	0.21	0.00	
Meghalaya	52.7 (51.0 to 54.4)	52.1 (50.6 to 53.8)	52.6 (51.2 to 54.1)	-1.2 (-3.8 to 2.0)	1.0 (-1.9 to 3.7)	-0.2 (-2.8 to 2.7)	-0.12	0.06	-0.01	
Assam	67.9 (66.0 to 69.5)	66.4 (64.5 to 68.2)	67.0 (65.2 to 58.7)	-2.2 (-4.5 to 0.4)	1.0 (-1.5 to 3.8)	-1.2 (-3.7 to 1.5)	-0.22	0.06	-0.05	
Chhattisgarh	55.5 (53.4 to 57.8)	55.0 (53.1 to 57.1)	56.7 (54.7 to 58.6)	-0.8 (-4.5 to 3.1)	3.1 (-0.9 to 7.4)	2.3 (-1.6 to 6.3)	-0.09	0.19	0.08	
Madhya Pradesh	58.2 (56.5 to 60.1)	56.1 (54.3 to 57.9)	58.3 (56.6 to 60.0)	-3.7 (-6.4 to -0.9)	4.0 (1.0 to 7.3)	0.2 (-2.1 to 2.5)	-0.37	0.24	0.01	
Odisha	58.6 (56.6 to 60.3)	57.9 (56.5 to 59.3)	58.9 (57.2 to 60.5)	-1.2 (-3.8 to 1.8)	1.8 (-1.6 to 4.5)	0.6 (-2.7 to 3.5)	-0.12	0.11	0.02	
Lower-middle ETI (92 million)		53.3 (51.9 to 54.6)	51.5 (50.1 to 52.8)	52.8 (51.5 to 54.2)	-3.4 (-5.5 to -1.4)	2.6 (0.5 to 4.8)	-0.8 (-2.7 to 1.1)	-0.34	0.16	-0.04
Arunachal Pradesh	50.2 (48.1 to 52.2)	49.5 (47.8 to 51.3)	50.6 (48.7 to 52.4)	-1.4 (-4.8 to 2.5)	2.2 (-1.2 to 5.7)	0.7 (-2.8 to 4.2)	-0.14	0.14	0.03	
Mizoram	41.9 (40.4 to 43.6)	40.6 (39.2 to 42.2)	41.4 (39.8 to 43.2)	-3.1 (-6.7 to 0.1)	2.1 (-1.8 to 5.9)	-1.1 (-4.9 to 2.8)	-0.31	0.12	-0.04	
Nagaland	38.9 (37.4 to 40.7)	38.0 (36.6 to 39.4)	37.5 (36.1 to 38.9)	-2.4 (-6.0 to 1.6)	1.3 (-5.1 to 2.4)	-3.7 (-7.2 to 0.3)	-0.24	-0.08	-0.14	
Uttarakhand	49.9 (48.4 to 51.7)	48.7 (47.0 to 50.4)	50.3 (48.6 to 52.0)	-2.5 (-5.5 to 0.6)	3.3 (0.2 to 7.1)	0.7 (-2.6 to 3.7)	-0.25	0.20	0.03	
Gujarat	55.1 (53.6 to 56.9)	53.3 (52.1 to 54.6)	54.7 (53.3 to 56.2)	-3.2 (-6.1 to -0.8)	2.6 (0.0 to 5.2)	-0.7 (-3.0 to 1.6)	-0.33	0.16	-0.03	
Tripura	57.0 (55.2 to 58.8)	55.7 (54.0 to 57.2)	57.8 (56.1 to 59.4)	-2.3 (-5.3 to 0.4)	3.8 (0.7 to 6.7)	1.4 (-1.9 to 4.2)	-0.23	0.23	0.05	
Sikkim	52.0 (50.3 to 53.9)	51.5 (49.6 to 53.3)	52.2 (50.2 to 54.2)	-1.0 (-4.7 to 2.6)	1.4 (-2.0 to 4.8)	0.4 (-3.4 to 4.4)	-0.11	0.08	0.01	
Manipur	33.1 (31.4 to 34.8)	32.1 (30.3 to 33.8)	33.9 (32.2 to 35.5)	-3.1 (-7.8 to 2.0)	5.5 (0.6 to 10.7)	2.2 (-2.6 to 7.0)	-0.31	0.34	0.09	
Higher-middle ETI (446 million)		54.5 (53.2 to 55.8)	53.1 (51.9 to 54.2)	54.8 (53.5 to 56.1)	-2.7 (-4.1 to -1.5)	3.3 (1.8 to 5.2)	0.6 (-9.7 to 1.8)	-0.26	0.20	0.02
Haryana	57.5 (56.0 to 58.9)	56.4 (54.6 to 58.1)	58.0 (56.3 to 59.4)	-1.9 (-4.3 to 0.7)	2.8 (0.0 to 5.6)	0.9 (-1.7 to 3.7)	-0.19	0.17	0.04	
Delhi	48.8 (46.8 to 50.5)	46.3 (44.8 to 48.2)	48.4 (46.5 to 50.6)	-5.2 (-8.8 to -1.1)	4.7 (0.3 to 9.7)	-0.8 (-4.8 to 3.1)	-0.53	0.28	-0.03	
Telangana	55.1 (53.5 to 56.6)	54.3 (52.8 to 55.8)	56.0 (54.3 to 57.6)	-1.5 (-4.0 to 1.1)	3.1 (0.3 to 5.8)	1.6 (-1.0 to 4.2)	-0.14	0.19	0.06	
Andhra Pradesh	58.2 (56.6 to 59.9)	57.1 (55.6 to 58.7)	58.5 (57.0 to 60.0)	-1.8 (-4.1 to 0.3)	2.5 (0.0 to 4.9)	0.6 (-1.6 to 2.9)	-0.19	0.15	0.02	
Jammu and Kashmir	47.2 (45.5 to 48.7)	46.7 (45.4 to 47.9)	47.7 (46.1 to 49.3)	-1.2 (-4.4 to 2.1)	2.2 (-1.2 to 5.4)	1.0 (-2.8 to 4.4)	-0.12	0.14	0.04	
Karnataka	51.9 (50.4 to 53.5)	49.7 (48.4 to 51.0)	52.3 (50.7 to 53.7)	-4.3 (-7.1 to -1.6)	5.2 (2.2 to 8.4)	0.7 (-2.1 to 3.2)	-0.44	0.32	0.03	
West Bengal	58.1 (56.4 to 59.9)	57.6 (56.0 to 59.4)	58.5 (56.8 to 60.2)	-0.8 (-3.5 to 1.8)	1.5 (-1.4 to 4.4)	0.6 (-2.1 to 3.4)	-0.09	0.10	0.03	
Maharashtra	51.6 (50.0 to 53.2)	49.6 (48.3 to 51.2)	52.1 (50.6 to 53.7)	-3.9 (-6.7 to -1.4)	5.0 (2.0 to 8.3)	0.9 (-1.7 to 3.3)	-0.39	0.31	0.04	
Union territories other than Delhi	55.1 (53.5 to 56.6)	53.4 (51.7 to 55.1)	54.6 (52.7 to 56.6)	-3.0 (-5.9 to -0.4)	2.4 (-0.7 to 5.5)	0.7 (-3.3 to 2.1)	-0.30	0.14	-0.03	
High ETI (152 million)		50.7 (49.4 to 52.0)	48.9 (47.6 to 50.2)	50.6 (49.3 to 51.9)	-3.5 (-5.2 to -1.7)	3.4 (1.5 to 5.3)	-0.2 (-2.1 to 1.5)	-0.36	0.21	-0.01
Himachal Pradesh	43.5 (41.6 to 45.3)	42.1 (40.6 to 43.5)	44.7 (42.7 to 46.4)	-3.1 (-7.1 to 1.7)	6.1 (1.7 to 11.0)	2.7 (-1.1 to 7.5)	-0.32	0.38	0.11	
Punjab	52.8 (51.1 to 54.7)	51.7 (50.2 to 53.5)	53.2 (51.7 to 54.7)	-2.0 (-5.0 to 1.3)	2.8 (-0.2 to 5.6)	0.7 (-1.8 to 3.4)	-0.21	0.18	0.03	
Tamil Nadu	57.8 (56.3 to 59.4)	55.3 (54.0 to 56.7)	56.8 (55.2 to 58.3)	-4.4 (-6.5 to -1.9)	2.7 (0.2 to 4.9)	-1.9 (-4.4 to 0.8)	-0.45	0.17	-0.07	
Goa	40.4 (38.6 to 42.0)	39.9 (38.3 to 41.4)	41.0 (39.2 to 42.8)	-1.3 (-5.0 to 2.3)	2.7 (-1.1 to 6.6)	1.4 (-2.6 to 5.3)	-0.13	0.17	0.06	
Kerala	37.6 (35.8 to 39.7)	36.4 (34.7 to 38.3)	37.2 (35.4 to 38.9)	-3.2 (-7.7 to 1.8)	2.2 (-3.2 to 8.2)	-1.1 (-5.6 to 3.2)	-0.32	0.14	-0.04	

ETI=epidemiological transition level.

Table 8: Prevalence of overweight in children aged 2-4 years for India and states grouped by epidemiological transition level, 1990-2016

	Prevalence			Percent change			Annualised percent change		
	1990 (95% uncertainty interval)	2000 (95% uncertainty interval)	2016 (95% uncertainty interval)	1990-2000 (95% uncertainty interval)	2000-2016 (95% uncertainty interval)	1990-2016 (95% uncertainty interval)	1990-2000	2000-2016	1990-2016
States of India grouped by ETL									
India (1316 million)	7.6 (4.4 to 11.9)	7.3 (4.2 to 11.5)	11.8 (7.1 to 17.9)	-3.4 (-21.6 to 19.6)	60.7 (27.4 to 100.3)	55.1 (22.2 to 96.1)	-0.35	3.01	1.70
Low ETL (626 million)	6.4 (4.8 to 8.6)	5.4 (4.1 to 7.3)	7.8 (5.8 to 10.5)	-15.3 (-38.6 to 20.8)	44.6 (-2.2 to 117.8)	22.4 (-18.6 to 83.2)	-1.65	2.30	0.76
Bihar	6.8 (3.3 to 12.9)	4.3 (2.1 to 8.1)	5.8 (2.8 to 10.0)	-36.9 (-70.0 to 33.6)	34.9 (-43.0 to 235.2)	-14.8 (-65.9 to 114.1)	-4.50	1.89	-0.62
Jharkhand	4.6 (2.2 to 8.2)	4.1 (2.0 to 7.4)	6.0 (3.0 to 10.9)	-12.4 (-57.3 to 93.5)	47.5 (-40.0 to 251.9)	29.3 (-51.2 to 210.7)	-1.31	2.46	0.99
Uttar Pradesh	7.7 (3.9 to 13.7)	6.6 (3.6 to 11.3)	9.3 (4.6 to 16.0)	-13.5 (-58.3 to 88.8)	40.1 (-39.2 to 210.1)	21.1 (-51.6 to 184.0)	-1.44	2.13	0.74
Rajasthan	5.6 (2.9 to 9.9)	5.6 (2.9 to 10.1)	8.3 (4.1 to 14.3)	-1.3 (-55.2 to 108.1)	49.7 (-39.4 to 245.8)	47.7 (-39.5 to 236.0)	-0.13	2.55	1.51
Meghalaya	12.8 (6.6 to 22.5)	1.3 (7.5 to 21.1)	19.4 (13.2 to 27.6)	1.3 (-47.7 to 101.8)	49.3 (-20.6 to 196.0)	51.3 (-21.8 to 230.9)	0.13	2.54	1.61
Assam	7.3 (3.6 to 13.2)	6.6 (3.4 to 11.4)	9.5 (4.7 to 17.2)	-9.6 (-55.1 to 86.8)	45.6 (-39.9 to 257.0)	31.6 (-46.3 to 232.6)	-1.01	2.38	1.06
Chhattisgarh	6.7 (3.1 to 13.5)	5.2 (2.6 to 9.6)	9.0 (4.4 to 16.3)	-22.7 (-64.0 to 85.9)	73.1 (-32.7 to 301.5)	33.9 (-50.3 to 252.1)	-2.54	3.49	1.13
Madhya Pradesh	4.3 (2.1 to 7.8)	3.8 (1.8 to 7.1)	5.4 (2.6 to 9.8)	-11.4 (-58.1 to 83.4)	42.1 (-40.2 to 246.9)	25.9 (-48.8 to 203.9)	-1.21	2.22	0.89
Odisha	5.3 (2.5 to 9.6)	4.8 (2.7 to 8.1)	8.4 (4.4 to 14.7)	-8.1 (-55.4 to 96.6)	73.3 (-28.6 to 283.6)	59.2 (-32 to 287.6)	-0.84	3.49	1.81
Lower-middle ETL (92 million)									
Arunachal Pradesh	6.6 (4.3 to 10.2)	6.8 (4.7 to 9.9)	11.3 (7.7 to 16.9)	2.5 (-37.5 to 65.4)	65.7 (-0.2 to 180.7)	69.8 (-3.9 to 198.6)	0.24	3.23	2.07
Mizoram	12.5 (6.2 to 22.4)	13.3 (7.7 to 22.0)	26.3 (18.9 to 35)	6.3 (-44.4 to 110.9)	97.7 (4.2 to 261.3)	110.2 (10.0 to 355.9)	0.62	4.35	2.90
Nagaland	13.4 (7.1 to 23)	14.0 (8.0 to 22.2)	16.5 (11.0 to 24.1)	4.6 (-47.1 to 104.5)	17.6 (-38.6 to 133.7)	23.1 (-38.4 to 156.4)	0.45	1.02	0.80
Uttarakhand	14.7 (7.4 to 24.9)	15.3 (9.0 to 23.8)	18.2 (12.0 to 25.7)	3.8 (-45.6 to 102.1)	19.5 (12.5 to 118.9)	24.0 (-36.0 to 169.6)	0.37	1.12	0.83
Gujarat	6.4 (3.1 to 12.1)	6.2 (3.1 to 11.3)	13.6 (6.9 to 25.2)	-3.4 (-58.6 to 113.5)	120.2 (-6.4 to 442.9)	112.8 (-12.5 to 440.9)	-0.34	5.06	2.95
Tripura	6.0 (3.2 to 10.9)	6.0 (3.2 to 10.2)	9.4 (4.9 to 16.7)	-0.5 (-52.3 to 115.1)	57.4 (-33.4 to 263.2)	56.6 (-36.2 to 286.1)	-0.05	2.88	1.74
Sikkim	7.8 (3.9 to 14.0)	7.7 (4.1 to 13.1)	16.2 (10.9 to 23.8)	-1.0 (-50.6 to 107.4)	110.4 (5.5 to 339.5)	108.3 (4.8 to 359.1)	-0.10	4.76	2.86
Manipur	6.6 (3.3 to 12.0)	6.4 (3.6 to 10.7)	18.0 (11.2 to 27.6)	20.9 (-31.4 to 130.2)	29.0 (-61.5 to 39.0)	-14.1 (-57.3 to 82.9)	1.92	-2.12	-0.58
Higher-middle ETL (446 million)									
Haryana	9.1 (6.8 to 12.1)	9.6 (7.4 to 12.4)	18.1 (14.7 to 22.2)	5.3 (-24.5 to 48.4)	88.7 (37.9 to 158.4)	98.6 (40.0 to 182.3)	0.51	4.03	2.66
Delhi	4.2 (2.0 to 7.6)	4.9 (2.5 to 8.9)	15.8 (9.4 to 23.9)	18.8 (-44.2 to 148.4)	219.3 (45.6 to 584.5)	279.4 (70.8 to 767.2)	1.74	7.53	5.26
Telangana	12.4 (6.1 to 22.5)	12.6 (6.4 to 21.3)	26.2 (16.9 to 37.8)	1.4 (-51.8 to 101.7)	108.4 (7.3 to 351.6)	111.2 (24.4 to 356.8)	0.14	4.70	2.92
Andhra Pradesh	16.2 (8.3 to 28.4)	17.0 (8.7 to 29.8)	28.4 (19.0 to 39.1)	5.1 (-49.5 to 112.5)	67.0 (-15.9 to 254.0)	75.6 (-12.3 to 274.5)	0.50	3.26	2.19
Jammu and Kashmir	7.7 (3.8 to 13.2)	8.8 (4.8 to 15.4)	18.9 (10.9 to 29.4)	14.3 (-45.2 to 139.2)	116.3 (-5.1 to 360.3)	147.3 (14.4 to 470.9)	1.35	4.94	3.54
Karnataka	8.0 (3.8 to 14.6)	7.6 (4.0 to 12.7)	12.7 (6.3 to 22.2)	-5.5 (-55.2 to 102.9)	67.8 (-27.2 to 284.4)	58.5 (-34 to 269.6)	-0.57	3.29	1.79
West Bengal	5.9 (2.7 to 9.8)	5.9 (2.9 to 10.8)	17.3 (10.5 to 26.1)	3.7 (-47.5 to 105.6)	62.9 (-28.3 to 270.5)	68.9 (-30.8 to 313.5)	0.36	3.10	2.04
Maharashtra	12.2 (6.3 to 20.9)	12.5 (6.9 to 20.9)	18.5 (11.0 to 28.8)	2.5 (-50.1 to 116.3)	47.6 (-31.2 to 214.0)	51.2 (-29.3 to 242.0)	0.25	2.46	1.60
Union territories other than Delhi	16.9 (9.3 to 29.4)	14.2 (7.3 to 25.1)	25.4 (12.9 to 41.2)	-16.0 (-59.2 to 73.3)	78.7 (-21.2 to 290.1)	50.1 (-37.2 to 239.1)	-1.73	1.90	0.49
High ETL (152 million)									
Himachal Pradesh	3.9 (1.8 to 6.6)	5.1 (2.5 to 9.1)	21.6 (14.6 to 31.1)	31.1 (-38.6 to 182.2)	326.4 (115.2 to 846.7)	459.1 (181.5 to 1071.8)	2.75	9.49	6.84
Punjab	7.1 (3.6 to 13.1)	7.5 (3.7 to 13.5)	13.1 (7.6 to 19.9)	5.3 (-48.4 to 123.9)	75.2 (-15.2 to 289.9)	84.5 (-13.3 to 314.6)	0.52	3.57	2.38
Tamil Nadu	9.0 (4.6 to 15.9)	10.1 (5.0 to 18.3)	15.7 (8.4 to 26.6)	12.1 (-47.0 to 142.0)	55.7 (-31.1 to 279.8)	74.5 (-22.1 to 320.1)	1.15	2.81	2.17
Goa	12.2 (6.2 to 20.9)	13.9 (7.5 to 22.9)	25.6 (16.8 to 37.9)	14.2 (-39.3 to 128.7)	84.8 (-6.9 to 281.6)	110.9 (4.1 to 343.2)	1.33	3.91	2.91
Kerala	7.1 (3.5 to 12.7)	8.3 (4.2 to 15.1)	17.3 (-43.6 to 148.8)	180.3 (34.2 to 515.1)	228.8 (58.8 to 640.9)	1.61	6.65	4.68	

ETL=epidemiological transition level.