

BMJ Open

How well do mothers recall their own and their infants' perinatal events? a two-district study using cross-sectional stratified random sampling in Bihar, India

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-031289.R3
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Valadez, Joseph; Liverpool School of Tropical Medicine, International Public Health Devkota, Baburam; Liverpool School of Tropical Medicine, International Public Health Jeffery, Caroline; Liverpool School of Tropical Medicine Hadden, Wilbur; University of Maryland at College Park, Sociology
Primary Subject Heading:	Global health
Secondary Subject Heading:	Epidemiology, Global health, Public health, Research methods
Keywords:	Public health < INFECTIOUS DISEASES, maternal recall, Bihar, India, perinatal, surveys, LQAS

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 How well do mothers recall their own and their infants' perinatal events? a two-district
4 study using cross-sectional stratified random sampling in Bihar, India

5
6
7
8 Joseph J Valadez^{1*}, Baburam Devkota¹, Caroline Jeffery¹, Wilbur C. Hadden²

9
10 ¹ Department of International Public Health, Liverpool School of Tropical Medicine,
11 Liverpool, United Kingdom

12 ² Department of Sociology, Center for Innovation, University of Maryland, USA

13 * Corresponding author: Professor Joseph J Valadez, Department of International Public
14 Health, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA
15 United Kingdom. Email: joseph.valadez@lstmed.ac.uk; Mobile: +44(0)754 8523717

16
17 Manuscript word length: 2255

Abstract

Objective: Global monitoring of maternal, newborn and child health (MNCH) programmes use self-reported data subject to recall error which may lead to incorrect decisions for improving health services and wasted resources. To minimize this risk, samples of mothers of infants aged 0-2 and 3-5 months are sometimes used. We test whether a single sample of mothers of infants 0-5 months provides the same information.

Design: An annual MNCH household survey in 2-districts of Bihar, India (N=6million).

Participants: Independent samples (n=475 each) of mothers of infants 0-5, 0-2 and 3-5 months.

Outcome measures: Main analyses compare responses from the 0-5 and 0-2 month samples with Mantel-Haenszel-Cochran statistics using 51 indicators in 2-districts.

Results: No measurable differences detected in 79.4% (81/102) comparisons; 20.6% (21/102) display differences for the main comparison. Sub-analyses produce similar results. A difference detected for exclusive breastfeeding is due to premature complementary-feeding by older infants. Measurable differences were detected in 33% (8/24) of the indicators on Front Line Worker (FLW) support, 26.9% (7/26) of indicators of birth preparedness and place of birth and

1 attendant, and 9.5% (4/42) of the indicators on neonatal
2 and antenatal care.

3 **Conclusions:** Differences in FLW visits and compliance with
4 their advice may be due to seasonal effects: mothers of
5 older infants 3-5 months were pregnant during the dry
6 season; mothers of infants 0-2 months were pregnant during
7 the monsoons, making transportation difficult. Useful
8 coverage estimates can be obtained by sampling mothers with
9 infants 0-5 months as with two samples suggesting that
10 mothers of young infants recall their own perinatal events
11 and those of their children. For some indicators (e.g.,
12 exclusive breast feeding) it may be necessary to adjust
13 targets. Excessive stratification wastes resources, does
14 not improve the quality of information, and increases the
15 burden placed on data collectors and communities, which can
16 increase non-sampling error.

17 18 **STRENGTHS AND LIMITATIONS OF THIS STUDY**

- 19 • Strength: The data were produced using stratified random sampling with no apparent
20 design effect leading to an efficient use of information.
- 21 • Strength: Data were collected from female participants by female data collectors
22 which is likely to have reduced non-sampling error.

- 1 • Strength: The large study population covers a large geographical area, reducing the
2 likelihood that the results are pertinent only to a small group of mothers with infants,
3 and may be generalizable.
- 4 • Strength: Both weighted and unweighted results are presented giving strength to the
5 conclusions.
- 6 • Limitation: Due to insufficient overlap of variables in the 0-5 month sample and
7 the 3-5 month sample, comparison between the 3-5 month sample vis a vis the
8 0-5 month sample was not possible.

9 **Key Words:**

10 Maternal recall, surveys, maternal health, MNCH, LQAS,
11 Bihar, India

12
13 **INTRODUCTION**

14 The progress toward United Nations' Sustainable Development
15 Goal (SDG) 3 is measured with 9 targets, including the
16 Maternal Mortality Ratio (MMR) and the under 5 mortality
17 rate (U5MR) (1, 2). In India, Bihar is one of the largest
18 (population 110 million) and poorest (53% of households are
19 in the lowest wealth index quintile of India (3)) states
20 with high child and maternal mortality (U5MR=54,
21 MMR=208) (4), and is a priority for donor support for health
22 systems strengthening (see (5) for an evaluation of the
23 health care system in Bihar).

1 To accelerate progress toward achieving SDG 3, state
2 governments in India pursue programmes of community-based
3 care (see (6, 7) for descriptions and assessments of this
4 approach). Since 2011, the Bihar Ministry of Health has
5 supported an Integrated Family Health Initiative to improve
6 the availability, quality and use of prenatal, perinatal
7 and postnatal care for mothers and infants (8).

8 The usual way to monitor progress toward achieving these
9 goals is with household surveys. Perhaps the most commonly
10 used surveys are cluster sample surveys such as the
11 Demographic and Health Surveys and the Multiple Indicator
12 Cluster Surveys (9, 10). An alternative design is Lot
13 Quality Assurance Sampling (LQAS), which provides
14 comparable data but is decentralized to local health
15 services organisations and more useful for management and
16 programme planning(11). Several states in India find it
17 benefits their programs (12). Surveys rely on the reports
18 of mothers of infants and young children, but these reports
19 are subject to several sources of potential error and bias
20 through interviewees not knowing, forgetting, and having
21 memory errors (13, 14). Studies have shown both that
22 mothers can accurately report significant facts about the
23 birth and care of their children many years after the event
24 (15), but also that even immediately after giving birth

1 mothers may misreport details (16-18). Studies of mothers
2 recall of their children's vaccination status concluded
3 that due to offsetting errors of maternal reports, the
4 resulting data accurately measured vaccination rates (19);
5 the pattern of error revealed that mothers whose children
6 are up-to-date or nearly so tended to underestimate their
7 child's vaccination status whilst mothers whose children
8 have few vaccinations, overestimate their coverage.
9 To improve the validity of collected data, knowledge,
10 practice and coverage surveys have used samples of mothers
11 of infants 0-11 months of age or 0-5 months of age and
12 children 6-11 months of age. In Bihar, local organisations
13 departed from this convention of sampling among these three
14 cohorts of children under one year of age and have been
15 monitoring their programs' progress by sampling five
16 dedicated cohorts: mothers of children 0-2, 3-5, 6-8, 9-11
17 and 12-23 months old with indicators focused on antenatal
18 care, safe delivery practices, infant and young child
19 feeding practices, immunisation, treatment seeking, and
20 more. To avoid the possibility of maternal recall error,
21 each of the five cohorts was asked questions particularly
22 relevant to a child's specific age group.

23 In countries such as India with high maternal and
24 child mortality rates, regular monitoring of related health

1 service coverage is critical to reducing these rates.
2 However, survey designs should be affordable and
3 sustainable for local health systems; they should also
4 produce precise, unbiased estimates (20). In this study, we
5 explore whether information is gained by sampling cohorts
6 of children aged 0-2 and 3-5 months or whether sample sizes
7 can be reduced by 50% by creating one sample cohort aged 0-
8 5 months.

9 The research question we address is: "Do the health service
10 delivery coverage estimates from a sample of mothers of
11 infants aged 0-5 months differ from those obtained from a
12 sample of mothers of infants ages 0-2 months?" A corollary
13 to this question is: "Do mothers of infants 3-5 months of
14 age display more recall error relative to mothers of
15 infants 0-2 months of age for antenatal, delivery or young
16 infant health practices?" We compare district coverage
17 estimates obtained from two independent samples of infants
18 0-2 months and 0-5 months of age. The implications of this
19 study are important for health systems researchers needing
20 results to appraise and improve their programmes.

21 **METHODS**

22 To answer this question, we collected information from a
23 sample of mothers with infants 0-5 months old and a sample

1 of mothers with infants 0-2 month old in two districts.
2 This study took place within the context of a larger survey
3 that also sampled children 3-5, 6-8, 9-11 and 12-23 months
4 of age. These four latter samples used questionnaires with
5 variables that either did not overlap at all or overlapped
6 on very few indicators with the questionnaires used to
7 interview the 0-5 month and 0-2 month samples. Due to this
8 constraint, in this study, we only use the two
9 aforementioned groups to assess the measurement of the
10 indicators and refer only to them for the remainder of this
11 paper. The household sampling design we used is a
12 stratified random sample (21). Within each district, the
13 strata are administrative units of the health system, which
14 in Bihar is called a *block*. Within each block the primary
15 sampling unit is the Anganwadi Centre (Community Health
16 Subcentre) Catchment Area (ACCA); 19 ACCA are selected from each block
17 with probability proportional to size. From each ACCA one respondent is
18 randomly selected from each age-group under study using
19 segmentation sampling (22, 23). The sample size 19 for each block is
20 chosen to maximize the probability of correctly classifying a block with reference to
21 performance targets on health related indicators (95% reliability) while balancing the
22 probability (10% margin of error) of incorrectly classifying a block and thereby
23 failing to recognize either the accomplishments of local health care delivery systems or

1 the local population's health care needs (22). For this purpose, principles of Lot Quality
2 Assurance Sampling were used along with established probability tables (24-26).
3 There are 14 and 11 blocks in Gopalganj and Aurangabad (N=6
4 million), the two districts selected for this study,
5 respectively. The total sample sizes are: (a) Gopalganj: 19
6 x 14 blocks= 266 infants 0-2 months and 266 infants 0-5
7 months, and (b) Aurangabad: 19 x 11 blocks= 209 infants 0-2
8 months and 209 infants 0-5 months. The 0-5 month old sample
9 is 60% 0-2 months old and 40% 3-5 months old.

10 Using summary data from each of the two samples we analyse
11 the data with Cochran-Mantel-Haenszel (CMH) (27) tests for
12 51 dichotomous indicators (See Supplementary File Table S1)
13 common to the two samples. The CMH tests theoretically have
14 a Chi-square probability distribution with 1 degree of
15 freedom. With a sufficient number of respondents or a
16 sufficient number of blocks the CMH test is equivalent to a
17 conditional logistic regression (28: 114-115). In this
18 analysis both the number of respondents and the number of
19 blocks only approach sufficiency. Consequently, the
20 calculated chi-squares and probabilities must be considered
21 as approximations of their true values.

22 We calculate both unweighted and weighted estimates. The
23 unweighted estimates permit the results from smaller blocks
24 to have equal weight vis à vis larger ones. Since the

1 research question concerns an analysis of which age cohort
2 is most informative, the weighted estimates may not be as
3 useful as the unweighted ones. However, the weighted
4 estimates provide better point estimates of the indicators
5 at the district level. The effect of the weights on the
6 Chi-square statistics is to increase the contribution of
7 the larger blocks and decrease the contribution of the
8 smaller blocks. Hence, we report both sets of results (See
9 Supplementary File Tables S2-S3).

10 The Chi-square probability distribution puts the
11 differences between the districts on a probability scale
12 (See Supplementary File Table S2). To determine meaningful
13 differences in responses between the two age cohorts we
14 used a probability of 0.05 as a cut-off value and consider
15 differences with probabilities less than 0.05 to be
16 possibly meaningful and those with larger probabilities to
17 be likely due to sampling errors. With 102 comparisons (51
18 indicators weighted or unweighted) we must expect some to
19 exceed this cut-off by chance alone. If all of the
20 comparisons were independent, we might randomly find 5
21 differences, but many of the indicators measure related
22 events (e.g., number of ANC visits and tetanus toxoid
23 vaccinations) and the weighted and unweighted estimates are
24 similar, so these indicators are not all independent and it

1 is not possible to calculate an expected number of
2 differences nor is it appropriate to interpret these
3 probabilities as measures of "statistical significance".

4 **Patient and Public Involvement**

5 This study does not involve patients. Also, the public was
6 not involved in the design, conduct and reporting of the
7 research. The public was engaged as interviewees. To
8 ensure local engagement we coordinated with the Bihar
9 Ministry of Health, local implementing non-governmental
10 organizations and our donor. We also shared the results
11 with them and offered further dissemination of results.

12 **RESULTS**

13 We find a high level of agreement between the two samples
14 (Table 1). Out of 102 weighted and unweighted comparisons
15 between the estimates from the 0-2 month and 0-5 month
16 samples there is no probable difference in 81 (79.4%) in
17 both the unweighted and weighted estimates. We detect that
18 probable differences exist for 13 comparisons (12.7%). For
19 the remaining eight comparisons the weighted and unweighted
20 estimates disagree. The weighted estimates find seven
21 differences that the unweighted estimates do not; the
22 unweighted estimates find one difference that the weighted
23 estimates do not find.

1

Table 1. Number of indicators by probability of a difference between the 0-2 months and the 0-5 months samples for weighted and unweighted samples						
Unweighted	Weighted					
	Aurangabad		Gopalganj		Total	
	\geq	$< .05$	\geq	$< .05$	\geq	$< .05$
	.05		.05		.05	
$\geq .05$	40	3	41	4	81	7
$< .05$	0	8	1	5	1	13

2

3

4 For different health service domains, the number of
 5 indicator comparisons varies from two (Exclusive
 6 Breastfeeding-EBF) to 24 concerning home visits by Front
 7 Line Worker (FLW) support (Table 2). The two principal FLW
 8 are Anganwadi workers and Accredited Social Health
 9 Activists (ASHA).

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Table 2. Number of indicator comparisons by subject domain showing a measurable difference using weighted and unweighted estimates of 0-2 months and 0-5 months samples

Health Service Domain	Total comparisons	No measurable difference between 0-2 and 0-5 months results	Measurable difference between 0-2 and 0-5 months results			
			Both	Unweighted only	Weighted only	Percent (%) indicators with different results
Antenatal care	22	21	0	0	1	5
Place of birth & attendant	8	6	1	0	1	25
Birth preparedness	18	13	3	0	2	28
Front Line Worker support	24	16	6	0	2	33
Maternal health	8	8	0	0	0	0
Neonatal care	20	17	1	1	1	15
Exclusive Breastfeeding	2	0	2	0	0	100
Totals	102	81	13	1	7	21.9

1
2
3 1
4
5 2 In the FLW support domain 33% of comparisons have probable
6
7 3 differences. The neonatal health domain has 20 comparisons
8
9 4 and the birth preparedness domain has 18; in these domains
10
11 5 15% and 28% show probable differences, respectively. Place
12
13 6 of birth and attendant, and maternal health each have eight
14
15 7 comparisons with 25%, or two comparisons, and 0
16
17 8 comparisons, respectively, showing a possible difference.
18
19 9 The differences between the two samples cluster around home
20
21 10 visits from FLW and behaviours associated with birth
22
23 11 preparedness and neonatal care. Details of these
24
25 12 differences are listed in Table 3.
26
27 13 For two indicators, both the weighted and unweighted
28
29 14 estimates display probable differences between the 0-2 and
30
31 15 0-5 months samples in both districts. For indicator #52,
32
33 16 the proportion exclusively breastfeeding, the 0-5 months
34
35 17 cohort has the lower estimate, and indicator #24, the
36
37 18 proportion of mothers visited by an ASHA at least once
38
39 19 during their last pregnancy, the 0-5 months sample gives
40
41 20 the higher estimate, about 74%, compared to 63% in the 0-2
42
43 21 months sample (See Supplementary Tables S2-3).
44
45 22
46
47 23 Additional analyses comparing subsamples of mothers of
48
49 24 infants 0-2 months and 3-5 months from the 0-5 months
50
51
52
53
54
55
56
57
58
59
60

1
2
3 1 sample, the sample of mothers of infants 0-2 months and the
4
5 2 subsample of infants 0-2 month, and the sample of mothers
6
7 3 of infants 3-5 months and the subsample of infants 3-5
8
9 4 months produced similar results (See Supplementary Text and
10
11
12 5 Tables S4a-b, Tables S5a-b and Table S6).
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Indicators by health service domain showing measurement differences between 0-2 months and 0-5 months samples

Health Service Domain and Indicator	Indicator No.	District	Weighted Coverage (%)		p-value		Estimate type
			0-2 months	0-5 months	Unweighted Estimate	Weighted Estimate	
Antenatal care							
Proportion of mothers of infants (0-2/0-5 months) registered during their last pregnancy	1	Aurangabad	85.2	77.6	0.0552	0.0365	Weighted
Place of birth & attendant							
Proportion of mothers of infants (0-2/0-5 months) whose last child was delivered at a public facility	38	Gopalganj	51.2	61.6	0.0363	0.0159	Both
Proportion of mothers of infants (0-2/0-5 months) whose last child was delivered at a health facility (private or public facility)	37	Gopalganj	78.9	85.5	0.0643	0.0459	Weighted
Birth preparedness							
Proportion of mothers of infants (0-2/0-5 months) who planned transportation to health facility in their last pregnancy (home & institutional delivery)	15	Gopalganj	45.7	56.0	0.0266	0.0158	Both
Proportion of mothers of infants (0-2/0-5 months) who identified persons to care for the baby immediately after birth (home + institutional delivery)	17	Gopalganj	51.8	62.6	0.0255	0.0103	Both
Proportion of mothers of infants (0-2/0-5 months) who planned for institutional delivery and	23	Aurangabad	62.5	47.0	0.0039	0.0052	Both

1	identified person to accompany her during the delivery							
2								
3	Proportion of mothers who planned for institutional delivery of infants (0-2/0-5 months) who had a new blade & thread for their delivery	19	Aurangabad	23.5	14.5	0.062	0.0429	Weighted
4								
5								
6								
7								
8	Proportion of mothers who planned institutional delivery of infants (0-2/0-5 months) who arranged clean cloth for mothers and baby	21	Aurangabad	43.6	31.2	0.0546	0.0137	Weighted
9								
10								
11								
12								
13								
14	Front Line Worker support							
15	Proportion of mothers of infants (0-2/0-5 months) who were visited by ASHA at least once during their last pregnancy	24	Aurangabad	62.2	75.2	0.0023	0.0042	Both
16			Gopalganj	63.5	73.0	0.0284	0.0175	Both
17								
18								
19	Proportion of mothers of infants (0-2/0-5 months) visited at home by FLWs at least once during their last pregnancy	26	Aurangabad	63.5	76.7	0.0021	0.0032	Both
20								
21								
22								
23								
24	Proportion of mothers of infants (0-2/0-5 months) visited at home by ASHA within 24 hours of last delivery	31	Aurangabad	29.9	44.9	0.0009	0.0016	Both
25								
26								
27								
28	Proportion of mothers of infants (0-2/0-5 months) visited at home by any FLW within 24 hours of last delivery	33	Aurangabad	32.2	46.7	0.0015	0.0026	Both
29								
30								
31								
32	Proportion of mothers of infants (0-2/0-5 months) visited at home by any FLW within first week of last delivery	35	Aurangabad	44.5	59.3	0.0018	0.0026	Both
33								
34								
35								
36	Proportion of mothers of infants (0-2/0-5 months) visited at home by any AWW within the first week of the last delivery	34	Gopalganj	14.4	8.9	0.0959	0.0471	Weighted
37								
38								
39								
40	Proportion of mothers of infants (0-2/0-5 months) visited by	27	Gopalganj	52.6	61.1	0.0617	0.0449	Weighted
41								
42								
43								
44								
45								
46								
47								

1	ASHAs at least once during their last trimester of pregnancy							
2	Infant care							
3	Proportion of infants aged (0-2/0-5 months) who were delivered at home continued with dry cord care	51	Aurangabad	78.0	45.4	0.0001	0.0006	Both
4			Gopalganj	63.7	41.1	0.0431	0.0627	Unweighted
5	Proportion of infants aged (0-2/0-5 months) weighed after birth (Public facility/Private facility/Home)	48	Gopalganj	70.7	78.2	0.0727	0.0464	Weighted
6	Exclusive Breastfeeding							
7	Proportion of infants (0-2/0-5 months) breast-fed in the past 24 hours (Exclusively Breast-Fed)	52	Aurangabad	69.2	59.7	0.0229	0.0411	Both
8			Gopalganj	82.1	68.4	0.0001	0.0003	Both

1
2
3 1
4
56 2 **DISCUSSION**
78
9 3 **Statement of principal findings**
10

11 4 There are no measurable differences in coverage estimates
12
13 5 for 79.4% (81 comparisons) of the indicator comparisons
14
15 6 between the samples of mothers with infants 0-2 months old
16
17 7 versus mothers of infants 0-5 months old; 12.7% (13
18
19 8 comparisons) display measurable differences. The remaining
20
21 9 7.8% (eight comparisons) display discrepancies between the
22
23 10 weighted and unweighted estimates.
24
25
26
27

28 11 **Strengths and weaknesses of the study**
29

30 12 The strengths of this study are that it compares estimates
31
32 13 from two independent samples and that there are many
33
34 14 estimates from diverse domains. The weaknesses of this
35
36 15 study are that the data were collected in only 2 districts
37
38 16 of 1 state in India and in different months of a single
39
40 17 year, and that indicators from the sample of mothers of 3-5
41
42 18 month old infants comparable to those of the 0-2 month old
43
44 19 infants, using the same questionnaire, were not collected.
45
46 20 Supplemental analyses comparing 0-2 and 3-5 month
47
48 21 subsamples of the 0-5 sample did not uncover evidence of
49
50 22 bias due to the combination of these two age groups.
51
52
53
54
55
56
57
58
59
60

1 **Strengths and weaknesses in relation to other studies**

2 Other studies of maternal recall bias have sought a "gold
3 standard" to represent reality and to evaluate measures.
4 Our study, of course, is interested in reality, but this
5 study compares alternative measures needed to assess the
6 Bihar health program. It also uses a complete sample of
7 the age grouping under study rather than just a sub-sample
8 of a larger age grouping. A weakness of this approach is
9 that the analysis does not result in a formal statistical
10 test; our conclusion is based on the weight of the
11 evidence.

12 **Meaning of the study**

13 The evidence indicates that samples of the broader group
14 yield comparable results to those of the narrower age
15 group. It is not necessary to double the total sample by
16 measuring independently 0-2 months and 3-5 months cohorts
17 of children. These results also tend to dispel the
18 hypothesis that maternal recall is problematic for mothers
19 during the first 6 months following delivery. Our results
20 are more consistent with conclusions presented in earlier
21 research (15) and they support those organizations
22 collecting data with 0-5 month cohorts.

1
2
3 1 Indicator #52, EBF, displayed two comparisons measuring
4
5 2 decreases in both districts. This is not surprising as
6
7 3 fewer infants are expected to be exclusively breast fed in
8
9 4 a sample ranging from 0 to 5 months than a sample ranging 0
10
11 to 2 months; mothers introduce complementary feeding and
12
13 5 liquids as infants age despite this being a health risk.
14
15 6 This difference could be accommodated by adjusting
16
17 7 expectations and targets for the indicator.
18
19 8

9 **Unanswered questions and future research**

10 Further investigation and consideration of the differences
11 is warranted. The eight differences found in the FLW
12 support indicators deserve more scrutiny. Seven show higher
13 estimates for the 0-5 cohort and one has a higher estimate
14 for the 0-2 month cohort. The former seven differences may
15 be due to excessive rainfall during the July-September
16 (monthly average 318.95mm, range:195.99-395.8mm) versus the
17 lesser rainfall during June-April (monthly average 25.52mm,
18 range: 0.3-27.88mm) which in the last trimester may have
19 reduced the access of ASHA in the 0-2 month cohort (29).
20 Indicators such as these may be particularly sensitive to
21 rainfall and may explain why more mothers in the 0-5 month
22 cohort displayed higher FLW visitation estimates since FLW

1 were not impeded by the monsoon and the resulting muddy
2 roads.
3 Differences in birth preparedness and institutional birth
4 may be a consequence of differences in rainfall or in FLW
5 support; the results signal a need for more careful
6 planning when transportation is difficult and decreases the
7 effectiveness of FLW by reducing their access to women. Or,
8 some of these differences may just be due to noise in the
9 data.

11 **CONCLUSIONS**

12 Overall, the answer to the research question, "Can one get
13 the same district coverage estimates from a sample of
14 mothers of infants aged 0-5 months as from a sample of
15 mothers of infants ages 0-2 months?" is yes. This result
16 can be paraphrased as: mothers do not display increased
17 recall errors of their perinatal health care behaviour in a
18 cohort of mothers with infants 0-5 months as compared with
19 mothers with younger infants. Substantial resources and
20 effort can be saved using a survey design that avoids
21 needless expenses to collect data that provides
22 insubstantial amounts of information. It also reduces the
23 burden on data collectors and community participants.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Fatigue to both groups can result in needless non-sampling
2 error.

For peer review only

ACKNOWLEDGMENTS

We gratefully acknowledge the essential roles of Hemant Das, Alok Prahdan and Sanjay Biswa for their careful field work supporting the LQAS survey and the HMIS data retrieval. We thank Prof Imelda Bates, Prof Brian Faragher and Nancy Vollmer for their valuable feedback on an earlier version of this manuscript. This work was supported by the Bill and Melinda Gates Foundation (Investment ID OPP1142889).

ETHICS

The Ethical Committees of the Indian Institute of Public Health (No IIPHB-IEC-2016/010) and the Liverpool School of Tropical Medicine Research Ethics Committee approved the protocol, study instruments and consent procedures for the data collection of the household surveys (Research Protocol 16-023).

COMPETING INTERESTS

The authors have read the statement on competing interests and declare that they have none.

AUTHORS' CONTRIBUTIONS

JJV, BD developed the research question and survey design; WH, CJ carried out the statistical analyses; JJV obtained the funding and donor support for the research; BD trained and managed the survey teams in Bihar; JJV, WH, CJ interpreted the data; CJ responsible for data curation; all authors wrote and reviewed the paper.

FUNDING

This research was funded by the Bill and Melinda Gates Foundation Investment ID OPP1142889

DATA SHARING STATEMENT

Data are available upon reasonable request.

REFERENCES

1. UNSTATS. Official list of MDG indicators: United Nations Statistics Division, Department of Economic and Social Affairs; 2005 [Available from: <https://unstats.un.org/unsd/mdg/Host.aspx?Content=Indicators/OfficialList.htm>].
2. UN DESA. Sustainable Development Goals: Division for Sustainable Development Goals, United Nations Department of Economic and Social Affairs; 2015 [Available from: <https://www.un.org/sustainabledevelopment/sustainable-development-goals/>].
3. International Institute for Population Sciences (IIPS) and ICF. National Family Health Survey (NFHS-4), 2015-16: India. Mumbai: IIPS; 2017.
4. MOSPI. Millennium Development Goals India Country Report 2015. New Delhi: Social Statistics Division, Central Statistics Office, Ministry of Statistics and Programme Implementation; 2015.
5. Karvande S, Sonawane D, Chavan S, Mistry N. What does quality of care mean for maternal health providers from two vulnerable states of India? Case study of Bihar and Jharkhand. *Journal of Health, Population and Nutrition*. 2016;35(6):1-10.
6. Mohan P, Kishore B, Singh S, Bahl R, Puri A, Kumar R. Assessment of implementation of integrated management of neonatal and childhood illness in India. *Journal of Health, Population and Nutrition*. 2011;29(6):639-38.
7. Neogi SB, Sharma J, Chauhan M, Khanna R, Chokshi M, Srivastava, et al. Care of newborn in the community and at home. *Journal of Perinatology*. 2016;36(Suppl 3):s13-7.
8. CARE Family Health. Integrated Family Health Initiative: Catalysing change for healthy communities. 2013.
9. Blanc AK, Warren CE, McCarthy KJ, Kimani J, Ndwiga C, RamaRao S. Assessing the validity of indicators of the quality of maternal and newborn health care in Kenya. *Journal of Global Health*. 2016;6(1):1-11.
10. Eisele TP, Rhoda DA, Cutts FT, Keating J, Ren R, Barros AJD, et al. Measuring coverage in MNCH: Total survey error and the interpretation of intervention coverage estimates from household surveys. *PLoS Medicine*. 2013;10(5):e1001386.
11. Anoke SC, Mwai P, Jeffery C, Valadez JJ, Pagano M. Comparing two survey methods of measuring health-related indicators: Lot Quality Assurance Sampling and Demographic Health Surveys. *Tropical Medicine and International Health*. 2015;20(12):1756-70.

12. Valadez JJ, Devkota B, Pradhan MM, Meherda P, Sonal GS, Dhariwal A, et al. Improving malaria treatment and prevention in India by aiding district managers to manage their programmes with local information: a trial assessing the impact of Lot Quality Assurance Sampling on programme outcomes. *Trop Med Int Health*. 2014.
13. Tourangeau R. Remembering what happened: Memory errors and survey reports. In: Stone AA, Jobe JB, Bachrach CA, Cain VS, Kurtzman HS, editors. *The Science of Self-report : Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum; 2000.
14. Bradburn NM. Temporal representation and event dating. In: Stone AA, Jobe JB, Bachrach CA, Cain VS, Kurtzman HS, editors. *The Science of Self-report : Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum; 2000.
15. Tomeo CA, Rich-Edwards JW, Michels KB, Berkey CS, Hunter DJ, Frazier AL, et al. Reproducibility and validity of maternal recall of pregnancy-related events. *Epidemiology*. 1999;10(6):774-7.
16. Elkadry E, Kenton K, White P, Creech S, Brubaker L. Do mothers remember key events during labor? *American Journal of Obstetrics and Gynecology*. 2003;189(1):195-200.
17. Li R, Scanlon KS, Serdula MK. The validity and reliability of maternal recall of breastfeeding practice. *Nutrition Reviews*. 2005;63(4):103-10.
18. Miles M, Ryman TK, Dietz V, Zell E, Luman ET. Validity of vaccination cards and parental recall to estimate vaccination coverage: A systematic review of the literature. *Vaccine*. 2013;31:1560-8.
19. Valadez JJ, Weld LH. Maternal recall error of child vaccination status in a developing nation. *Am J Public Health*. 1992;82(1):120-2.
20. Chan M, Kazatchkine M, Lob-Levyt J, Obaid T, Schweizer J, Sidibe M, et al. Meeting the demand for results and accountability: a call for action on health data from eight global health agencies. *PLoS Med*. 2010;7(1):e1000223.
21. Valadez JJ, Weiss W, Leburg C, Davis R. *Assessing Community Health Programs: A Trainer's Guide. Using LQAS for Baseline Surveys and Regular Monitoring*. 2nd ed. St Albans: Teaching-aids At Low Cost; 2007.
22. Turner AG, Magnani RJ, Shuaib M. A not quite as quick but much cleaner alternative to the Expanded Programme on Immunization (EPI) Cluster Survey design. *International Journal of Epidemiology*. 1996;25(4):198-203.
23. Davis RH, Valadez JJ. Improving the collection of knowledge, attitude and practice data with community surveys: a comparison of two second-stage sampling methods. *Health Policy Planning*. 2014;29(8):1054-60.
24. Dodge HF, Romig HG. *Sampling inspection tables: single and double sampling*. 2nd ed. New York: John Wiley & Sons; 1959.
25. Robertson SE, Joseph J V. Global review of health care surveys using lot quality assurance sampling (LQAS), 1984–2004. *Social Science & Medicine*. 2006;63:1648-60.
26. Valadez JJ. *Assessing child survival programmes in developing countries: testing lot quality assurance sampling*. Boston: Harvard University Press; 1991.
27. Mantel N. Chi-square tests with one degree of freedom: extensions of the Mantel-Haenszel procedure. *J Am Stat Assoc*. 1963;58(303):690-700.
28. Agresti A. *An Introduction to Categorical Data Analysis*. 2nd ed. Hoboken, New Jersey: Wiley-Interscience; John Wiley & Sons; 2007.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

29. World Weather Online. Patna Monthly Climate Averages, Bihar, India 2018. Available from: <https://www.worldweatheronline.com/patna-weather-averages/bihar/in.aspx>.

For peer review only

Supplementary Text for Tables S4a through S6

We make four comparisons to inform an evaluation of these data. The first compares the 0-5 month sample to the 0-2 month sample and is presented in Table 2 in the main text and Tables S2-3. In a second comparison, indicators estimated for the 0-2 month old **subsample** are compared to the 3-5 month old **subsample** to assess the internal consistency of estimates for the entire 0-5 sample (Table 4a-b). In the third comparison, the 0-2 **subsample** of the 0-5 sample is compared to the 0-2 month **sample** to assess the sampling variability for this age group (Tables S5a-b). Finally, in the fourth comparison, the 3-5 **subsample** of the 0-5 sample is compared to the 3-5 month **sample** (Table S6). This last comparison is limited because the 3-5 month sample collected limited data; there are only 3 indicators common to the two samples.

Table S4a has the complete results for the second comparison with point estimates for the two **subsamples** 0-2 and 3-5 of the 0-5 sample, differences and estimated confidence intervals for the estimates and differences of 104 comparisons. Fourteen indicators for which the estimated confidence interval of the difference does not include zero are listed in Table S4b. The table has 2 panels. In the top panel are 6 indicators for which a difference is also reported in Table S2. Table S2 compares the 0-2 month sample and the 0-5 month sample. For each of these indicators the reported difference is in the same direction. Two of these indicators are indicator 52, exclusive infant breast feeding, in the 2 provinces. These differences are not surprising as it is common for mothers to introduce supplemental foods as infants age. The authors speculate in their paper that the timing of the monsoon may have reduced some of the indicators for the younger infants. Of the remaining four indicators of the top panel, three of the indicator differences are negative, lower for the younger infants and might plausibly be related to a monsoon. In the bottom panel of Table S4b are 8 indicators where zero is not in the confidence interval of the difference between subsamples, which suggests difference, but the results between two full-samples in Tables S2 and S3 suggest there is no meaningful difference, except that one of them, indicator 37 in Gopalganj; it does show a difference in Table S2. Of these 8, the difference between samples is in the same direction five times and in a different direction three times. For the 5 indicators where the differences are in the same direction, the absolute value of the differences between 0-2 subsample and the 3-5 subsample are larger than the absolute values of the differences between the 0-2 month sample and the 3-5 subsample. For these 5 indicators the 0-2 month **sample** is more like the 3-5 **subsample** than the 0-2 **subsample**. For the 3 indicators where the differences are in the different directions the 0-2 **subsample** closely resembles the 0-2 month **sample**; two of the estimates are different by less than 1 percent and the third by 2.4 percent.

A third comparison is in Tables S5a-b. In Table S5a the 0-2 **subsample** is compared to the 0-2 month **sample**. The expectation here would be that there are no differences because these two samples are designed to represent the same population. However, this expectation is not met; there are 10 differences where the confidence interval for the difference does not contain 0. These differences are listed in Table S5b. Nine of these 10 indicators are also among the differences in Table S3 and Table 3 in the paper. For 3 of these 9 differences the 3-5 subsample is closer to the 0-2 month sample than it is the 0-2 subsample (indicators 15, 17-2, 51); and for 3 indicators the 3-5 subsample is further from the 0-2 month sample than the 0-2 subsample (indicators 31, 33, 35). For 3 indicators the 0-2 and 3-5 subsample indicator values are nearly equal (indicators 23, 24, 26). For the final indicator in this list (indicator 17-1, the indicator for which zero is in the confidence interval of the difference between the 0-2

1
2
3 month sample and the 0-5 sample in Table S3) the difference between the indicator values for
4 the 0-2 and 3-5 subsamples and the 0-2 month sample are about equal in magnitude but have
5 opposite signs.
6

7
8 Results from the fourth comparison, comparing the 3-5 **subsample** of the 0-5 **sample** to the
9 3-5 month sample are limited, as noted above. There are only 3 indicators in the 2 districts –
10 six comparisons. Zero is within the confidence interval of 5 of the differences (Table S6).
11

12 We note in the paper that in making this number of comparisons one must expect that some
13 will be large enough to be considered meaningful by chance alone. In the above analysis
14 there are 3 comparisons of 51 indicators in 2 provinces producing 16, 14 and 10 differences
15 and a fourth comparison with 1 difference in 6. In their paper the authors find that some of
16 these differences are readily understood and others may be interpreted effects of monsoon
17 rains. Post hoc interpretation is risky here; many of these differences may be noise in the
18 data. There is evidence in the third comparison to support this position with 3 differences
19 moving the 0-5 **sample** closer to the 0-2 month **sample**, 3 moving it away, and 4 not moving
20 it one way or the other. Furthermore, 9 of the 10 differences in the third comparison are also
21 differences in the first comparison, suggesting that about half the differences between the 0-2
22 month sample and the 0-5 sample may be due to differences between the 0-2 month **sample**
23 and the 0-2 **subsample** of the 0-5 sample.
24
25
26

27 The second comparison, comparing the 0-2 subsample to the 3-5 subsample, provides
28 evidence both undermining and supporting the conclusion that a 0-5 sample will provide the
29 same answers as two samples 0-2 and 3-5. On the one hand 5 of the fourteen differences in
30 the third comparison are also in the first, the comparison of the 0-5 sample to the 0-2 sample,
31 suggesting that the inclusion of 3-5 month olds in the 0-5 sample might contribute to the
32 differences. On the other hand, in 5 out of the 8 indicators that are not different between the
33 0-2 and 0-5 samples, the 3-5 subsample more closely matches the 0-2 month sample than
34 does the 0-2 subsample of the 0-5 sample. That this is true in general and not only for these
35 extreme differences is suggested by the mean absolute differences between samples which
36 are 5.2 for the differences between the 0-2 month **sample** and 3-5 month **subsample** and 6.6
37 for the differences between the 0-2 month **subsample** and the 3-5 month **subsample**.
38
39
40

41 Finally, the overall result of all these comparisons is the same as that of the comparison
42 presented in the main text of the paper: in each comparison 85% or more of the indicators
43 show no differences between the samples. The most consistent evidence of difference is
44 between the 0-2 **subsample** of the 0-5 sample and the 0-2 month **sample**.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table S1. Subject domain and indicators labels

Domain ^a	Indicator	Indicator text
1	1	Proportion of mothers of infants (0-2/0-5) months who were registered during their last pregnancy
1	2	Proportion of mothers of infants (0-2/0-5) months whose last pregnancy was registered in the first three months of pregnancy
1	3	Proportion of mothers of infants (0-2/0-5) months attended at least one ANC visit during their last pregnancy
1	4	Proportion of mothers of infants (0-2/0-5) months attended 3 or more ANC visits during their last pregnancy
1	5	Proportion of mothers of infants (0-2/0-5) months attended 4 or more ANC visits during their last pregnancy
1	6	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC visit where BP was checked during her last pregnancy
1	7	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one sonography was performed during her last pregnancy
1	8	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one abdominal examination was performed during her last pregnancy
1	9	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one urine test was performed during her last pregnancy
1	10	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one blood test was performed during her last pregnancy
1	40	Proportion of mothers of infants (0-2/0-5) months who have used an ambulance to reach an institution for last delivery
2	11	Proportion of mothers of infants (0-2/0-5) months who were protected against tetanus in their last pregnancy (Neonatal TT)
2	12	Proportion of mothers of infants (0-2/0-5) months who received two or more doses of TT injection in their last pregnancy
2	13	Proportion of mothers of infants (0-2/0-5) months who received IFA for 100 days or more during their pregnancy
2	14	Proportion of mothers of infants (0-2/0-5) months consumed IFA for 100 days or more in their pregnancy
3	15	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who planned transportation to health facility in their last pregnancy
3	16	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have identified anybody who would donate blood in the case of emergency in their last pregnancy

1			
2	3	17	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have identified persons who would take care of the baby immediately after birth
3			
4	3	18	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have arranged new blade & thread for their last delivery
5			
6	3	19	Proportion of mothers who planned for institutional delivery of infants (0-2/0-5) months who arranged new blade & thread for their delivery
7			
8	3	20	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have arranged clean cloth for mothers and baby
9			
10	3	21	Proportion of mothers who planned institutional delivery of infants (0-2/0-5) months who arranged clean cloth for mothers and baby
11			
12	3	22	Proportion of mothers of infants (0-2/0-5) months who have saved money for the delivery
13			
14	3	23	Proportion of mothers who planned for institutional delivery of infants (0-2/0-5) months identified person to accompany her during the delivery
15			
16	4	24	Proportion of mothers of infants (0-2/0-5) months who were visited by ASHA at least once during their last pregnancy
17			
18	4	25	Proportion of mothers of infants (0-2/0-5) months who were visited by AWW at least once during their last pregnancy
19			
20	4	26	Proportion of mothers of infants (0-2/0-5) months who were visited by FLWs at least once during their last pregnancy
21			
22	4	27	Proportion of mothers of infants (0-2/0-5 months) visited by ASHAs at least once during their last trimester of pregnancy
23			
24	4	28	Proportion of mothers of infants (0-2/0-5) months who were visited by AWW at least once during their last pregnancy
25			
26	4	29	Proportion of mothers of infants (0-2/0-5) months who were visited by any FLW in the last trimester during their last pregnancy
27			
28	4	30	Proportion of mothers of infants (0-2/0-5) months who were visited 2 or more times by any FLW in the last trimester during their last pregnancy
29			
30	4	31	Proportion of mothers of infants (0-2/0-5) months who were visited home by ASHA within 24 hours of last delivery
31			
32	4	32	Proportion of mothers of infants (0-2/0-5) months who were visited home by AWW within 24 hours of last delivery
33			
34	4	33	Proportion of mothers of infants (0-2/0-5) months who were visited home by any FLW within 24 hours of last delivery
35			
36	4	34	Proportion of mothers of infants (0-2/0-5) months who were visited home by any AWW within the first week of last delivery
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			

4	35	Proportion of mothers of infants (0-2/0-5) months who were visited home by any FLW within first week of last delivery
5	37	Proportion of mothers of infants (0-2/0-5) months whose last child was delivered at a health facility (private or public facility)
5	38	Proportion of mothers of infants (0-2/0-5) months whose last child was delivered at public facility
5	39	Proportion of mothers (home delivery) of infants (0-2/0-5) months who had home delivery attended by skilled birth attendant (SBA)
5	39.5	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who had home delivery attended by skill birth attendant (SBA)
6	42	Proportion of infants (home + institutional delivery) aged (0-2/0-5) months with nothing applied to the umbilical cord after cutting and tying in their last delivery
6	43	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have delivered baby practiced skin to skin care (STSC) immediately after birth
6	44	Proportion of mothers with institutional delivery of infants (0-2/0-5) months who continued skin to skin care (STSC) at home
6	45	Proportion of mothers with home delivery of infants (0-2/0-5) months who continued skin to skin care (STSC) later
6	46	Proportion of mothers of infants (0-2/0-5) months who were breastfed within one hour of birth
6	47	Proportion of infants aged (0-2/0-5) months who have received delayed bath (between 48 hours and before 7 days after birth) (Public facility/Private facility/Home)
6	48	Proportion of infants aged (0-2/0-5) months who were weighed after birth (Public facility/Private facility/Home)
6	49	Proportion of infants aged (0-2/0-5) months who were delivered at HF received dry cord care
6	50	Proportion of infants aged (0-2/0-5) months who were delivered at HF continued with dry cord care
6	51	Proportion of infants aged (0-2/0-5) months who were delivered at home continued with dry cord care
7	52	Proportion of infants (0-2/0-5) months who were breast-fed in the past 24 hours (Exclusively Breast-fed)

a. 1 Antenatal care, 2 Maternal health, 3 Birth preparedness, 4 Front Line Worker Support, 6 Place of birth & attendant, 7 Exclusive Breastfeeding

Table S2. Indicators from two samples in Aurangabad and Gopalganj of mothers of infants aged 0-2 months compared to 0-5 months: Estimates, Cochran-Mantel-Haenszel Chi-squares, and probabilities: Bihar 2015

Domain ^a	Indicator ^b	District ^c	Unweighted				Weighted			
			Percent 0-2	Percent 0-5	Chi-square	Probability	Percent 0-2	Percent 0-5	Chi-square	Probability
1	1	1	86.6	79.9	3.6775	0.0552	85.2	77.6	4.3758	0.0365
1	1	2	85.3	85.3	0.0000	1.0000	85.0	85.4	0.0187	0.8911
1	2	1	44	40.7	0.4996	0.4797	43.0	38.3	1.0147	0.3138
1	2	2	47.7	47.0	0.0304	0.8616	47.5	48.2	0.0212	0.8843
1	3	1	84.7	80.9	1.2398	0.2655	83.8	80.4	0.9592	0.3274
1	3	2	81.2	78.6	0.5985	0.4391	80.6	77.8	0.6424	0.4229
1	4	1	60.3	63.6	0.4931	0.4826	61.1	63.1	0.1792	0.6720
1	4	2	59.4	63.2	0.8252	0.3637	57.4	61.0	0.7211	0.3958
1	5	1	42.1	45.0	0.3491	0.5546	43.2	45.7	0.2561	0.6128
1	5	2	44.7	47.7	0.4866	0.4854	43.4	46.9	0.6302	0.4273
1	6	1	59.8	59.3	0.0107	0.9177	58.9	60.1	0.0644	0.7996
1	6	2	67.7	63.5	1.0487	0.3058	66.3	63.3	0.5176	0.4719
1	7	1	42.6	39.2	0.5043	0.4776	42.0	39.5	0.2824	0.5952
1	7	2	70.3	73.3	0.6025	0.4376	69.2	72.6	0.7537	0.3853
1	8	1	44.0	45.9	0.1685	0.6815	42.7	44.8	0.2028	0.6525
1	8	2	53.4	54.1	0.0322	0.8576	52.4	54.5	0.2438	0.6215
1	9	1	45.9	42.6	0.5004	0.4793	45.8	41.7	0.7523	0.3857
1	9	2	42.9	43.6	0.0321	0.8579	42.8	43.5	0.0231	0.8792
1	10	1	67.0	59.8	2.6555	0.1032	67.2	59.6	2.9924	0.0837
1	10	2	55.3	58.3	0.5191	0.4712	54.8	58.5	0.7674	0.3810
1	40	1	11.5	17.7	3.4797	0.0621	10.6	16.0	2.8329	0.0924
1	40	2	8.6	10.2	0.3686	0.5438	8.5	9.5	0.1374	0.7109
2	11	1	98.6	98.1	0.1451	0.7033	98.5	98.1	0.0844	0.7715
2	11	2	97.0	96.6	0.0614	0.8044	96.9	97.1	0.0131	0.9089
2	12	1	68.4	66.5	0.1763	0.6746	69.3	67.5	0.1626	0.6868
2	12	2	79.7	79.3	0.0116	0.9141	81.9	80.4	0.2143	0.6434
2	13	1	3.8	2.4	0.7384	0.3902	4.0	2.6	0.6021	0.4378
2	13	2	3.4	5.6	1.5745	0.2096	3.0	5.1	1.6016	0.2057

Table S2. Indicators from mothers two samples in Aurangabad and Gopalganj of mothers of infants aged 0-2 months compared to 0-5 months: Estimates, Cochran-Mantel-Haenszel Chi-squares, and probabilities: Bihar 2015 - Continued

Domain ^a	Indicator ^b	District ^c	Unweighted				Weighted			
			Percent 0-2	Percent 0-5	Chi- square	Probability	Percent 0-2	Percent 0-5	Chi-square	Probability
2	14	1	1.4	1.9	0.1474	0.7010	1.6	2.0	0.1025	0.7489
2	14	2	1.9	3.8	1.7161	0.1902	1.7	3.6	1.8535	0.1734
3	15	1	26.3	34.0	3.1510	0.0759	25.9	33.3	2.9594	0.0854
3	15	2	47.0	56.4	4.9192	0.0266	45.7	56.0	5.8262	0.0158
3	16	1	4.3	3.8	0.0639	0.8004	4.3	3.6	0.1410	0.7073
3	16	2	10.2	13.9	1.8263	0.1766	9.7	14.3	2.8157	0.0933
3	17	1	48.8	56.5	2.5852	0.1079	48.5	55.6	2.2155	0.1366
3	17	2	53.4	62.8	4.9914	0.0255	51.8	62.6	6.5790	0.0103
3	18	1	24.9	22.5	0.3601	0.5485	26.1	23.4	0.4531	0.5009
3	18	2	39.5	45.1	2.2194	0.1363	38.2	43.7	2.1362	0.1439
3	19	1	22.6	14.4	3.4821	0.0620	23.5	14.5	4.0983	0.0429
3	19	2	45.2	40.7	1.1508	0.2834	43.4	39.4	0.6771	0.4106
3	20	1	40.7	37.8	0.4565	0.4993	44.1	39.5	1.1927	0.2748
3	20	2	51.9	57.1	1.8538	0.1733	50.0	55.9	2.3539	0.1250
3	21	1	40.1	30.4	3.6953	0.0546	43.6	31.2	6.0749	0.0137
3	21	2	58.1	52.8	1.7277	0.1887	55.6	51.6	0.8220	0.3646
3	22	1	65.1	59.3	1.5561	0.2122	65.5	59.4	1.7643	0.1841
3	22	2	62.4	68.4	2.2445	0.1341	60.2	68.0	3.6903	0.0547
3	23	1	65.0	49.0	8.3301	0.0039	62.5	47.0	7.8078	0.0052
3	23	2	68.8	63.0	1.8993	0.1682	68.0	62.5	1.5725	0.2098
4	24	1	61.2	75.1	9.2638	0.0023	62.2	75.2	8.2048	0.0042
4	24	2	65.8	74.4	4.8055	0.0284	63.5	73.0	5.6444	0.0175
4	25	1	11.5	12.9	0.2096	0.6471	11.1	12.5	0.2064	0.6496
4	25	2	20.7	18.4	0.4319	0.5111	19.3	19.1	0.0019	0.9656
4	26	1	62.7	76.6	9.4840	0.0021	63.5	76.7	8.6936	0.0032
4	26	2	68.4	75.9	3.8007	0.0512	66.9	74.3	3.5169	0.0607
4	27	1	42.1	47.8	1.4400	0.2301	42.3	47.6	1.2194	0.2695
4	27	2	54.5	62.4	3.4903	0.0617	52.6	61.1	4.0230	0.0449

Table S2. Indicators from mothers two samples in Aurangabad and Gopalganj of mothers of infants aged 0-2 months compared to 0-5 months: Estimates, Cochran-Mantel-Haenszel Chi-squares, and probabilities: Bihar 2015 - Continued

Domain ^a	Indicator ^b	District ^c	Unweighted				Weighted			
			Percent 0-2	Percent 0-5	Chi-square	Probability	Percent 0-2	Percent 0-5	Chi-square	Probability
4	28	1	8.6	8.1	0.0323	0.8575	8.2	7.6	0.0561	0.8127
4	28	2	16.2	13.9	0.5345	0.4647	15.4	13.8	0.2732	0.6012
4	29	1	43.1	48.8	1.4361	0.2308	43.3	48.5	1.1846	0.2764
4	29	2	57.5	64.7	2.9183	0.0876	55.9	63.5	3.2352	0.0721
4	30	1	29.7	35.4	1.6374	0.2007	28.7	34.2	1.5497	0.2132
4	30	2	48.1	53	1.2686	0.2600	46.9	51.4	1.0908	0.2963
4	31	1	30.1	45.9	10.9581	0.0000	29.9	44.9	9.9713	0.0016
4	31	2	47.4	53.0	1.6849	0.1943	44.5	52.7	3.6136	0.0573
4	32	1	5.3	3.8	0.4948	0.4818	5.3	3.6	0.7401	0.3896
4	32	2	9.8	9.0	0.0897	0.7646	9.1	9.9	0.0974	0.7550
4	33	1	32.5	47.8	10.1305	0.0015	32.2	46.7	9.0939	0.0026
4	33	2	50.8	56.4	1.7133	0.1906	48.2	56.2	3.5264	0.0604
4	34	1	6.2	10.0	2.0808	0.1492	6.2	9.9	1.9617	0.1613
4	34	2	13.5	9.0	2.7721	0.0959	14.4	8.9	3.9431	0.0471
4	35	1	45.0	60.3	9.7398	0.0018	44.5	59.3	9.0930	0.0026
4	35	2	64.7	69.5	1.4491	0.2287	62.5	69.0	2.5201	0.1124
5	37	1	73.2	76.6	0.6496	0.4203	72.9	74.9	0.2182	0.6404
5	37	2	80.1	86.1	3.4220	0.0643	78.9	85.5	3.9858	0.0459
5	38	1	55.5	56.5	0.0389	0.8436	54.7	55.3	0.0117	0.9139
5	38	2	52.3	61.3	4.3834	0.0363	51.2	61.6	5.8090	0.0159
5	39	1	5.7	0	3.3034	0.0691	5.4	0	3.6637	0.0556
5	39	2	10.2	8.3	0.2393	0.6247	9.1	6.6	0.3451	0.5569
5	39.5	1	1.4	0	0.0003	0.0833	1.4	0	2.8978	0.0887
5	39.5	2	1.9	1.1	0.5139	0.4735	1.8	0.9	0.7629	0.3824
6	42	1	42.6	47.4	1.0165	0.3134	43.8	47.6	0.6553	0.4182
6	42	2	41.4	46.2	1.3134	0.2518	41.8	46.3	1.1228	0.2893
6	43	1	24.9	29.7	1.3262	0.2495	25.6	30.4	1.3008	0.2541
6	43	2	27.1	29.7	0.4968	0.4809	26.9	31.2	1.2586	0.2619

Table S2. Indicators from mothers two samples in Aurangabad and Gopalganj of mothers of infants aged 0-2 months compared to 0-5 months: Estimates, Cochran-Mantel-Haenszel Chi-squares, and probabilities: Bihar 2015 - Continued										
Domain ^a	Indicator ^b	District ^c	Unweighted				Weighted			
			Percent 0-2	Percent 0-5	Chi-square	Probability	Percent 0-2	Percent 0-5	Chi-square	Probability
6	44	1	9.8	12.5	1.2833	0.2573	12.2	13.0	0.3958	0.5293
6	44	2	21.1	20.1	0.0586	0.8087	21.6	18.7	0.5178	0.4718
6	45	1	7.5	10.2	0.6321	0.4266	7.8	11.1	0.8502	0.3565
6	45	2	8.2	5.6	0.4009	0.5266	5.2	8.3	0.1670	0.6828
6	46	1	55.5	58.9	0.4904	0.4837	56.6	59.2	0.2960	0.5864
6	46	2	62.8	68.0	1.6764	0.1954	61.7	66.3	1.2736	0.2591
6	47	1	34.0	32.1	0.1768	0.6741	35.9	32.8	0.4369	0.5086
6	47	2	67.3	66.5	0.0344	0.8528	66.6	68.3	0.1745	0.6762
6	48	1	65.6	67.5	0.1782	0.6729	64.9	66.6	0.1312	0.7172
6	48	2	71.1	77.8	3.2208	0.0727	70.7	78.2	3.9664	0.0464
6	49	1	62.7	59.4	0.1537	0.6950	64.5	61.2	0.1691	0.6809
6	49	2	68.1	67.2	0.0166	0.8974	67.3	66.8	0.0011	0.9735
6	50	1	49.0	55.0	1.2527	0.2630	48.4	54.7	1.3188	0.2508
6	50	2	56.8	55.9	0.0964	0.7562	56.1	55.5	0.0615	0.8042
6	51	1	81.1	44.9	14.9736	0.0000	78.0	45.4	11.7395	0.0000
6	51	2	63.3	38.9	4.0901	0.0431	63.7	41.1	3.4654	0.0627
7	52	1	70.3	59.8	5.1727	0.0229	69.2	59.7	4.1723	0.0411
7	52	2	82.3	67.3	16.2014	0.0000	82.1	68.4	13.3711	0.0000
a . 1 Antenatal care, 2 Maternal health, 3 Birth preparedness, 4 FLW Support, 5 Place of birth & attendant, 6 Neonatal Health, 7 Exclusive breastfeeding										
b . For text see Table S1										
c. 1 Aurangabad, 2 Gopalganj										

Indicator ^a	District ^b	0-2 Month Sample			0-5 Month Sample			Difference		
		Point estimate	Confidence interval ^c		Point estimate	Confidence interval ^c		Point estimate	Confidence interval	
			Lower	Upper		Lower	Upper		Lower	Upper
1	1	85.2	80.2	90.2	77.6	72.6	82.6	7.6	0.1	15.2
1	2	85.0	80.5	89.6	85.4	80.9	90.0	-0.4	-6.7	5.9
2	1	43.0	36.3	49.8	38.3	31.5	45.0	4.7	-4.5	14.0
2	2	47.5	41.5	53.6	48.2	42.1	54.2	-0.6	-9.5	8.2
3	1	83.8	78.8	88.8	80.4	75.4	85.4	3.4	-3.6	10.4
3	2	80.6	75.7	85.5	77.8	72.9	82.7	2.8	-4.5	10.0
4	1	61.1	54.4	67.8	63.1	56.4	69.9	-2.0	-11.4	7.3
4	2	57.4	51.2	63.6	61.0	54.8	67.2	-3.5	-12.3	5.2
5	1	43.2	36.4	50.1	45.7	38.9	52.6	-2.5	-12.2	7.2
5	2	43.4	37.1	49.8	46.9	40.5	53.2	-3.4	-12.4	5.6
6	1	58.9	52.3	65.6	60.1	53.5	66.7	-1.2	-10.4	8.0
6	2	66.3	60.2	72.4	63.3	57.2	69.4	2.9	-5.7	11.6
7	1	42.0	35.3	48.8	39.5	32.7	46.3	2.5	-7.0	12.1
7	2	69.2	63.2	75.2	72.6	66.6	78.6	-3.4	-11.7	4.9
8	1	42.7	36.1	49.3	44.8	38.2	51.4	-2.1	-11.4	7.2
8	2	52.4	46.2	58.7	54.5	48.2	60.8	-2.1	-10.9	6.7
9	1	45.8	39.0	52.6	41.7	34.9	48.4	4.1	-5.4	13.6
9	2	42.8	36.6	49.1	43.5	37.3	49.7	-0.6	-9.5	8.2
10	1	67.2	60.9	73.6	59.6	53.2	65.9	7.6	-1.2	16.5
10	2	54.8	48.5	61.1	58.5	52.2	64.8	-3.7	-12.5	5.2
11	1	98.5	96.7	100.2	98.1	96.4	99.9	0.4	-2.2	2.9
11	2	96.9	94.8	99.1	97.1	94.9	99.3	-0.2	-3.0	2.7
12	1	69.3	63.0	75.7	67.5	61.2	73.9	1.8	-7.2	10.8
12	2	81.9	77.3	86.6	80.4	75.7	85.0	1.6	-5.2	8.4
13	1	4.0	1.2	6.7	2.6	-0.1	5.4	1.3	-2.2	4.9
13	2	3.0	1.0	5.0	5.1	3.1	7.1	-2.2	-5.5	1.2
14	1	1.6	-0.2	3.4	2.0	0.2	3.8	-0.4	-3.1	2.3
14	2	1.7	0.2	3.2	3.6	2.1	5.1	-1.9	-4.7	0.9
15	1	25.9	20.1	31.6	33.3	27.5	39.0	-7.4	-15.9	1.1
15	2	45.7	39.6	51.8	56.0	49.9	62.1	-10.3	-19.0	-1.5
16	1	4.3	1.5	7.0	3.6	0.8	6.3	0.7	-3.0	4.4
16	2	9.7	6.1	13.3	14.3	10.8	17.9	-4.7	-10.5	1.1
17	1	48.5	41.8	55.2	55.6	48.9	62.3	-7.1	-16.5	2.4
17	2	51.8	45.6	58.0	62.6	56.4	68.8	-10.8	-19.6	-2.1
18	1	26.1	20.2	32.1	23.4	17.5	29.4	2.7	-5.5	10.9
18	2	38.2	33.0	43.3	43.7	38.5	48.8	-5.5	-13.2	2.2
19	1	23.5	16.4	30.7	14.5	7.3	21.7	9.1	0.4	17.7
19	2	43.4	37.3	49.5	39.4	33.3	45.5	4.0	-4.6	12.6
20	1	44.1	38.3	50.0	39.5	33.7	45.4	4.6	-3.7	12.9
20	2	50.0	44.7	55.4	55.9	50.5	61.3	-5.9	-13.7	2.0
21	1	43.6	36.7	50.5	31.2	24.3	38.1	12.4	3.3	21.5
21	2	55.6	49.5	61.7	51.6	45.5	57.7	4.0	-4.6	12.6
22	1	65.5	59.0	72.0	59.4	52.9	65.9	6.1	-3.0	15.2
22	2	60.2	54.0	66.3	68.0	61.8	74.2	-7.8	-16.4	0.7
23	1	62.5	54.9	70.2	47.0	39.3	54.7	15.5	5.4	25.7
23	2	68.0	61.2	74.8	62.5	55.7	69.3	5.5	-3.8	14.8

1	24	1	62.2	55.5	68.9	75.2	68.5	81.9	-13.0	-22.0	-4.1
2	24	2	63.5	57.4	69.7	73.0	66.9	79.2	-9.5	-17.9	-1.1
3	25	1	11.1	7.0	15.1	12.5	8.4	16.5	-1.4	-7.5	4.7
4	25	2	19.3	14.4	24.2	19.1	14.3	24.0	0.1	-6.9	7.2
5	26	1	63.5	56.8	70.1	76.7	70.1	83.4	-13.2	-22.1	-4.4
6	26	2	66.9	60.9	73.0	74.3	68.3	80.3	-7.4	-15.7	1.0
7	27	1	42.3	35.6	49.1	47.6	40.8	54.4	-5.3	-14.8	4.3
8	27	2	52.6	46.3	58.9	61.1	54.8	67.4	-8.5	-17.2	0.3
9	28	1	8.2	4.6	11.8	7.6	4.0	11.2	0.6	-4.4	5.7
10	28	2	15.4	11.0	19.8	13.8	9.4	18.3	1.6	-4.7	7.8
11	29	1	43.3	36.5	50.1	48.5	41.7	55.3	-5.2	-14.8	4.3
12	29	2	55.9	49.7	62.2	63.5	57.2	69.8	-7.5	-16.3	1.2
13	30	1	28.7	22.6	34.7	34.2	28.1	40.2	-5.5	-14.3	3.2
14	30	2	46.9	40.5	53.3	51.4	45.0	57.8	-4.5	-13.5	4.5
15	31	1	29.9	23.5	36.3	44.9	38.5	51.3	-15.0	-24.4	-5.6
16	31	2	44.5	38.2	50.8	52.7	46.5	59.0	-8.2	-17.2	0.7
17	32	1	5.3	2.2	8.5	3.6	0.4	6.7	1.7	-2.3	5.8
18	32	2	9.1	5.7	12.6	9.9	6.4	13.4	-0.8	-6.0	4.4
19	33	1	32.2	25.8	38.7	46.7	40.2	53.2	-14.5	-23.9	-5.0
20	33	2	48.2	41.9	54.5	56.2	49.9	62.5	-8.1	-16.9	0.8
21	34	1	6.2	2.8	9.6	9.9	6.5	13.3	-3.7	-8.9	1.5
22	34	2	14.4	9.8	19.0	8.9	4.4	13.5	5.5	-0.3	11.3
23	35	1	44.5	37.7	51.4	59.3	52.5	66.2	-14.8	-24.4	-5.1
24	35	2	62.5	56.4	68.7	69.0	62.9	75.2	-6.5	-15.1	2.1
25	37	1	72.9	66.7	79.1	74.9	68.7	81.0	-2.0	-10.5	6.5
26	37	2	78.9	73.5	84.2	85.5	80.2	90.8	-6.6	-13.7	0.4
27	38	1	54.7	47.8	61.7	55.3	48.4	62.2	-0.5	-10.3	9.2
28	38	2	51.2	44.7	57.6	61.6	55.2	68.0	-10.4	-19.4	-1.5
29	39	1	5.4	-0.9	11.8	0.0	-6.4	6.4	5.4	-0.9	11.8
30	39	2	9.1	1.3	16.9	6.6	-1.2	14.4	2.5	-9.1	14.1
31	39.5	1	1.4	-0.2	3.0	0.0	-1.6	1.6	1.4	-0.2	3.0
32	39.5	2	1.8	0.2	3.4	0.9	-0.6	2.5	0.9	-1.0	2.8
33	40	1	10.6	6.5	14.7	16.0	12.0	20.1	-5.4	-11.7	0.8
34	40	2	8.5	4.9	12.2	9.5	5.8	13.1	-0.9	-5.9	4.1
35	42	1	43.8	36.9	50.6	47.6	40.7	54.5	-3.8	-13.3	5.6
36	42	2	41.8	35.6	48.1	46.3	40.1	52.6	-4.5	-13.4	4.4
37	43	1	25.6	19.8	31.5	30.4	24.6	36.2	-4.8	-13.3	3.7
38	43	2	26.9	21.5	32.4	31.2	25.7	36.6	-4.2	-12.0	3.5
39	44	1	12.2	7.3	17.1	13.0	8.1	17.9	-0.8	-7.7	6.1
40	44	2	21.6	16.3	26.8	18.7	13.4	24.0	2.8	-4.4	10.1
41	45	1	7.8	0.4	15.2	11.1	3.7	18.5	-3.3	-16.2	9.6
42	45	2	5.2	0.3	10.1	8.3	3.4	13.2	-3.1	-17.4	11.2
43	46	1	56.6	49.9	63.2	59.2	52.5	65.8	-2.6	-12.2	7.0
44	46	2	61.7	55.5	67.9	66.3	60.1	72.6	-4.6	-13.3	4.1
45	47	1	35.9	29.3	42.5	32.8	26.2	39.5	3.0	-6.2	12.3
46	47	2	66.6	60.6	72.7	68.3	62.3	74.4	-1.7	-10.1	6.7
47	48	1	64.9	58.5	71.4	66.6	60.1	73.0	-1.6	-10.8	7.5
48	48	2	70.7	64.9	76.5	78.2	72.4	84.0	-7.5	-15.2	0.2
49	49	1	64.5	57.1	71.8	61.2	53.9	68.5	3.3	-6.7	13.3
50	49	2	67.3	60.6	73.9	66.8	60.1	73.4	0.5	-8.9	9.9
51	50	1	48.4	40.3	56.4	54.7	46.7	62.8	-6.4	-17.4	4.7
52	50	2	56.1	49.0	63.2	55.5	48.4	62.6	0.7	-9.2	10.5

1	51	1	78.0	65.9	90.2	45.4	33.3	57.6	32.6	13.0	52.2
2	51	2	63.7	48.8	78.5	41.1	26.2	56.0	22.5	-2.2	47.2
3	52	1	69.2	62.7	75.6	59.7	53.3	66.1	9.5	0.2	18.8
4	52	2	82.1	77.1	87.0	68.4	63.5	73.4	13.6	6.0	21.2
5	a. For text see Table S1										
6	b. 1 Aurangabad, 2 Gopalganj										
7	c. Estimated with Stata command svy										
8											
9											
10											
11											
12											
13											
14											
15											
16											
17											
18											
19											
20											
21											
22											
23											
24											
25											
26											
27											
28											
29											
30											
31											
32											
33											
34											
35											
36											
37											
38											
39											
40											
41											
42											
43											
44											
45											
46											
47											
48											
49											
50											
51											
52											
53											
54											
55											
56											
57											
58											
59											
60											

Table S4a: Point estimates and confidence intervals of indicators for subsamples of 0-2 and 3-5 month old infants in two districts: Bihar, India, 2015

Indicator ^a	District ^b	0-2 month subsample			3-5 month subsample			Difference		
		Point estimate	Confidence interval ^c		Point estimate	Confidence interval ^c		Point estimate	Confidence interval	
			Lower	Upper		Lower	Upper		Lower	Upper
1	1	79.2	71.9	86.5	75.2	66.0	84.4	4.0	-7.7	15.7
1	2	80.8	74.4	87.2	92.3	87.5	97.2	-11.5	-19.5	-3.5
2	1	36.1	27.8	44.4	41.5	31.7	51.3	-5.4	-18.2	7.4
2	2	46.5	38.1	54.8	50.6	40.3	61.0	-4.2	-17.5	9.1
3	1	81.0	75.0	87.1	79.5	71.5	87.5	1.5	-8.5	11.6
3	2	79.9	73.4	86.5	74.7	65.8	83.7	5.2	-5.9	16.3
4	1	61.8	53.1	70.4	65.2	55.0	75.3	-3.4	-16.7	10.0
4	2	62.7	54.7	70.7	58.4	48.4	68.3	4.4	-8.4	17.1
5	1	45.0	36.2	53.9	46.8	35.5	58.0	-1.7	-16.0	12.6
5	2	49.1	40.8	57.4	43.5	33.3	53.7	5.6	-7.6	18.7
6	1	60.3	51.9	68.7	59.8	49.6	70.0	0.5	-12.7	13.7
6	2	68.5	60.8	76.1	55.8	45.7	65.8	12.7	0.0	25.4
7	1	40.3	31.8	48.8	38.4	27.4	49.3	1.9	-11.9	15.8
7	2	74.5	67.2	81.9	69.8	60.3	79.3	4.7	-7.3	16.8
8	1	49.3	41.2	57.3	38.2	27.9	48.5	11.1	-2.0	24.2
8	2	60.0	52.0	67.9	46.4	36.7	56.2	13.6	1.0	26.1
9	1	39.8	31.3	48.4	44.4	33.9	54.8	-4.5	-18.0	9.0
9	2	44.7	36.6	52.8	41.7	31.5	51.9	3.0	-10.0	16.0
10	1	61.9	54.4	69.4	56.1	45.3	66.9	5.7	-7.4	18.9
10	2	63.9	56.1	71.7	50.5	40.3	60.6	13.5	0.7	26.2
11	1	98.4	96.3	100.6	97.6	94.3	100.9	0.8	-3.1	4.8
11	2	96.7	94.1	99.3	97.7	95.1	100.3	-1.0	-4.7	2.7
12	1	65.0	56.4	73.6	71.3	61.8	80.7	-6.3	-19.1	6.5
12	2	77.8	71.1	84.6	84.1	76.7	91.4	-6.2	-16.2	3.8
13	1	3.3	0.1	6.5	1.6	-1.5	4.8	1.7	-2.8	6.2
13	2	6.7	2.6	10.9	2.8	0.0	5.6	3.9	-1.1	8.9
14	1	2.2	-0.4	4.8	1.6	-1.5	4.8	0.6	-3.5	4.7
14	2	4.5	1.0	8.0	2.3	-0.4	4.9	2.2	-2.2	6.6
15	1	35.2	26.8	43.6	30.4	20.9	39.9	4.8	-7.8	17.5
15	2	56.8	48.7	64.9	54.8	44.3	65.3	2.0	-11.3	15.2
16	1	5.3	1.4	9.3	0.9	-0.8	2.6	4.5	0.2	8.7
16	2	16.6	10.2	23.0	11.0	4.8	17.1	5.7	-3.2	14.5
17	1	59.6	51.4	67.9	49.5	38.7	60.3	10.1	-3.5	23.7
17	2	64.4	56.5	72.4	59.9	49.8	70.0	4.5	-8.3	17.4
18	1	24.2	16.8	31.6	22.2	13.9	30.5	2.0	-9.1	13.1
18	2	45.9	38.1	53.8	40.3	32.0	48.6	5.7	-5.7	17.1
19	1	17.9	11.2	24.5	9.2	2.4	16.1	8.6	-0.9	18.2
19	2	40.5	32.4	48.6	37.7	29.0	46.4	2.8	-9.0	14.6
20	1	42.5	34.8	50.2	35.1	26.0	44.2	7.4	-4.5	19.4
20	2	59.4	51.7	67.0	50.7	42.3	59.1	8.6	-2.7	20.0
21	1	37.0	29.2	44.8	22.1	13.1	31.2	14.9	2.9	26.8
21	2	54.2	46.1	62.3	47.7	38.9	56.6	6.5	-5.6	18.5
22	1	62.5	54.0	70.9	54.7	44.8	64.6	7.8	-5.2	20.8
22	2	69.7	62.3	77.1	65.5	55.4	75.5	4.2	-8.2	16.7
23	1	47.5	39.1	55.9	46.1	35.0	57.3	1.4	-12.5	15.3

1	23	2	64.9	56.9	73.0	58.9	48.4	69.5	6.0	-7.3	19.3
2	24	1	75.1	67.4	82.9	75.3	66.0	84.7	-0.2	-12.4	11.9
3	24	2	67.6	59.7	75.5	81.0	72.7	89.3	-13.4	-24.8	-1.9
4	25	1	12.1	6.1	18.0	13.1	5.9	20.2	-1.0	-10.3	8.3
5	25	2	16.7	10.3	23.1	22.7	13.8	31.6	-6.0	-16.9	4.9
6	26	1	76.2	68.6	83.8	77.5	68.5	86.6	-1.3	-13.1	10.5
7	26	2	69.4	61.6	77.2	81.6	73.3	89.8	-12.2	-23.6	-0.8
8	27	1	50.4	41.5	59.3	43.4	33.0	53.9	6.9	-6.8	20.7
9	27	2	57.2	48.9	65.4	66.9	57.7	76.0	-9.7	-22.0	2.7
10	28	1	8.6	3.7	13.6	6.0	1.3	10.6	2.7	-4.1	9.5
11	28	2	9.8	5.1	14.5	19.7	11.4	28.1	-9.9	-19.5	-0.3
12	29	1	50.4	41.5	59.3	45.6	35.4	55.9	4.8	-8.8	18.4
13	29	2	59.2	51.1	67.4	69.7	60.5	79.0	-10.5	-22.9	1.8
14	30	1	32.7	24.5	41.0	36.3	25.9	46.7	-3.6	-16.9	9.7
15	30	2	49.2	41.0	57.4	54.7	44.3	65.1	-5.5	-18.8	7.7
16	31	1	42.0	33.0	51.1	49.2	39.0	59.5	-7.2	-20.9	6.5
17	31	2	48.6	40.3	56.8	58.9	48.8	69.0	-10.3	-23.3	2.7
18	32	1	3.9	0.4	7.3	3.2	-0.4	6.8	0.7	-4.3	5.7
19	32	2	9.4	4.2	14.5	10.7	4.7	16.8	-1.4	-9.3	6.5
20	33	1	43.6	34.5	52.6	51.4	41.1	61.7	-7.8	-21.6	5.9
21	33	2	52.9	44.8	61.0	61.1	51.1	71.2	-8.2	-21.1	4.7
22	34	1	7.0	2.5	11.5	14.2	7.0	21.4	-7.2	-15.7	1.2
23	34	2	8.6	3.8	13.4	9.5	4.0	15.0	-0.9	-8.2	6.4
24	35	1	57.2	48.2	66.2	62.5	52.7	72.4	-5.3	-18.7	8.0
25	35	2	63.2	55.1	71.4	77.6	68.6	86.6	-14.3	-26.5	-2.2
26	37	1	76.6	69.1	84.1	72.3	62.7	81.8	4.3	-7.8	16.4
27	37	2	81.6	74.9	88.3	91.3	85.8	96.8	-9.7	-18.4	-1.0
28	38	1	59.0	49.9	68.0	49.7	39.3	60.2	9.2	-4.6	23.1
29	38	2	56.1	47.8	64.4	69.7	61.2	78.2	-13.6	-25.5	-1.7
30	39	1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
31	39	2	5.8	-3.1	14.7	9.6	-40.1	59.3	-3.8	-54.3	46.7
32	39.5	1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
33	39.5	2	1.1	-0.4	2.6	0.7	-0.7	2.1	0.3	-1.7	2.4
34	40	1	17.6	10.9	24.4	13.6	7.5	19.8	4.0	-5.2	13.2
35	40	2	8.0	3.9	12.1	11.6	5.6	17.7	-3.7	-11.0	3.6
36	42	1	52.6	44.3	61.0	40.1	29.6	50.6	12.5	-0.9	25.9
37	42	2	47.5	39.4	55.6	44.6	34.4	54.8	2.9	-10.2	16.0
38	43	1	35.6	27.4	43.8	22.6	13.6	31.5	13.1	0.9	25.2
39	43	2	35.5	28.0	43.0	24.8	16.4	33.1	10.8	-0.5	22.0
40	44	1	10.1	4.2	16.0	17.6	9.5	25.7	-7.5	-17.6	2.5
41	44	2	20.6	14.1	27.0	16.2	8.7	23.8	4.3	-5.6	14.3
42	45	1	9.0	-4.7	22.7	13.7	-0.7	28.1	-4.7	-24.6	15.2
43	45	2	6.6	-9.9	23.2	14.4	14.4	14.4	-7.7	-24.3	8.8
44	46	1	63.4	54.6	72.2	52.8	41.5	64.0	10.7	-3.6	24.9
45	46	2	68.9	61.1	76.7	62.5	52.6	72.4	6.4	-6.2	19.0
46	47	1	33.0	24.7	41.4	32.5	22.2	42.9	0.5	-12.8	13.8
47	47	2	70.4	62.8	77.9	65.3	55.9	74.8	5.1	-7.1	17.2
48	48	1	68.8	60.6	77.0	63.2	52.5	73.8	5.7	-7.8	19.1
49	48	2	76.2	69.3	83.0	81.3	73.9	88.6	-5.1	-15.2	5.0
50	49	1	66.5	57.9	75.1	52.7	41.0	64.3	13.8	-0.6	28.3
51	49	2	68.6	59.9	77.3	64.4	53.8	75.0	4.2	-9.5	17.9
52	50	1	51.7	42.1	61.4	59.5	47.6	71.5	-7.8	-23.1	7.6

1	50	2	55.2	46.3	64.1	55.8	45.1	66.6	-0.7	-14.6	13.3
2	51	1	33.2	14.6	51.7	61.0	36.7	85.4	-27.9	-58.5	2.7
3	51	2	40.6	17.5	63.7	43.0	-6.7	92.7	-2.4	-57.2	52.4
4	52	1	73.0	65.1	80.9	39.7	28.6	50.7	33.3	19.8	46.9
5	52	2	83.8	78.2	89.3	45.8	35.6	55.9	38.0	26.4	49.5

6 a. For text see Table S1

7 b. 1 Aurangabad, 2 Gopalganj

8 c. Estimated with Stata command svy

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table S4b: Indicators in Table S4a where zero is not in the estimated confidence interval of difference between 0-2 and 3-5 month subsamples							
Indicator	District ^a	Point estimate		Difference	Confidence interval ^b		
		0-2 month	3-5 month		Lower	Upper	
In Table S4a and in Table S2							
21	Proportion of mothers who planned institutional delivery of infants (0-2/0-5) months who arranged clean cloth for mothers and baby	1	37.0	22.1	14.9	2.9	26.8
24	Proportion of mothers of infants (0-2/0-5) months who were visited by ASHA at least once during their last pregnancy	2	67.6	81.0	-13.4	-24.8	-1.9
37	Proportion of mothers of infants (0-2/0-5) months whose last child was delivered at a health facility (private or public facility)	2	81.6	91.3	-9.7	-18.4	-1.0
38	Proportion of mothers of infants (0-2/0-5) months whose last child was delivered at public facility	2	56.1	69.7	-13.6	-25.5	-1.7
52	Proportion of infants (0-2/0-5) months who were breast-fed in the past 24 hours (Exclusively Breast-fed)	1	73.0	39.7	33.3	19.8	46.9
52		2	83.8	45.8	38.0	26.4	49.5
In Table S4a but not in Table S2							
1	Proportion of mothers of infants (0-2/0-5) months who were registered during their last pregnancy	2	80.8	92.3	-11.5	-19.5	-3.5
8	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one abdominal examination was performed during her last pregnancy	2	60.0	46.4	13.6	1.0	26.1
10	Proportion of mothers of infants (0-2/0-5) months who attended at least one ANC where at least one blood test was performed during her last pregnancy	2	63.9	50.5	13.5	0.7	26.2
16	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have identified anybody who would donate blood in the case of emergency in their last pregnancy	1	5.3	0.9	4.5	0.2	8.7
26	Proportion of mothers of infants (0-2/0-5) months who were visited by FLWs at least once during their last pregnancy	2	69.4	81.6	-12.2	-23.6	-0.8
28	Proportion of mothers of infants (0-2/0-5) months who were visited by AWW in the last trimester during their last pregnancy	2	9.8	19.7	-9.9	-19.5	-0.3
35	Proportion of mothers of infants (0-2/0-5) months who were visited home by any FLW within first week of last delivery	2	63.2	77.6	-14.3	-26.5	-2.2

1	43	Proportion of mothers (home + institutional delivery) of infants (0-2/0-5)	1	35.6	22.6	13.1	0.9	25.2
2		months who have delivered baby practiced skin to skin care (STSC)						
3		immediately after birth						
4	a.	1 = Aurangabad; 2 = Gopalganj						
5	b.	Calculated from standard errors for point estimates estimated with Stata command svy						

For peer review only

Table S5b: Point estimates of weighted indicators and differences between a sample of mothers of 0-2 month old infants and a subsample of mothers of 0-2 month old infants from a sample of mothers of 0-5 month old infants in two districts of Bihar, India

Indicator	District ^a	Point estimates		Difference		
		Subsample	Sample	Estimate	Confidence interval ^b	
					Lower	Upper
15 Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who planned transportation to health facility in their last pregnancy	2	56.8	45.7	11.1	0.9	21.2
17 Proportion of mothers (home + institutional delivery) of infants (0-2/0-5) months who have identified persons who would take care of the baby immediately after birth	1	59.6	48.5	11.1	0.5	21.7
	2	64.4	51.8	12.7	2.6	22.7
23 Proportion of mothers who planned for institutional delivery of infants (0-2/0-5) months identified person to accompany her during the delivery	1	47.5	62.5	-15.0	-26.4	-3.6
24 Proportion of mothers of infants (0-2/0-5) months who were visited by ASHA at least once during their last pregnancy	1	75.1	62.2	12.9	2.7	23.2
26 Proportion of mothers of infants (0-2/0-5) months who were visited by FLWs at least once during their last pregnancy	1	76.2	63.5	12.7	2.6	22.8
31 Proportion of mothers of infants (0-2/0-5) months who were visited home by ASHA within 24 hours of last delivery	1	42.0	29.9	12.1	1.1	23.2
33 Proportion of mothers of infants (0-2/0-5) months who were visited home by any FLW within 24 hours of last delivery	1	43.6	32.2	11.3	0.2	22.5
35 Proportion of mothers of infants (0-2/0-5) months who were visited home by any FLW within first week of last delivery	1	57.2	44.5	12.7	1.3	24.0
51 Proportion of infants aged (0-2/0-5) months who were delivered at home continued with dry cord care	1	33.2	78.0	-44.9	-67.0	-22.7

a. 1 = Aurangabad; 2 = Gopalganj
 b. Calculated from standard errors for point estimates estimated with Stata command svy

Table S5a: Point estimates and confidence intervals of indicators for a sample 0-2 month old infants and a subsample of 0-2 month old infants from a sample of 0-5 month old infants in two districts: Bihar, India, 2015

Indicator ^a	District ^b	0-2 month subsample			0-2 month sample			Difference		
		Point estimate	Confidence interval ^c		Point estimate	Confidence interval ^c		Point estimate	Confidence interval	
			Lower	Upper		Lower	Upper		Lower	Upper
1	1	79.2	71.9	86.5	85.2	80.2	90.2	-6.0	-14.8	2.8
1	2	80.8	74.4	87.2	85.0	80.5	89.6	-4.2	-12.1	3.6
2	1	36.1	27.8	44.4	43.0	36.3	49.8	-6.9	-17.6	3.8
2	2	46.5	38.1	54.8	47.5	41.5	53.6	-1.1	-11.3	9.2
3	1	81.0	75.0	87.1	83.8	78.8	88.8	-2.8	-10.7	5.1
3	2	79.9	73.4	86.5	80.6	75.7	85.5	-0.7	-8.8	7.5
4	1	61.8	53.1	70.4	61.1	54.4	67.8	0.7	-10.3	11.6
4	2	62.7	54.7	70.7	57.4	51.2	63.6	5.3	-4.8	15.4
5	1	45.0	36.2	53.9	43.2	36.4	50.1	1.8	-9.4	13.0
5	2	49.1	40.8	57.4	43.4	37.1	49.8	5.7	-4.8	16.2
6	1	60.3	51.9	68.7	58.9	52.3	65.6	1.4	-9.3	12.1
6	2	68.5	60.8	76.1	66.3	60.2	72.4	2.2	-7.6	12.0
7	1	40.3	31.8	48.8	42.0	35.3	48.8	-1.7	-12.6	9.2
7	2	74.5	67.2	81.9	69.2	63.2	75.2	5.3	-4.2	14.8
8	1	49.3	41.2	57.3	42.7	36.1	49.3	6.5	-3.9	16.9
8	2	60.0	52.0	67.9	52.4	46.2	58.7	7.6	-2.6	17.7
9	1	39.8	31.3	48.4	45.8	39.0	52.6	-5.9	-16.8	5.0
9	2	44.7	36.6	52.8	42.8	36.6	49.1	1.8	-8.3	12.0
10	1	61.9	54.4	69.4	67.2	60.9	73.6	-5.3	-15.2	4.5
10	2	63.9	56.1	71.7	54.8	48.5	61.1	9.1	-0.9	19.1
11	1	98.4	96.3	100.6	98.5	96.7	100.2	-0.0	-2.8	2.8
11	2	96.7	94.1	99.3	96.9	94.8	99.1	-0.2	-3.7	3.2
12	1	65.0	56.4	73.6	69.3	63.0	75.7	-4.3	-15.0	6.4
12	2	77.8	71.1	84.6	81.9	77.3	86.6	-4.1	-12.3	4.1
13	1	3.3	0.1	6.5	4.0	1.2	6.7	-0.6	-4.9	3.6
13	2	6.7	2.6	10.9	3.0	1.0	5.0	3.7	-0.9	8.3
14	1	2.2	-0.4	4.8	1.6	-0.2	3.4	0.7	-2.5	3.8
14	2	4.5	1.0	8.0	1.7	0.2	3.2	2.8	-1.1	6.7
15	1	35.2	26.8	43.6	25.9	20.1	31.6	9.3	-0.8	19.5
15	2	56.8	48.7	64.9	45.7	39.6	51.8	11.1	0.9	21.2
16	1	5.3	1.4	9.3	4.3	1.5	7.0	1.1	-3.7	5.9
16	2	16.6	10.2	23.0	9.7	6.1	13.3	7.0	-0.4	14.3
17	1	59.6	51.4	67.9	48.5	41.8	55.2	11.1	0.5	21.7
17	2	64.4	56.5	72.4	51.8	45.6	58.0	12.7	2.6	22.7
18	1	24.2	16.8	31.6	26.1	20.2	32.1	-1.9	-11.4	7.6
18	2	45.9	38.1	53.8	38.2	33.0	43.3	7.8	-1.6	17.1
19	1	17.9	11.2	24.5	23.5	16.4	30.7	-5.7	-15.5	4.1
19	2	40.5	32.4	48.6	43.4	37.3	49.5	-2.9	-13.0	7.3
20	1	42.5	34.8	50.2	44.1	38.3	50.0	-1.6	-11.3	8.0
20	2	59.4	51.7	67.0	50.0	44.7	55.4	9.3	0.0	18.7
21	1	37.0	29.2	44.8	43.6	36.7	50.5	-6.6	-17.0	3.8
21	2	54.2	46.1	62.3	55.6	49.5	61.7	-1.4	-11.6	8.7
22	1	62.5	54.0	70.9	65.5	59.0	72.0	-3.0	-13.7	7.7
22	2	69.7	62.3	77.1	60.2	54.0	66.3	9.5	-0.1	19.2

1	23	1	47.5	39.1	55.9	62.5	54.9	70.2	-15.0	-26.4	-3.6
2	23	2	64.9	56.9	73.0	68.0	61.2	74.8	-3.1	-13.7	7.4
3	24	1	75.1	67.4	82.9	62.2	55.5	68.9	12.9	2.7	23.2
4	24	2	67.6	59.7	75.5	63.5	57.4	69.7	4.1	-5.9	14.1
5	25	1	12.1	6.1	18.0	11.1	7.0	15.1	1.0	-6.2	8.2
6	25	2	16.7	10.3	23.1	19.3	14.4	24.2	-2.6	-10.6	5.5
7	26	1	76.2	68.6	83.8	63.5	56.8	70.1	12.7	2.6	22.8
8	26	2	69.4	61.6	77.2	66.9	60.9	73.0	2.4	-7.4	12.3
9	27	1	50.4	41.5	59.3	42.3	35.6	49.1	8.1	-3.1	19.3
10	27	2	57.2	48.9	65.4	52.6	46.3	58.9	4.6	-5.8	15.0
11	28	1	8.6	3.7	13.6	8.2	4.6	11.8	0.5	-5.7	6.6
12	28	2	9.8	5.1	14.5	15.4	11.0	19.8	-5.6	-12.0	0.8
13	29	1	50.4	41.5	59.3	43.3	36.5	50.1	7.1	-4.1	18.3
14	29	2	59.2	51.1	67.4	55.9	49.7	62.2	3.3	-7.0	13.6
15	30	1	32.7	24.5	41.0	28.7	22.6	34.7	4.1	-6.1	14.3
16	30	2	49.2	41.0	57.4	46.9	40.5	53.3	2.3	-8.1	12.7
17	31	1	42.0	33.0	51.1	29.9	23.5	36.3	12.1	1.1	23.2
18	31	2	48.6	40.3	56.8	44.5	38.2	50.8	4.1	-6.3	14.5
19	32	1	3.9	0.4	7.3	5.3	2.2	8.5	-1.5	-6.2	3.2
20	32	2	9.4	4.2	14.5	9.1	5.7	12.6	0.2	-6.0	6.4
21	33	1	43.6	34.5	52.6	32.2	25.8	38.7	11.3	0.2	22.5
22	33	2	52.9	44.8	61.0	48.2	41.9	54.5	4.7	-5.5	15.0
23	34	1	7.0	2.5	11.5	6.2	2.8	9.6	0.8	-4.8	6.4
24	34	2	8.6	3.8	13.4	14.4	9.8	19.0	-5.8	-12.5	0.8
25	35	1	57.2	48.2	66.2	44.5	37.7	51.4	12.7	1.3	24.0
26	35	2	63.2	55.1	71.4	62.5	56.4	68.7	0.7	-9.5	10.9
27	37	1	76.6	69.1	84.1	72.9	66.7	79.1	3.7	-6.0	13.4
28	37	2	81.6	74.9	88.3	78.9	73.5	84.2	2.7	-5.8	11.3
29	38	1	59.0	49.9	68.0	54.7	47.8	61.7	4.2	-7.2	15.6
30	38	2	56.1	47.8	64.4	51.2	44.7	57.6	4.9	-5.6	15.4
31	39	1	0.0	0.0	0.0	5.4	-0.9	11.8	-5.4	-11.8	0.9
32	39	2	5.8	-3.1	14.7	9.1	1.3	16.9	-3.3	-15.1	8.5
33	40	1	0.0	0.0	0.0	1.4	-0.2	3.0	-1.4	-3.0	0.2
34	40	2	1.1	-0.4	2.6	1.8	0.2	3.4	-0.7	-2.9	1.4
35	40	1	17.6	10.9	24.4	10.6	6.5	14.7	7.0	-0.9	14.9
36	40	2	8.0	3.9	12.1	8.5	4.9	12.2	-0.6	-6.0	4.9
37	42	1	52.6	44.3	61.0	43.8	36.9	50.6	8.9	-2.0	19.7
38	42	2	47.5	39.4	55.6	41.8	35.6	48.1	5.7	-4.6	15.9
39	43	1	35.6	27.4	43.8	25.6	19.8	31.5	10.0	-0.1	20.1
40	43	2	35.5	28.0	43.0	26.9	21.5	32.4	8.6	-0.7	17.8
41	44	1	10.1	4.2	16.0	12.2	7.3	17.1	-2.1	-9.8	5.6
42	44	2	20.6	14.1	27.0	21.6	16.3	26.8	-1.0	-9.3	7.3
43	45	1	9.0	-4.7	22.7	7.8	0.4	15.2	1.2	-14.3	16.8
44	45	2	6.6	-9.9	23.2	5.2	0.3	10.1	1.4	-15.8	18.7
45	46	1	63.4	54.6	72.2	56.6	49.9	63.2	6.9	-4.2	17.9
46	46	2	68.9	61.1	76.7	61.7	55.5	67.9	7.2	-2.8	17.2
47	47	1	33.0	24.7	41.4	35.9	29.3	42.5	-2.8	-13.5	7.8
48	47	2	70.4	62.8	77.9	66.6	60.6	72.7	3.7	-6.0	13.4
49	48	1	68.8	60.6	77.0	64.9	58.5	71.4	3.9	-6.5	14.3
50	48	2	76.2	69.3	83.0	70.7	64.9	76.5	5.4	-3.6	14.4
51	49	1	66.5	57.9	75.1	64.5	57.1	71.8	2.1	-9.2	13.3
52	49	2	68.6	59.9	77.3	67.3	60.6	73.9	1.3	-9.7	12.3

1	50	1	51.7	42.1	61.4	48.4	40.3	56.4	3.4	-9.2	15.9
2	50	2	55.2	46.3	64.1	56.1	49.0	63.2	-0.9	-12.3	10.4
3	51	1	33.2	14.6	51.7	78.0	65.9	90.2	-44.9	-67.0	-22.7
4	51	2	40.6	17.5	63.7	63.7	48.8	78.5	-23.0	-50.5	4.5
5	52	1	73.0	65.1	80.9	69.2	62.7	75.6	3.8	-6.3	14.0
6	52	2	83.8	78.2	89.3	82.1	77.1	87.0	1.7	-5.7	9.1

7 a. For text see Table S1

8 b. 1 Aurangabad, 2 Gopalganj

9 c. Estimated with Stata command svy

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table S6: Point estimates and confidence intervals of indicators for a sample of 3-5 month infants and subsample of 3-5 month infants in two districts: Bihar, India, 2015

Indicator ^a	District ^b	3-5 month subsample			3-5 month sample			Difference		
		Point estimate	Confidence interval ^c		Point estimate	Confidence interval ^c		Point estimate	Confidence interval	
			Lower	Upper		Lower	Upper		Lower	Upper
37	1	72.3	62.7	81.8	70.4	64.2	76.5	1.9	13.3	-9.5
37	2	91.3	85.8	96.8	86.2	81.8	90.6	5.1	12.2	-2.0
38	1	49.7	39.3	60.2	45.8	38.9	52.6	4.0	16.4	-8.5
38	2	69.7	61.2	78.2	64.4	58.3	70.5	5.4	15.8	-5.1
52	1	39.7	28.6	50.7	65.3	58.8	71.8	-25.6	-12.8	-38.4
52	2	45.8	35.6	55.9	55.2	49.1	61.2	-9.4	2.4	-21.2

a. For text see Table S1

b. 1 Aurangabad, 2 Gopalganj

c. Estimated with Stata command svy

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 How well do mothers recall their own and their infants' perinatal events? a two-district
4 study using cross-sectional stratified random sampling in Bihar, India
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Joseph J Valadez^{1*}, Baburam Devkota¹, Caroline Jeffery¹, Wilbur C. Hadden²

¹ Department of International Public Health, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

² Department of Sociology, Center for Innovation, University of Maryland, USA

* Corresponding author: Professor Joseph J Valadez, Department of International Public Health, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA United Kingdom. Email: joseph.valadez@lstmed.ac.uk; Mobile: +44(0)754 8523717

Manuscript word length: 2255

Abstract

Objective: Global monitoring of maternal, newborn and child health (MNCH) programmes use self-reported data subject to recall error which may lead to incorrect decisions for improving health services and wasted resources. To minimize this risk, samples of mothers of infants aged 0-2 and 3-5 months are sometimes used. We test whether a single sample of mothers of infants 0-5 months provides the same information.

Design: An annual MNCH household survey in 2-districts of Bihar, India (N=6million).

Participants: Independent samples (n=475 each) of mothers of infants 0-5, 0-2 and 3-5 months.

Outcome measures: Main analyses compare responses from the 0-5 and 0-2 month samples with Mantel-Haenszel-Cochran statistics using 51 indicators in 2-districts.

Results: No measurable differences detected in 79.4% (81/102) comparisons; 20.6% (21/102) display differences for the main comparison. Sub-analyses produce similar results. A difference detected for exclusive breastfeeding is due to premature complementary-feeding by older infants. Measurable differences were detected in 33% (8/24) of the indicators on Front Line Worker (FLW) support, 26.9% (7/26) of indicators of birth preparedness and place of birth and

1 attendant, and 9.5% (4/42) of the indicators on neonatal
2 and antenatal care.

3 **Conclusions:** Differences in FLW visits and compliance with
4 their advice may be due to seasonal effects: mothers of
5 older infants 3-5 months were pregnant during the dry
6 season; mothers of infants 0-2 months were pregnant during
7 the monsoons, making transportation difficult. Useful
8 coverage estimates can be obtained by sampling mothers with
9 infants 0-5 months as with two samples suggesting that
10 mothers of young infants recall their own perinatal events
11 and those of their children. For some indicators (e.g.,
12 exclusive breast feeding) it may be necessary to adjust
13 targets. Excessive stratification wastes resources, does
14 not improve the quality of information, and increases the
15 burden placed on data collectors and communities, which can
16 increase non-sampling error.

17 **INTRODUCTION**

18 ~~Global monitoring of maternal, newborn and child health~~
19 ~~(MNCH) programmes use self-reported data subject to recall~~
20 ~~error which leads to incorrect decisions for improving~~
21 ~~health services and wasted resources. To minimize this~~
22 ~~risk, samples of mothers of infants aged 0-2 and 3-5 months~~
23 ~~are sometimes used. We report a test of whether a single~~

1 ~~sample of mothers of infants 0-5 months provides the same~~
2 ~~information.~~

3 **METHODS**

4 ~~An annual MNCH household survey in two districts of Bihar,~~
5 ~~India (N=6million), collected data from independent samples~~
6 ~~of mothers of infants 0-5, 0-2 and 3-5 months. Analyses~~
7 ~~compare responses from the 0-5 and 0-2 month samples~~
8 ~~(n=950), 0-2 and 3-5 month subsamples (n=475), 0-2 month~~
9 ~~sample and 0-2 month subsample (n=7136759), and 3-5 month~~
10 ~~sample and 3-5 month subsample (n=713666) to assess recall~~
11 ~~bias using 51 indicators with Mantel-Haenszel-Cochran~~
12 ~~statistics.~~

13 **RESULTS**

14 ~~No measurable differences are detected in 79.4% of~~
15 ~~comparisons while 20.6% display differences for the main~~
16 ~~comparison of the 0-5 and 0-2 month samples. The other sub-~~
17 ~~analyses produced similar results. A difference detected~~
18 ~~for exclusive breastfeeding is due to early commencement of~~
19 ~~complementary feeding by older infants. There are~~
20 ~~measurable differences in a third of the indicators on~~
21 ~~front line worker (FLW) support, about a quarter of~~
22 ~~indicators of birth preparedness and place of birth and~~
23 ~~attendant, a sixth or less of indicators on neonatal and~~
24 ~~antenatal care. Differences in FLW visits and compliance~~

1 ~~with their advice may be due to seasonal effects: mothers~~
2 ~~of older infants 3-5 months were pregnant during the dry~~
3 ~~season; mothers of children 0-2 months were pregnant during~~
4 ~~the monsoons, making transportation difficult.~~

5 **CONCLUSIONS**

6 ~~Useful coverage estimates can be obtained by sampling~~
7 ~~mothers with infants 0-5 months as with two cohorts: 0-2~~
8 ~~and 3-5 months suggesting that mothers of young infants~~
9 ~~recall their own perinatal events and those of their~~
10 ~~children, but for some indicators, for example exclusive~~
11 ~~breast feeding, it may be necessary to adjust targets.~~
12 ~~Excessive stratification wastes resources, does not improve~~
13 ~~the quality of information, and increases the burden placed~~
14 ~~on data collectors and communities. This burden can~~
15 ~~increase non-sampling error.~~

16 **STRENGTHS AND LIMITATIONS OF THIS STUDY**

- 17 • Strength: The data were produced using stratified random sampling with no apparent
18 design effect leading to an efficient use of information.
- 19 • Strength: Data were collected from female participants by female data collectors
20 which is likely to have reduced non-sampling error.
- 21 • Strength: The large study population is very large covering a large geographical
22 area, reducing the likelihood that the results are pertinent only to a small group of
23 mothers with infants, and; results may be generalizable.

- 1 • Strength: Both weighted and unweighted results are presented giving strength to the
2 conclusions.

3 Limitation: Due to insufficient overlap of variables in the 0-5 month sample and
4 the 3-5 month sample, The study is confined to assessing data in the 0-2 month
5 sample vis a vis a 0-5 month sample of infants; a comparison between the 3-5
6 month sample vis a vis the 0-5 month sample was not possible. would have been
7 informative; however, insufficient overlap of variables in the 0-5 month sample
8 and 3-5 month sample prevented us for doing so.

9 • Limitation: Non-independence of indicators precludes formal statistical testing
10 of difference between samples Our study compares a 0-2 month subsample with
11 a 3-5 month subsample as an alternative.

12 •

13 **Key Words:**

14 Maternal recall, surveys, maternal health, MNCH, LQAS,
15 Bihar, India

17 **INTRODUCTION**

18 The progress toward United Nations' Sustainable Development
19 Goal (SDG) 3 -is measured with 9 targets, including the
20 Maternal Mortality Ratio (MMR) and the under 5 mortality
21 rate (U5MR)_(1, 2). In India, Bihar is one of the largest
22 (population 110 million) and poorest (53% of households are
23 in the lowest wealth index quintile of India_(3)) states

1 with high child and maternal mortality (U5MR=54,
2 MMR=208) (4) ~~(see (5) for an evaluation of the health care~~
3 ~~system in Bihar)~~, and is a priority for donor support for
4 health systems strengthening (see (5) for an evaluation of
5 the health care system in Bihar).

6 To accelerate progress toward achieving SDG 3, state
7 governments in India pursue programmes of community-based
8 care (see (6, 7) for descriptions and assessments of this
9 approach). Since 2011, the Bihar Ministry of Health has
10 supported an Integrated Family Health Initiative to improve
11 the availability, quality and use of prenatal, perinatal
12 and postnatal care for mothers and infants (8).

13 The usual way to monitor progress toward achieving these
14 goals is with household surveys. Perhaps the most commonly
15 used surveys are cluster sample surveys such as the
16 Demographic and Health Surveys and the Multiple Indicator
17 Cluster Surveys (9, 10). An alternative design is Lot
18 Quality Assurance Sampling (LQAS), which provides
19 comparable data but is decentralized to local health
20 services organisations and more useful for management and
21 programme planning (11). Several states in India find it
22 benefits their programs (12). Surveys rely on the reports
23 of mothers of infants and young children, but these reports
24 are subject to several sources of potential error and bias

1 through interviewees not knowing, forgetting, and having
2 memory errors_(13, 14). Studies have shown both that
3 mothers can accurately report significant facts about the
4 birth and care of their children many years after the event
5 (15), but also that even immediately after giving birth
6 mothers may misreport details_(16-18). Studies of maternal
7 mothers recall of ~~her~~-their children's vaccination status
8 concluded that due to offsetting errors of maternal
9 reports, the resulting data accurately measured vaccination
10 rates_(19); the pattern of error revealed that mothers
11 whose children are up-to-date or nearly so tended to
12 underestimate their child's vaccination status whilst
13 mothers whose children have few vaccinations, overestimate
14 their coverage.

15 To improve the validity of collected data, knowledge,
16 practice and coverage surveys have used samples of mothers
17 of infants 0-11 months of age or 0-5 months of age and
18 children 6-11 months of age. In Bihar, local organisations
19 departed from this convention of sampling among these three
20 cohorts of children under one year of age and have been
21 monitoring their programs' progress by sampling five
22 dedicated cohorts: mothers of children 0-2, 3-5, 6-8, 9-11
23 and 12-23 months old with indicators focused on antenatal
24 care, safe delivery practices, infant and young child

1 feeding practices, immunisation, treatment seeking, and
2 more. To avoid the possibility of maternal recall error,
3 each of the five cohorts was asked questions particularly
4 relevant to a child's specific age group.

5 In countries such as India with high maternal and
6 child mortality rates, regular monitoring of related health
7 service coverage is critical to reducing these rates.
8 However, ~~the~~ survey designs should be affordable and
9 sustainable for local health systems; they should also
10 produce precise, unbiased estimates₂₀. In this study, we
11 explore whether information is gained by sampling cohorts
12 of children aged 0-2 and 3-5 months or whether ~~the~~ sample
13 sizes_s can be reduced by 50% by creating one sample cohort
14 aged 0-5 months.

15 The research question we address is: "Do the health service
16 delivery coverage estimates from a sample of mothers of
17 infants aged 0-5 months differ from those obtained from a
18 sample of mothers of infants ages 0-2 months?" A corollary
19 to this question is: "Do mothers of infants 3-5 months of
20 age display more recall error relative to mothers of
21 infants 0-2 months of age for antenatal, delivery or young
22 infant health practices?" We compare district coverage
23 estimates obtained from two independent samples of infants
24 0-2 months and 0-5 months of age. The implications of this

1 study are important for health systems researchers needing
2 results to appraise and improve their programmes.

3 **METHODS**

4 To answer this question, we collected information from a
5 sample of mothers with infants 0-5 months old and a sample
6 of mothers with infants 0-2 month old ~~infants~~ in two
7 districts. This study took place within the context of a
8 larger survey that also sampled children 3-5, 6-8, 9-11 and
9 12-23 months of age. These four latter samples used
10 questionnaires with variables that either did not overlap
11 at all or overlapped on very few indicators with the
12 questionnaires used to interview the 0-5 month and 0-2
13 month samples. Due to this constraint, in this study, we
14 only use the two aforementioned groups- to assess the
15 measurement of the indicators and refer only to them for
16 the remainder of this paper. The household sampling design
17 we used is a stratified random sample_(21). Within each
18 district, the strata are administrative units of the health
19 system, which in Bihar is called a *block*. Within each block
20 the primary sampling unit is the Anganwadi Centre
21 (Community Health Subcentre) Catchment Area (ACCA); 19 ACCA
22 are selected from each block with probability proportional to size. From each ACCA
23 one respondent is randomly selected from each age-group

1 under study using segmentation sampling_(22, 23). The sample
2 size 19 for each block is chosen to maximize the probability of correctly classifying a
3 block with reference to performance targets on health related indicators (95% reliability)
4 while balancing the probability (10% margin of error) of incorrectly classifying a
5 block and thereby failing to recognize either the accomplishments of local health care
6 delivery systems or the local population's health care needs₍₂₂₎. For this purpose,
7 principles of Lot Quality Assurance Sampling were used along with established
8 probability tables (24-26).

9 There are 14 and 11 blocks in Gopalganj and Aurangabad (N=6
10 million), the two districts selected for this study,
11 respectively. The total sample sizes are: (a) Gopalganj: 19
12 x 14 blocks= 266 infants 0-2 months and 266 infants 0-5
13 months, and (b) Aurangabad: 19 x 11 blocks= 209 infants 0-2
14 months and 209 infants 0-5 months. The 0-5 month old sample
15 is 60% 0-2 months old and 40% 3-5 months old.

16 Using summary data from each of the two samples we analyse
17 the data with Cochran-Mantel-Haenszel (CMH)₍₂₇₎ tests for
18 51 dichotomous indicators (See Supplementary File Table S1)
19 common to the two samples. The CMH tests theoretically have
20 a Chi-square probability distribution with 1 degree of
21 freedom. With a sufficient number of respondents or a
22 sufficient number of blocks, the CMH test is equivalent to
23 [a conditional logistic regression](#)_(28: 114-115). In this
24 analysis both the number of respondents and the number of

1 blocks only approach sufficiency. Consequently, the
2 calculated chi-squares and probabilities must be considered
3 as approximations of their true values.

4 We calculate both unweighted and weighted estimates. The
5 unweighted estimates permit the results from smaller blocks
6 to have equal weight vis à vis larger ones. Since the
7 research question concerns an analysis of which age cohort
8 is most informative, the weighted estimates may not be as
9 useful as the unweighted ones. However, the weighted
10 estimates provide better point estimates of the indicators
11 at the district level. The effect of the weights on the
12 Chi-square statistics is to increase the contribution of
13 the larger blocks and decrease the contribution of the
14 smaller blocks. Hence, we report both sets of results (See
15 Supplementary File Tables S2-S3).

16 The Chi-square probability distribution puts the
17 differences between the districts on a probability scale
18 (See Supplementary File Table S2). To determine meaningful
19 differences in responses between the two age cohorts we
20 used a probability of 0.05 as a cut-off value and consider
21 differences with probabilities less than 0.05 to be
22 possibly meaningful and those with larger probabilities to
23 be likely due to sampling errors. With 102 comparisons (51
24 indicators weighted or unweighted) we must expect some to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 exceed this cut-off by chance alone. If all of the
2 comparisons were independent, we might randomly find 5
3 differences, but many of the indicators measure related
4 events (e.g., number of ANC visits and tetanus toxoid
5 vaccinations) and the weighted and unweighted estimates are
6 similar, so these indicators are not all independent and it
7 is not possible to calculate an expected number of
8 differences nor is it appropriate to interpret these
9 probabilities as measures of "statistical significance".

10 **Patient and Public Involvement**

11 This study does not involve patients. Also, the public was
12 not involved in the design, conduct and reporting of the
13 research. The public was engaged as interviewees. To
14 ensure local engagement we coordinated with the Bihar
15 Ministry of Health, local implementing non-governmental
16 organizations and our donor. We also shared the results
17 with them and offered further dissemination of results.

18 **RESULTS**

19 We find a high level of agreement between the two samples
20 (Table 1). Out of 102 weighted and unweighted comparisons
21 between the estimates from the 0-2 month and 0-5 month
22 samples there is no probable difference in 81 (79.4%) in
23 both the unweighted and weighted estimates. We detect that

1 probable differences exist for 13 comparisons (12.7%). For
 2 the remaining eight comparisons the weighted and unweighted
 3 estimates disagree. The weighted estimates find seven
 4 differences that the unweighted estimates do not; the
 5 unweighted estimates find one difference that the weighted
 6 estimates do not find.

Table 1. Number of indicators by probability of a difference between the 0-2 months and the 0-5 months samples for weighted and unweighted samples

Unweighted	Weighted					
	Aurangabad		Gopalganj		Total	
	\geq .05	$<$.05	\geq .05	$<$.05	\geq .05	$<$.05
\geq .05	40	3	41	4	81	7
$<$.05	0	8	1	5	1	13

8
 9
 10 For different health service domains, the number of
 11 indicator comparisons varies from two (Exclusive
 12 Breastfeeding-EBF) to 24 concerning home visits by Front
 13 Line Worker (FLW) support (Table 2). The two principal FLW
 14 are Anganwadi workers and Accredited Social Health
 15 Activists (ASHA).

Table 2. Number of indicator comparisons by subject domain showing a measurable difference using weighted and unweighted estimates of 0-2 months and 0-5 months samples

Health Service Domain	Total comparisons	No measurable difference between 0-2 and 0-5 months results	Measurable difference between 0-2 and 0-5 months results			
			Both	Unweighted only	Weighted only	Percent (%) indicators with different results
Antenatal care	22	21	0	0	1	5
Place of birth & attendant	8	6	1	0	1	25
Birth preparedness	18	13	3	0	2	28
Front Line Worker support	24	16	6	0	2	33
Maternal health	8	8	0	0	0	0
Neonatal care	20	17	1	1	1	15
Exclusive Breastfeeding	2	0	2	0	0	100
Totals	102	81	13	1	7	21.9

1
2
3 1
4
5 2 In the FLW support domain 33% of comparisons have probable
6
7 3 differences. The neonatal health domain has 20 comparisons
8
9 4 and the birth preparedness domain has 18; in these domains
10
11 5 15% and 28% show probable differences, respectively. Place
12
13
14 6 of birth and attendant, and maternal health each have eight
15
16 7 comparisons with 25%, or two comparisons, and 0
17
18 8 comparisons, respectively, showing a possible difference.
19
20 9 The differences between the two samples cluster around home
21
22 10 visits from FLW and behaviours associated with birth
23
24 11 preparedness and neonatal care. Details of these
25
26 12 differences are listed in Table 3.
27
28 13 For two indicators, both the weighted and unweighted
29
30 14 estimates display probable differences between the 0-2 and
31
32 15 0-5 months samples in both districts. For indicator #52,
33
34 16 the proportion exclusively breastfeeding, the 0-5 months
35
36 17 cohort has the lower estimate, and indicator #24, the
37
38 18 proportion of mothers visited by an ASHA at least once
39
40 19 during their last pregnancy, the 0-5 months sample gives
41
42 20 the higher estimate, about 74%, compared to 63% in the 0-2
43
44 21 months sample (See Supplementary Tables S2-3).
45
46 22
47
48 23 Additional analyses comparing subsamples of mothers of
49
50 24 infants 0-2 months and 3-5 months from the 0-5 months
51
52
53
54
55
56
57
58
59
60

1
2
3 1 sample, the sample of mothers of infants 0-2 months and the
4
5 2 subsample of infants 0-2 month, and the sample of mothers
6
7 3 of infants 3-5 months and the subsample of infants 3-5
8
9 4 months produced similar results (See Supplementary Text and
10
11 5 Tables S4a-b, Tables S5a-b and Table S6).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Indicators by health service domain showing measurement differences between 0-2 months and 0-5 months samples

Health Service Domain and Indicator	Indicator No.	District	Weighted Coverage (%)		p-value		Estimate type
			0-2 months	0-5 months	Unweighted Estimate	Weighted Estimate	
Antenatal care							
Proportion of mothers of infants (0-2/0-5 months) registered during their last pregnancy	1	Aurangabad	85.2	77.6	0.0552	0.0365	Weighted
Place of birth & attendant							
Proportion of mothers of infants (0-2/0-5 months) whose last child was delivered at a public facility	38	Gopalganj	51.2	61.6	0.0363	0.0159	Both
Proportion of mothers of infants (0-2/0-5 months) whose last child was delivered at a health facility (private or public facility)	37	Gopalganj	78.9	85.5	0.0643	0.0459	Weighted
Birth preparedness							
Proportion of mothers of infants (0-2/0-5 months) who planned transportation to health facility in their last pregnancy (home & institutional delivery)	15	Gopalganj	45.7	56.0	0.0266	0.0158	Both
Proportion of mothers of infants (0-2/0-5 months) who identified persons to care for the baby immediately after birth (home + institutional delivery)	17	Gopalganj	51.8	62.6	0.0255	0.0103	Both
Proportion of mothers of infants (0-2/0-5 months) who planned for institutional delivery and	23	Aurangabad	62.5	47.0	0.0039	0.0052	Both

1	identified person to accompany her during the delivery							
2								
3	Proportion of mothers who planned for institutional delivery of infants (0-2/0-5 months) who had a new blade & thread for their delivery	19	Aurangabad	23.5	14.5	0.062	0.0429	Weighted
4								
5								
6								
7								
8	Proportion of mothers who planned institutional delivery of infants (0-2/0-5 months) who arranged clean cloth for mothers and baby	21	Aurangabad	43.6	31.2	0.0546	0.0137	Weighted
9								
10								
11								
12								
13								
14	Front Line Worker support							
15	Proportion of mothers of infants (0-2/0-5 months) who were visited by ASHA at least once during their last pregnancy	24	Aurangabad	62.2	75.2	0.0023	0.0042	Both
16			Gopalganj	63.5	73.0	0.0284	0.0175	Both
17								
18								
19	Proportion of mothers of infants (0-2/0-5 months) visited at home by FLWs at least once during their last pregnancy	26	Aurangabad	63.5	76.7	0.0021	0.0032	Both
20								
21								
22								
23	Proportion of mothers of infants (0-2/0-5 months) visited at home by ASHA within 24 hours of last delivery	31	Aurangabad	29.9	44.9	0.0009	0.0016	Both
24								
25								
26								
27								
28	Proportion of mothers of infants (0-2/0-5 months) visited at home by any FLW within 24 hours of last delivery	33	Aurangabad	32.2	46.7	0.0015	0.0026	Both
29								
30								
31								
32	Proportion of mothers of infants (0-2/0-5 months) visited at home by any FLW within first week of last delivery	35	Aurangabad	44.5	59.3	0.0018	0.0026	Both
33								
34								
35								
36	Proportion of mothers of infants (0-2/0-5 months) visited at home by any AWW within the first week of the last delivery	34	Gopalganj	14.4	8.9	0.0959	0.0471	Weighted
37								
38								
39								
40	Proportion of mothers of infants (0-2/0-5 months) visited by	27	Gopalganj	52.6	61.1	0.0617	0.0449	Weighted
41								
42								
43								
44								
45								
46								
47								

1	ASHAs at least once during their last trimester of pregnancy							
2	Infant care							
3	Proportion of infants aged (0-2/0-5 months) who were delivered at home continued with dry cord care	51	Aurangabad	78.0	45.4	0.0001	0.0006	Both
4			Gopalganj	63.7	41.1	0.0431	0.0627	Unweighted
5	Proportion of infants aged (0-2/0-5 months) weighed after birth (Public facility/Private facility/Home)	48	Gopalganj	70.7	78.2	0.0727	0.0464	Weighted
6								
7	Exclusive Breastfeeding							
8	Proportion of infants (0-2/0-5 months) breast-fed in the past 24 hours (Exclusively Breast-fed)	52	Aurangabad	69.2	59.7	0.0229	0.0411	Both
9			Gopalganj	82.1	68.4	0.0001	0.0003	Both

1
2
3 1
4
56 2 **DISCUSSION**
78
9 3 **Statement of principal findings**
10

11 4 There are no measurable differences in coverage estimates
12
13 5 for 79.4% (81 comparisons) of the indicator comparisons
14
15 6 between the samples of mothers with infants 0-2 months old
16
17 7 versus mothers of infants 0-5 months old; 12.7% (13
18
19 8 comparisons) display measurable differences. The remaining
20
21 9 7.8% (eight comparisons) display discrepancies between the
22
23
24
25 10 weighted and unweighted estimates.
26
27

28 11 **Strengths and weaknesses of the study**
29

30 12 The strengths of this study are that it compares estimates
31
32 13 from two independent samples and that there are many
33
34 14 estimates from diverse domains. The weaknesses of this
35
36 15 study are that the data were collected in only 2 districts
37
38 16 of 1 state in India and in different months of a single
39
40 17 year, and that indicators from the sample of mothers of -3-
41
42 18 5 month old infants comparable to those of the 0-2 month
43
44 19 old infants, using the same questionnaire, were not
45
46 20 collected. Supplemental analyses comparing 0-2 and 3-5
47
48 21 month subsamples of the 0-5 sample did not uncover evidence
49
50 22 of bias due to the combination of these two age groups.
51
52
53
54
55
56
57
58
59
60

1 **Strengths and weaknesses in relation to other studies**

2 Other studies of maternal recall bias have sought a "gold
3 standard" to represent reality and to evaluate measures.
4 Our study, of course, is interested in reality, but this
5 study compares alternative measures needed to assess the
6 Bihar health program. It also uses a complete sample of
7 the age grouping under study rather than just a sub-sample
8 of a larger age grouping. A weakness of this approach is
9 that the analysis does not result in a formal statistical
10 test; our conclusion is based on the weight of the
11 evidence.

12 **Meaning of the study**

13 The evidence indicates that samples of the broader group
14 yield comparable results to those of the narrower age
15 group. It is not necessary to double the total sample by
16 measuring independently 0-2 months and 3-5 months cohorts
17 of children. These results also tend to dispel the
18 hypothesis that maternal recall is problematic for mothers
19 during the first 6 months following delivery. Our results
20 are more consistent with conclusions presented in earlier
21 research_(15) and they support those organizations
22 collecting data with 0-5 month cohorts.

1 Indicator #52, EBF, displayed two comparisons measuring
2 decreases in both districts. This is not surprising as
3 fewer infants are expected to be exclusively breast fed in
4 a sample ranging from 0 to 5 months than a sample ranging 0
5 to 2 months; mothers introduce complementary feeding and
6 liquids as infants age despite this being a health risk.
7 This difference could be accommodated by adjusting
8 expectations and targets for the indicator.

9 **Unanswered questions and future research**

10 Further investigation and consideration of the differences
11 is warranted. The eight differences found in the FLW
12 support indicators deserve more scrutiny. Seven show higher
13 estimates for the 0-5 cohort and one has a higher estimate
14 for the 0-2 month cohort. – The former seven differences
15 may be due to excessive rainfall during the July-September
16 (monthly average 318.95mm, range:195.99-395.8mm) versus the
17 lesser rainfall during June-April (monthly average 25.52mm,
18 range: 0.3-27.88mm) which in the last trimester may have
19 reduced the access of ASHA in the 0-2 month cohort (29).
20 Indicators such as these may be particularly sensitive to
21 rainfall and may explain why more mothers in the 0-5 month
22 cohort displayed higher FLW visitation estimates since FLW

1 were not impeded by the monsoon and the resulting muddy
2 roads.
3 Differences in birth preparedness and institutional birth
4 may be a consequence of ~~the~~ differences in rainfall or in
5 FLW support; the results signal a with increases a response
6 to the needs for more careful planning when transportation
7 is difficult and decreases due to a reductions in the
8 educational effectiveness efforts of ~~the~~ FLW by reducing
9 their access to women. Or, some of these differences may
10 just be due to noise in the data.

12 CONCLUSIONS

13 Overall, the answer to the research question, "Can one get
14 the same district coverage estimates from a sample of
15 mothers of infants aged 0-5 months as from a sample of
16 mothers of infants ages 0-2 months?" is yes. This result
17 can be paraphrased as: mothers do not display increased
18 recall errors of their perinatal health care behaviour in a
19 cohort of mothers with infants 0-5 months as compared with
20 mothers with younger infants. Substantial resources and
21 effort can be saved using a survey design that avoids
22 needless expenses to collect data that provides
23 insubstantial amounts of information. It also reduces the
24 burden on data collectors and community participants.

1
2
3 1 Fatigue to both groups can result in needless non-sampling
4
5 2 error.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGMENTS

We gratefully acknowledge the essential roles of Hemant Das, Alok Prahdan and Sanjay Biswa for their careful field work supporting the LQAS survey and the HMIS data retrieval. We thank Prof Imelda Bates, Prof Brian Faragher and Nancy Vollmer for their valuable feedback on an earlier version of this manuscript. This work was supported by the Bill and Melinda Gates Foundation (Investment ID OPP1142889).

ETHICS

[The Ethical Committees of the Indian Institute of Public Health \(No IIPHB-IEC-2016/010\) and the Liverpool School of Tropical Medicine Research Ethics Committee approved the protocol, study instruments and consent procedures for the data collection of the household surveys \(Research Protocol 16-023\).](#)

~~The Ethical Committees of the authors' institutes approved the protocol, study instruments and consent procedures for the survey data collection.~~

COMPETING INTERESTS

The authors have read the statement on competing interests and declare that they have none.

AUTHORS' CONTRIBUTIONS

JJV, BD developed the research question and survey design; WH, CJ carried out the statistical analyses; JJV obtained

1
2
3 the funding and donor support for the research; BD trained
4 and managed the survey teams in Bihar; JJV, WH, CJ
5 interpreted the data; CJ responsible for data curation; all
6 authors wrote and reviewed the paper.
7

8 9 10 **COMPETING INTERESTS**

11 ~~The authors have no competing interests.~~
12

13 14 **FUNDING**

15 This research was funded by the Bill and Melinda Gates Foundation Investment ID
16 OPP1142889
17

18 19 20 **DATA SHARING STATEMENT**

21 Data are available upon reasonable request.
22
23

24 25 **REFERENCES**

- 26 1. UNSTATS. Official list of MDG indicators: United Nations Statistics Division,
27 Department of Economic and Social Affairs; 2005 [Available from:
28 <https://unstats.un.org/unsd/mdg/Host.aspx?Content=Indicators/OfficialList.htm>.
29
- 30 2. UN DESA. Sustainable Development Goals: Division for Sustainable
31 Development Goals, United Nations Department of Economic and Social Affairs; 2015
32 [Available from: [https://www.un.org/sustainabledevelopment/sustainable-development-
33 goals/](https://www.un.org/sustainabledevelopment/sustainable-development-goals/).
34
- 35 3. International Institute for Population Sciences (IIPS) and ICF. National Family
36 Health Survey (NFHS-4), 2015-16: India. Mumbai: IIPS; 2017.
- 37 4. MOSPI. Millennium Development Goals India Country Report 2015. New Delhi:
38 Social Statistics Division, Central Statistics Office, Ministry of Statistics and Programme
39 Implementation; 2015.
- 40 5. Karvande S, Sonawane D, Chavan S, Mistry N. What does quality of care mean
41 for maternal health providers from two vulnerable states of India? Case study of Bihar
42 and Jharkhand. *Journal of Health, Population and Nutrition*. 2016;35(6):1-10.
43
- 44 6. Mohan P, Kishore B, Singh S, Bahl R, Puri A, Kumar R. Assessment of
45 implementation of integrated management of neonatal and childhood illness in India.
46 *Journal of Health, Population and Nutrition*. 2011;29(6):639-38.
- 47 7. Neogi SB, Sharma J, Chauhan M, Khanna R, Chokshi M, Srivastava, et al. Care
48 of newborn in the community and at home. *Journal of Perinatology*. 2016;36(Suppl
49 3):s13-7.
50
- 51 8. CARE Family Health. Integrated Family Health Initiative: Catalysing change for
52 healthy communities. 2013.
- 53 9. Blanc AK, Warren CE, McCarthy KJ, Kimani J, Ndwiga C, RamaRao S.
54 Assessing the validity of indicators of the quality of maternal and newborn health care in
55 Kenya. *Journal of Global Health*. 2016;6(1):1-11.
56
57
58
59

10. Eisele TP, Rhoda DA, Cutts FT, Keating J, Ren R, Barros AJD, et al. Measuring coverage in MNCH: Total survey error and the interpretation of intervention coverage estimates from household surveys. *PLoS Medicine*. 2013;10(5):e1001386.
11. Anoke SC, Mwai P, Jeffery C, Valadez JJ, Pagano M. Comparing two survey methods of measuring health-related indicators: Lot Quality Assurance Sampling and Demographic Health Surveys. *Tropical Medicine and International Health*. 2015;20(12):1756-70.
12. Valadez JJ, Devkota B, Pradhan MM, Meherda P, Sonal GS, Dhariwal A, et al. Improving malaria treatment and prevention in India by aiding district managers to manage their programmes with local information: a trial assessing the impact of Lot Quality Assurance Sampling on programme outcomes. *Trop Med Int Health*. 2014.
13. Tourangeau R. Remembering what happened: Memory errors and survey reports. In: Stone AA, Jobe JB, Bachrach CA, Cain VS, Kurtzman HS, editors. *The Science of Self-report : Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum; 2000.
14. Bradburn NM. Temporal representation and event dating. In: Stone AA, Jobe JB, Bachrach CA, Cain VS, Kurtzman HS, editors. *The Science of Self-report : Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum; 2000.
15. Tomeo CA, Rich-Edwards JW, Michels KB, Berkey CS, Hunter DJ, Frazier AL, et al. Reproducibility and validity of maternal recall of pregnancy-related events. *Epidemiology*. 1999;10(6):774-7.
16. Elkadry E, Kenton K, White P, Creech S, Brubaker L. Do mothers remember key events during labor? *American Journal of Obstetrics and Gynecology*. 2003;189(1):195-200.
17. Li R, Scanlon KS, Serdula MK. The validity and reliability of maternal recall of breastfeeding practice. *Nutrition Reviews*. 2005;63(4):103-10.
18. Miles M, Ryman TK, Dietz V, Zell E, Luman ET. Validity of vaccination cards and parental recall to estimate vaccination coverage: A systematic review of the literature. *Vaccine*. 2013;31:1560-8.
19. Valadez JJ, Weld LH. Maternal recall error of child vaccination status in a developing nation. *Am J Public Health*. 1992;82(1):120-2.
20. Chan M, Kazatchkine M, Lob-Levyt J, Obaid T, Schweizer J, Sidibe M, et al. Meeting the demand for results and accountability: a call for action on health data from eight global health agencies. *PLoS Med*. 2010;7(1):e1000223.
21. Valadez JJ, Weiss W, Leburg C, Davis R. *Assessing Community Health Programs: A Trainer's Guide. Using LQAS for Baseline Surveys and Regular Monitoring*. 2nd ed. St Albans: Teaching-aids At Low Cost; 2007.
22. Turner AG, Magnani RJ, Shuaib M. A not quite as quick but much cleaner alternative to the Expanded Programme on Immunization (EPI) Cluster Survey design. *International Journal of Epidemiology*. 1996;25(4):198-203.
23. Davis RH, Valadez JJ. Improving the collection of knowledge, attitude and practice data with community surveys: a comparison of two second-stage sampling methods. *Health Policy Planning*. 2014;29(8):1054-60.
24. Dodge HF, Romig HG. *Sampling inspection tables: single and double sampling*. 2nd ed. New York: John Wiley & Sons; 1959.

- 1
2
3 25. Robertson SE, Joseph J V. Global review of health care surveys using lot quality
4 assurance sampling (LQAS), 1984–2004. *Social Science & Medicine*. 2006;63:1648-60.
5 26. Valadez JJ. Assessing child survival programmes in developing countries: testing
6 lot quality assurance sampling. Boston: Harvard University Press; 1991.
7 27. Mantel N. Chi-square tests with one degree of freedom: extensions of the Mantel-
8 Haenszel procedure. *J Am Stat Assoc*. 1963;58(303):690-700.
9 28. Agresti A. *An Introduction to Categorical Data Analysis*. 2nd ed. Hoboken, New
10 Jersey: Wiley-Interscience; John Wiley & Sons; 2007.
11 29. World Weather Online. Patna Monthly Climate Averages, Bihar, India 2018.8
12 [Available from: [https://www.worldweatheronline.com/patna-weather-](https://www.worldweatheronline.com/patna-weather-averages/bihar/in.aspx)
13 [averages/bihar/in.aspx](https://www.worldweatheronline.com/patna-weather-averages/bihar/in.aspx).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60