



OPEN ACCESS

Association between socioeconomic status and incident stroke in China

Weiju Zhou ¹, Ruoling Chen,¹ Alex Hopkins,¹ Yulong Wang,² Jie Tang,³ Xiangyan Chen,⁴ Angela Clifford,¹ Yuesong Pan,^{5,6} Ken Forthby,⁷ Jindong Ni,⁸ Duolao Wang,⁹ Eric Brunner ¹⁰

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jech-2019-213515>).

For numbered affiliations see end of article.

Correspondence to

Professor Ruoling Chen, Faculty of Education, Health and Wellbeing, University of Wolverhampton, Wolverhampton WV1 1LY, UK; R.Chen@wlv.ac.uk

Received 18 November 2019

Revised 31 January 2020

Accepted 3 February 2020

ABSTRACT

Background Little is known about the impact of socioeconomic status (SES) on incidence of stroke in China. This study aimed to examine the association of SES, which was measured by different indicators, with incidence of stroke and gender differences in the association.

Methods and results Two prospective cohort studies were conducted including 2852 participants aged ≥ 60 years in Anhui province and 3016 participants in four other provinces in China. During a median follow-up of 7.1 years, 211 incident stroke cases occurred in the Anhui cohort. The risk of stroke increased with living in rural areas (adjusted HR 2.49, 95% CI 1.19 to 5.22; women 3.64, 95% CI 1.17 to 11.32, men 2.23, 95% CI 0.81 to 6.19), but not significantly with educational level, occupational class, satisfactory income and financial problems (except for women with low education). The four-province cohort had 113 incident stroke cases over the 3.1 years' follow-up. The five SES indicators were not significantly associated with incident stroke (except for increased risk in men with high occupation), but additional measurement for actual income showed that incident stroke increased in women with low personal income and in men with high family income. Pooled data from the two cohorts demonstrated the impacts of rural living (1.66, 95% CI 1.08 to 2.57) and having high occupational class (1.56, 95% CI 1.01 to 2.38), and gender differences for women with low education (2.26, 95% CI 1.19 to 4.27).

Conclusions Rural living and being female with low SES are associated with increased stroke risk in China. Strategies to improve public health in the rural communities and gender-specific targets for health inequality should be an integral component of stroke interventions.

in LMICs. There are fewer studies undertaken in LMICs to assess the association of SES with incident stroke.

China is the largest LMIC and has the highest estimated lifetime risk of stroke worldwide.⁶ There are around 5.5 million new stroke cases occurring in China each year.⁵ Over the past decades, China has experienced rapid economic growth, along with an increasing gap in income between rich and poor.⁷ However, it is not known whether socioeconomic inequalities, including urban-rural differences,⁷ are associated with increased risk of stroke.

Previous studies in HICs showed some inconsistent findings of the association between SES and incident stroke.^{3,8} No study has assessed the impacts of different SES indicators on incidence of stroke simultaneously. Most of the studies did not adjust for other SES variables when examining incidence of stroke in relation to one measurement of SES.^{3,4,9} Few studies have investigated gender differences in the association of SES with incident stroke. In this paper we examine data from two community-based cohort studies in China to assess the impacts of multiple measurements of SES on the incidence of stroke and their gender differences.

METHODS

Studied populations were derived from the Anhui cohort study and the four-province cohort study in China.

Anhui cohort study

The methods of the baseline investigation and the follow-up in the Anhui cohort study have been fully described previously.¹⁰ Briefly, we randomly selected 1810 people aged ≥ 65 years who had lived for at least 5 years in Yiming subdistrict of Hefei city in 2001 and 1709 aged ≥ 60 years from all 16 villages in Tangdian district of Yingshang county in 2003. A total of 3336 older people participated in the study (urban $n=1736$), with a response rate of 94.8%. Informed consent was obtained from each participant. In the case of those who were unable to provide informed consent such as due to disability or limited education level, their next of kin or caregivers were invited to provide assent for participation.¹¹ The participants were interviewed at home by a trained survey team from the Anhui Medical University.

The main interview materials were a general health and risk factors questionnaire, and the Geriatric Mental Status (GMS)—a comprehensive semistructured mental state interview.¹² In the general health

INTRODUCTION

The burden of stroke has increased across the world, with the disability-adjusted life-years ranked at third up from fifth over the past 30 years.¹ Previous studies²⁻⁴ showed that low socioeconomic status (SES) was associated with increased incidence of stroke in the general population. The global burden of disease study⁵ demonstrated that people in low and middle-income countries (LMIC) have a higher risk of stroke than in high-income countries (HIC). Current knowledge of the impact of SES on incident stroke has been predominantly derived from HICs, which may not be applicable to those



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Zhou W, Chen R, Hopkins A, et al. *J Epidemiol Community Health* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jech-2019-213515

and risk factors record, we collected data relating to sociodemography, social networks and support, psychosocial aspects, cardiovascular disease (CVD) and other disease risk factors.^{10 12} We documented stroke from participants' self-report of having being diagnosed by a doctor.¹³ In older people self-reported doctor's diagnosis of stroke has been validated¹⁴ and our previous study showed that these older people with stroke had an HR of 1.91 (95% CI 1.45 to 2.52) for 5-year mortality compared with their non-stroke counterparts.¹³ Using the GMS interview data and the Automated Geriatric Examination for Computer Assisted Taxonomy¹² we diagnosed depression and dementia for each participant. According to standard procedures¹⁰ we measured systolic and diastolic blood pressure, weight and height, and waist circumferences for all participants at baseline interview (wave 1).

Baseline SES measurements

We measured SES from long-term residential location in urban and rural areas and from individual records of educational attainment level, occupational class, satisfactory income and a serious financial problem in the past 2 years.¹³ In the interview we asked each participant whether he or she attended any school or not. If answered with 'yes', they would be asked to specify either (1) primary school, (2) secondary school, (3) high secondary school/professional school or (4) university/college. Those without any formal school attainment were defined as illiterate. All participants gave their occupational class according to their current or last main job titles as a peasant, manual worker, officer/teacher, businessman, housewife or other. We asked each participant the questions 'Are you satisfied with your income?' (with answers either at 'very satisfied', 'satisfied', 'average' or 'poor') and 'Have you had any financial problems in the past two years' (with answers 'yes' or 'no').

Follow-up of cohort

One year after the baseline survey we reinterviewed 2608 participants (wave 2) using the same protocol as at baseline. From 2007 to 2009 we successfully re-examined 1757 cohort members (wave 3), and in 2010–2011 we carried out a wave 4 survey on surviving cohort members and reinterviewed 944 participants.¹⁰ In waves 3 and 4, we added in the 10/66 dementia algorithm research package and more risk factors including dietary intakes, passive smoking and personal and family incomes.¹¹ At each wave interview of the follow-up we documented incident stroke based on self-reported doctor diagnosis of stroke.

The vital status of the cohort members was monitored until December 2011. At each survey wave we conducted home visits to obtain information about participants' survival status through multiple sources including resident committees, family members, neighbours and friends. In total we identified 671 deaths during the follow-up of the cohort. Using a standard verbal autopsy questionnaire we interviewed the next of kin responsible for the deceased or reviewed the death certificate to ascertain causes of death,¹² including stroke diagnosis.^{10 13}

Four-province study

Using the same protocol as that in the Anhui study wave 3, we carried out a large-scale community-based household survey in four provinces of China (Guangdong, Heilongjiang, Shanghai and Shanxi) in 2008. Their methods of the baseline investigation have been fully described previously.¹¹ In brief, we randomly recruited no fewer than 500 residents aged ≥ 60 years from the urban and rural communities separately from each province. In total, 4314 participants completed the interview in 2008–2009, with an overall

response rate of 93.8%. We documented cardiovascular diseases and risk factors (CVDRF) and SES variables for each participant.¹¹

In 2010–2012 we followed up the cohort, having identified 259 deaths and reinterviewed 2892 survivals using the same questionnaire as that at baseline.

Statistical analysis

In the Anhui cohort study, we analysed the data from 2852 cohort members, after excluding 141 participants with stroke at baseline and 343 who were lost to follow-up. In the four-province cohort study, the data from 3016 cohort members were analysed, after excluding 166 participants who had stroke at baseline, 1102 who were lost to follow-up and 30 who had no SES indicator measured. We computed person-years at risk for cohort members to the end of follow-up, date of incident stroke, death or loss at follow-up. Annual incidence rate was expressed as cases per 1000 person-years. We described baseline characteristics and CVDRFs of participants using mean (SD) and percentage (%) and examined differences in their distributions between women and men at baseline using a one-way analysis of variance for continuous variables and a χ^2 test for categorical variables. We grouped individual SES variables into (1) low, (2) middle, and (3) high levels; in *Education*: (1) illiterate, (2) primary school, and (3) secondary school or higher; in *Occupational class*: (1) peasant, (2) manual labourer or housewife, and (3) official/teacher or business/other; in *Satisfactory income*: (1) poor or average, (2) satisfactory, and (3) very satisfactory. The *financial problem* in the past 2 years was recorded as 'yes' or 'no'. In the four-province study, we divided baseline annual personal income and family income into three groups according to their respective tertile cut-off points.

We employed multivariate adjusted Cox regression models to examine the risk of incident stroke in relation to each of SES indicators in the two cohorts, respectively. We calculated the HR and its 95% CIs for incidence of stroke. In the models we adjusted for age, sex, body mass index, smoking, alcohol consumption, marital status, frequency of visiting children/other relatives, hypertension, heart disease, diabetes, activity of daily living (ADL), and depression and dementia. When the covariables had missing data, we took them as a separate group for analysis. Apart from examining the independent effect of one SES indicator from other SES we included all other SES indicators in the models for full adjustment. In the full model data analysis, we investigated the impacts in women and men separately, and tested their gender differences using a ratio of 2 HRs as we did before.¹²

We pooled the two-cohort findings where possible using a meta-analysis method as we did before¹³ to assess the impacts of different SES measurements on incidence of stroke, so as to increase statistical power. All analyses were performed using SPSS V.24.0 (SPSS) and Stata V.14.0 (StataCorp, College Station, Texas, USA).

RESULTS

Anhui cohort study

Of 2852 participants, the average age was 71.7 years (SD 6.9), 51.8% were women and 48.2% lived in rural areas. The distribution of baseline characteristics of participants is shown in online supplementary table 1. Compared with men, women were more likely to have low levels of education, occupation and satisfactory income, be obese, be widowed and have high levels of depression and dementia, but were more likely to have never smoked, not drink, live with children and/or grandchildren and be visited daily by children or other relatives. There were no

Table 1 Number, rate and HR of incident stroke in older people from urban and rural areas in China: the Anhui cohort study

Urban-rural SES variable	Number of strokes/ participants	PYAR* (incidence)	HR ¹ 95% CI	HR ² 95% CI	HR ³ 95% CI	HR ⁴ 95% CI	HR ⁵ 95% CI	HR ⁶ 95% CI
Urban	100/1478	10210.1 (9.79)	1.00	1.00	1.00	1.00	1.00	1.00
Rural	111/1374	8427.6 (13.17)	1.88 1.38 to 2.56	2.07 1.45 to 2.96	1.58 1.00 to 2.48	2.27 1.24 to 4.17	2.44 1.41 to 4.24	2.49 1.19 to 5.22

HR¹: adjusted for age (continuous), sex, body mass index (BMI), smoking, alcohol consumption; HR²: adjusted for age (continuous), sex, BMI, smoking, alcohol consumption, marital status, frequency of visiting children or other relatives, hypertension, heart disease, diabetes, activity of daily living, depression and dementia; HR³: educational level variable was added in the model 2 for adjustment; HR⁴: occupational class variable was added in the model 2 for adjustment; HR⁵: satisfactory income variable was added in the model 2 for adjustment; HR⁶: educational level, occupational class and satisfactory income variables were added in the model 2 for adjustment.

*PYAR (incidence): person-year at risk (incidence rate); incidence rate per 1000 person-years.
SES, socioeconomic status.

significant gender differences in age, urban rurality, financial problem, ADL and cardiovascular comorbidities.

Over the 10 years' follow-up, 211 cases of stroke were identified. Table 1 shows the numbers, incidence and adjusted HRs of stroke in participants living in urban and rural areas. The HR for stroke in participants living in rural areas was significantly increased with adjustment for different sets of confounders,

including other SES indicators. Separate data analysis for women showed a fully adjusted HR of 3.64 (95% CI 1.17 to 11.32) and for men 2.23 (95% CI 0.81 to 6.19) (table 2).

There were no significant associations of education, occupation, satisfactory income and financial problem with incident stroke, although increased HRs in participants with low levels of education and occupation and a financial problem were

Table 2 HRs of incident stroke by socioeconomic status in women and men: the Anhui cohort study

SES variable	Anhui cohort					
	Women			Men		
	Number of strokes/ participants	PYAR (incidence)*	HR 95% CI†	Number of strokes/participants	PYAR (incidence)*	HR 95% CI†
Urban rurality						
Urban	49/770	5415.2 (9.05)	1.00	51/708	4794.8 (10.64)	1.00
Rural	51/707	4490.4 (11.36)	3.64 1.17 to 11.32	60/667	3937.3 (15.24)	2.23 0.81 to 6.19
Educational level‡						
High	19/421	3050.3 (6.23)	1.00	45/632	4307.8 (10.45)	1.00
Middle	10/183	1253.7 (7.98)	1.59 0.66 to 3.85	16/178	1116.1 (14.34)	1.07 0.52 to 2.20
Low	71/873	5601.6 (12.67)	3.68 1.70 to 7.97	50/565	3308.1 (15.11)	0.93 0.44 to 1.98
Occupational class‡						
High	29/470	3333.9 (8.70)	1.00	43/632	4341.0 (9.91)	1.00
Middle	22/299	2074.2 (10.61)	0.73 0.36 to 1.47	14/129	788.1 (17.76)	1.53 0.77 to 3.06
Low	49/708	4497.5 (10.89)	0.43 0.16 to 1.18	54/614	3603.0 (14.99)	1.20 0.49 to 2.94
Satisfactory income‡						
High	13/155	1066.5 (12.19)	1.00	12/139	960.4 (12.49)	1.00
Middle	41/682	4742.0 (8.65)	0.62 0.32 to 1.20	48/705	4637.4 (10.35)	0.74 0.38 to 1.44
Low	46/640	4097.1 (11.23)	0.62 0.30 to 1.28	51/531	3134.3 (16.27)	0.83 0.40 to 1.73
Financial problem in the past 2 years						
No	54/803	5618.0 (9.61)	1.00	52/744	4964.8 (10.47)	1.00
Yes	46/674	4287.5 (10.73)	0.68 0.31 to 1.49	59/631	3767.3 (15.66)	1.74 0.77 to 3.92

*PYAR (incidence): person-year at risk (incidence rate); incidence rate per 1000 person-years.

†Models are adjusted for: age (continuous), sex, body mass index (BMI), smoking, alcohol consumption, marital status, frequency of visiting children or other relatives, hypertension, heart disease, diabetes, activity of daily living, depression and dementia, and urban rurality, education, occupation, satisfactory income (or financial problem).

‡Classification of low, middle and high levels in SES variables: *education* was classified as follows: (1) low: illiterate, (2) middle: primary school, and (3) high: secondary school or higher; *occupational class* was classified as follows: (1) low: peasant, (2) middle: manual labourer or housewife, and (3) high: official/teacher or business/other; *satisfactory income* was classified as follows: (1) low: poor or average, (2) middle: satisfactory, and (3) high: very satisfactory.
SES, socioeconomic status.

Table 3 HRs of incident stroke by occupational class in women and men: the four-province cohort study

Occupational class*	Four-province cohort					
	Women			Men		
	Number of strokes/participants	PYAR† (incidence)	HR‡ 95% CI	Number of strokes/participants	PYAR† (incidence)	HR‡ 95% CI
High	6/235	690.1 (8.69)	1.00	20/358	1029.0 (19.44)	1.00
Middle	15/541	1624.9 (9.23)	1.01 0.35 to 2.93	18/267	797.8 (22.56)	1.20 0.59 to 2.44
Low	26/912	2836.1 (9.17)	0.69 0.22 to 2.12	28/703	2185.7 (12.81)	0.46§ 0.23 to 0.93

*Classification of low, middle and high levels in *occupational class*: (1) low: peasant, (2) middle: manual labourer or housewife, and (3) high: official/teacher or business/other.

†PYAR (incidence): person-year at risk (incidence rate); incidence rate per 1000 person-years.

‡Models are adjusted for: age (continuous), sex, body mass index (BMI), smoking, alcohol consumption, marital status, frequency of visiting children or other relatives, hypertension, heart disease, diabetes, activity of daily living, depression and dementia, and urban rurality, education, satisfactory income.

§HR for incident stroke in men with high occupation versus low occupation was 2.17 (95% CI 1.08 to 4.35).

significant before adjustment for other SES (online supplementary table 2). However, separate data analysis by gender showed that there were significantly increased risks of stroke in women with low education (table 2).

Four-province cohort study

Of 3016 participants, 113 cases of incident stroke occurred over the 3 years' follow-up. Compared with those living in urban areas, participants in rural areas had a fully adjusted HR of 1.34 (95% CI 0.78 to 2.30) of incident stroke (online supplementary table 3). There were no significant associations of education, satisfactory income and financial problem with incident stroke, except for high occupational class appearing associated with increased stroke (online supplementary table 3). Separate data analysis for women and men showed findings similar to those for the combined sample and there were no gender differences in these associations (data not shown), except for men with high occupation having increased the risk of stroke (2.17, 95% CI 1.08 to 4.35) (table 3).

Table 4 shows the numbers, incidence and adjusted HRs of stroke in participants with different levels of annual personal

and family incomes. Significantly increased risk of incident stroke was seen in women with low annual personal income (3.05, 95% CI 1.17 to 8.00) and in men with high family income (2.38, 95% CI 1.14 to 4.76 vs middle income; $p=0.037$ in three group differences).

Pooled data from the Anhui cohort and the four-province cohort

Figure 1 shows pooled data of HRs of incident stroke in people with low SES from the two cohorts. There was a significantly increased risk of stroke in participants living in rural areas, before and after adjustment for other SES variables. The associations of low education and financial problems with incident stroke were significant in the analysis before adjustment for other SES variables. High occupational class was associated with increased risk of stroke after adjustment for other SES variables. Other HRs in figure 1 were not statistically significant.

Pooled data for women and men separately showed that women with rural living or low education had significantly increased risk of stroke, and other HRs were not statistically

Table 4 Number, rate and HR of incident stroke in older people with actual income in China: the four-province cohort study

Annual income (¥)	All participants			Women			Men		
	Number of strokes/participants	PYAR* (incidence)	HR† 95% CI	Number of strokes/participants	PYAR* (incidence)	HR† 95% CI	Number of strokes/participants	PYAR* (incidence)	HR† 95% CI
Personal									
≥10000	48/1377	4061.7 (11.82)	1.00	12/682	2019.9 (5.94)	1.00	36/695	2041.8 (17.63)	1.00
4800 to <10000	21/568	1744.3 (12.04)	1.14 0.60 to 2.19	10/334	1008.1 (9.92)	2.14 0.78 to 5.91	11/234	736.2 (14.94)	0.87 0.37 to 2.07
0 to <4800	43/969	3078.2 (13.97)	1.54 0.83 to 2.86	24/603	1932.9 (12.42)	3.05 1.17 to 8.00	19/366	1145.3 (16.59)	0.96 0.41 to 2.24
Family (per person)									
≥12000	53/1133	3575.4 (14.82)	1.00	15/595	1914.1 (7.84)	1.00	38/538	1661.3 (22.87)	1.00‡
4800 to <12000	25/737	2307.1 (10.84)	0.66 0.40 to 1.10	13/419	1308.5 (9.94)	1.02 0.45 to 2.33	12/318	998.6 (12.02)	0.42 0.21 to 0.88
0 to <4800	24/755	2174.7 (11.04)	0.75 0.37 to 1.50	13/416	1199.4 (10.84)	1.71 0.62 to 4.71	11/339	975.2 (11.28)	0.37 0.13 to 1.07

The top tertile of actual personal and family incomes was treated as the high level. Of 3016 eligible participants in the four-province cohort study, 102 missing occurred in annual personal income and 391 missing in family annual income per person.

*PYAR (incidence): person-year at risk (incidence rate); incidence rate per 1000 person-years.

†Models are adjusted for: age (continuous), sex, body mass index (BMI), smoking, alcohol consumption, marital status, frequency of visiting children or other relatives, hypertension, heart disease, diabetes, activity of daily living, depression and dementia, and urban rurality, education, occupation.

‡The overall p value for family income in men was 0.037.




Figure 1 Pooled HRs for the risk of incident stroke in low socioeconomic status (SES) before and after adjustment for SES from the Anhui cohort and the four-province cohort studies.

significant (table 5). There were no gender differences in the associations of rural living, education, occupation, satisfactory income and financial problems with incident stroke (table 5).

DISCUSSION

Our community-based cohort study examined the impact of different measurements of SES on incidence of stroke in China. It demonstrated that inequality in the risk of stroke was largely for people who lived in rural areas. Women who had low education or low personal income also had increased risk of stroke, as did men with high occupational class or family income.

The association of low education with increased risk of stroke has been observed in the HICs.¹⁵ In Australia women with least

education versus highest education had increased risk of stroke.¹⁶ Our study also found that women with illiteracy had significantly increased risk of stroke compared with their counterparts who attained at least primary school. However, our data from the male population did not show a significant association of low educational level with increased risk of stroke. It may be because other factors, for example, smoking and alcohol drinking in Chinese men, play a more important role in the aetiology of stroke, covering the impact of low education on risk of stroke.

Our study showed that high level of occupational class was associated with increased incidence of stroke, mainly in men. This is different from those in previous studies undertaken in HICs.⁴ One of the reasons for it could be attributed to epidemiological transition in the early stages that those with high occupational class are more likely to have adverse lifestyles including smoking¹⁷ and drinking,¹⁸ sedentary behaviour¹⁹ and psychological stress,²⁰ especially men. These lifestyles are associated with increased risk of stroke.

Our data of the four-province study showed that men with high family income had an increased risk of stroke in comparison to their counterparts with low and middle family income. This could be explained by lifestyle changes among them; the Chinese men with high income experienced and adopted unhealthy lifestyles, such as smoking and drinking alcohol.²¹ The pattern of high income associated with high risk of stroke was also observed in HICs. In USA, the incidence of stroke in people aged above 75 was increased with higher income (HR 2.33, 95% CI 1.16 to 4.55).³ In France, the Three-City Study demonstrated that in those aged over 65 years there was an 80% increased risk of ischaemic stroke in those with higher income compared with lower income (HR 1.77, 95% CI 1.20 to 2.61).⁹ In our study, high income is partly in line with high occupational class, and to a large extent a man's earnings and occupation would determine his family's SES. Therefore, the explanations relating to some risk factors for incident stroke in high occupational class also could be applied to men with high family income.

The current study did not find a significant association of satisfactory income or financial problem with incidence of stroke. Satisfactory level of income involved people's feeling about income, while the financial problem in the past 2 years reflected 'the financial crisis'. These proxies could be good for examining the risk of mental disorders and other chronic diseases.^{22 23} In this study, we analysed the data of the actual income from the four-province study and found that older women with low personal income had increased risk of stroke. The finding is consistent with other studies.²⁴

Compared with other indicators of SES above, the impact of the rural-urban disparity on incident stroke has rarely been studied. Based on existing literature, we have found few published studies of examining differences in incidence rate of stroke between rural and urban areas. In a large-scale population-based cohort study in Portugal, Correia *et al* examined the data from an older population aged 65+ years and found that the annual incidence of stroke per 1000 population was higher in rural (from 9.5 to 20.2) than in urban areas (from 6.8 to 10.9).²⁵ Data from young and middle-aged populations in a study in China also showed that incidence of stroke in rural areas was significantly greater than that of their urban counterparts.²⁶ However in sub-Saharan Africa, a community-based study of the Tanzania Stroke Incidence Project found that there was higher incidence of stroke in older people living in urban versus rural areas, which could be due to high prevalence of stroke risk factors at a community level in the urban

Table 5 Pooled HR of incident stroke in different SES indicators among participants from the Anhui cohort and the four-province cohort studies

SES variable	Women		Men		Ratio of HRs in women versus men	
	HR*	95% CI	HR*	95% CI	RHR†	P value
Urban rurality						
Urban	1.00		1.00			
Rural	2.39	1.18 to 4.83	1.52	0.84 to 2.75	1.57	0.335
Educational level						
High	1.00		1.00			
Middle	0.95	0.47 to 1.94	1.14	0.67 to 1.93	0.83	0.686
Low	2.26	1.19 to 4.27	0.98	0.55 to 1.74	2.31	0.057
Occupational class						
High	1.00		1.00			
Middle	0.81	0.45 to 1.45	1.36	0.83 to 2.23	0.60	0.185
Low	0.53	0.25 to 1.12	0.66	0.38 to 1.15	0.80	0.645
Satisfactory income						
High	1.00		1.00			
Middle	0.70	0.39 to 1.25	1.07	0.44 to 2.61	0.65	0.434
Low	0.73	0.39 to 1.37	0.96	0.52 to 1.77	0.76	0.541
Financial problem in the past 2 years						
No	1.00		1.00			
Yes	0.65	0.31 to 1.36	1.45	0.73 to 2.87	0.45	0.119

*Models are adjusted for: age (continuous), sex, body mass index (BMI), smoking, alcohol consumption, marital status, frequency of visiting children or other relatives, hypertension, heart disease, diabetes, activity of daily living, depression and dementia, and urban rurality, education, occupation, satisfactory income (or financial problem).

†Those RHR: all $p > 0.05$.

RHR, ratio of HR; SES, socioeconomic status.

region.²⁷ In spite of these differences between rural and urban in the incidence of stroke, all three studies did not adjust for confounders to assess the rural-urban disparities in the risk of stroke, and the residual effects could therefore not be removed. Our study, with adjustment for many important confounders, showed that older people living in the rural areas in China had significantly increased risk of stroke.

There was a wide inequality between rural and urban in China due to disparities in education, employment opportunities, income, political rights, social welfare and healthcare services.⁷ In rural areas some risk factors for stroke were common and uncontrolled.²⁶ A health survey in China showed that people living in rural areas had higher prevalence of hypertension and that they were particularly unaware, leaving high levels of untreated and uncontrolled hypertension.²⁸ Primary care and preventive care access remain poor in rural areas,²⁹ including lack of facilities such as CT or MRI scan for the diagnosis of stroke.³⁰ People who live in rural areas might not receive adequate preventive interventions for high-risk factors and thus would have increased risk of stroke.³¹ These could be some possible explanations for an association of living in rural areas with increased incidence of stroke.

In China, apart from the rural-urban inequality, low income has been associated with reduced healthcare use. Luo *et al* found that compared with those from lower income families, people from higher income families were more likely to use outpatient services (OR 1.55, 95% CI 1.05 to 2.29) and inpatient services (1.40, 95% CI 1.00 to 1.98).³² Previous studies showed that there were no gender differences in outpatient visits, but among those with low income women used outpatient services more frequently than men.³² These could partially explain the findings of the current study.

Our study has identified that women were more evident in the disparities of incident stroke in terms of levels of education and income, apart from rurality. The women in our study were born in a turbulent age when most of them had no opportunities to attain schooling and work outside due to many factors, for example, poverty, Chinese tradition. Also, female education was rarely given priority due to poverty and patriarchal attitudes,³³ thus causing a high illiterate rate in older women.³⁴ Older women in China are still among the most economically disadvantaged population groups. Consequently, once they experienced some economic problems, they would have increased risk of incidence of stroke.

Strengths and weaknesses of the study

The main contribution of this study, beyond the intrinsic importance of studying inequality in stroke risk in the largest and most populous LMIC, lies in what it tells us about the simultaneous impacts of multiple measurements of SES on incidence of stroke and their gender differences. As far as we know, our study is the first to report gender differences in the association of SES with incident stroke, addressing that women had higher inequality in incidence of stroke in relation to low SES. Our study included important covariables for adjustment such as ADL and depression and thus the confounding effect would be minimised. Particularly, we adjusted for each of these SES indicators (rurality, education, occupation and income), and thus their residual effects from other SES indicators were removed. Our study has limitations. First, although each of the two cohorts has enough participants to examine incidence of stroke in relation to SES, multiple adjustment analysis gave HR estimates a wide 95% CI, reducing the statistical significance. However, our pooled data could increase the study power. Second, we do

not have data of stroke subtypes in the cohort for analysis, and it is unclear whether or not there were significant differences in the impacts of SES on ischaemic and haemorrhagic strokes. However, studies in other countries showed similar effects of SES on incidence of ischaemic and haemorrhagic strokes.^{35 36} In China the prevalence of hypertension and the proportion of intracerebral haemorrhage (ICH)²⁶ are higher than those in Western countries,³⁷ and previous studies suggested that hypertension has more effects on increased risk of ICH than on acute ischaemic stroke.³⁸ It is unclear what the effects are of both prevalent hypertension and ICH on the association between SES and incident stroke in this study. However, we adjusted for multiple CVD risk factors, including hypertension in the data analysis, and thus the confounding effect of hypertension on the association has been minimised. Nevertheless, we need further studies focused on this issue in China, including detailed data collection using brain scan for stroke subtypes to clarify the effects of these SES indicators on the incident risk of each stroke subtype. Third, in the two cohorts we analysed self-reported satisfactory income associated with incident stroke. The satisfactory income is the person's perception of her/his own income and not the actual income. Although there is significant correlation between this measurement and actual salary income (online supplementary table 4), we did not find its association with risk of stroke. However, in the four-province cohort we used actual income from personal and family and found their associations with incident stroke in women and men. Fourth, the four-province cohort study had a 27% loss to follow-up rate. This was because during that period China had many residential area houses reformed and rearranged due to rapid development, and some participants moved home and we could not trace their new house address. In the Anhui cohort, we also had a number of cohort members lost to follow-up, for example, at wave 3 only 1757 were reinterviewed (excluding 565 deaths), which gave a 30% loss to follow-up rate, although the overall loss to follow-up rate across the Anhui cohort wave 2–4 interviews and identified mortality was lower (10.7%).¹² We do not know whether those lost to follow-up had higher or lower level of incident stroke than other cohort members. The bias, if any, could be in any direction for the association of SES with incident stroke. However, the loss to follow-up rate in our study is similar to those in some studies undertaken in Western countries,³⁹ and there is need of further studies with lower loss to follow-up rate to examine the association of SES with incident stroke in China. Fifth, caution should be exercised in generalising the findings to the whole population of China. The impact of SES on the risk of stroke in our cohort study consists of participants aged ≥ 60 years old in urban and rural communities. Those aged < 60 years need to be included in further research.

CONCLUSION

This study demonstrates inequality in the risk of stroke in terms of SES in China. The rurality was a main resource of incident stroke inequality. There were gender differences in the impact of SES on stroke risk; in women increased risk of stroke was associated with low education and low personal income, while in men it was with high occupational class and high family income. The gender-specific strategies and preventive interventions of health promotion targeting people living in rural areas, through reducing socioeconomic deprivation, would be helpful in campaigns to reduce stroke incidence in China. Increasing public health education and improving stroke care access in rural areas, particularly in women, and constructing

balanced policies for rural seniors could have large impacts in reducing the burden of stroke in China.

What is already known on this subject

- ▶ Many studies in high-income countries (HIC) showed that low socioeconomic status (SES) was associated with increased incidence of stroke. There is a lack of data from low and middle-income countries and no study has been done in China where the stroke is the number one killer to examine the association of SES with incidence of stroke.
- ▶ The findings of previous studies in HICs were not consistent in assessing the impact of SES on incidence of stroke. Few studies have used multiple measurements of SES to examine their associations with incident stroke simultaneously. No study has been done to investigate gender differences in the associations.

What this study adds

- ▶ In China, living in rural areas was associated with increased risk of incident stroke in terms of SES.
- ▶ In women, those with low levels of education and personal income had a higher risk of incident stroke than their counterparts.
- ▶ In men, those with high levels of occupational class and family income had a higher risk of incident stroke than their counterparts.
- ▶ Health promotion in rural areas and gender-specific targets would help reduce health inequality and incidence of stroke in China.

Author affiliations

- ¹Faculty of Education, Health and Wellbeing, University of Wolverhampton, Wolverhampton, UK
- ²Department of Rehabilitation, Shenzhen Second People's Hospital, Shenzhen University First Affiliated Hospital, Shenzhen, Guangdong, China
- ³Department of Preventive Medicine, School of Public Health, Guangzhou Medical University, Guangzhou, Guangdong, China
- ⁴Department of Health Technology and Informatics, The Hong Kong Polytechnic University, Hong Kong, China
- ⁵Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China
- ⁶China National Clinical Research Center for Neurological Diseases, Beijing, China
- ⁷Royal Wolverhampton Hospitals NHS Trust, Wolverhampton, UK
- ⁸Department of Epidemiology and Biostatistics, School of Public Health, Guangdong Medical University, Dongguan, Guangdong, China
- ⁹Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK
- ¹⁰Department of Public Health and Epidemiology, University College London, London, UK

Acknowledgements The authors thank the participants and all who were involved in the Anhui cohort and the four-province cohort studies. Dr Yulong Wang is the recipient of a grant from Shenzhen Health and Family Planning Commission, Shenzhen, Guangdong, China for the training of rehabilitation ability of severe stroke patients (project no. SM201623A).

Contributors RC (guarantor) had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. RC obtained funding, conceptualised and designed the study, supervised the analyses and revised the manuscript. WZ had full access to all the data in the study, managed the database, performed the analyses and drafted the initial manuscript. AH, YW, XC, AC, KF and JN commented and revised the manuscript and interpreted the findings. JT, YP and DW provided statistical support and input, interpreted the data and commented on the manuscript. EB commented and supported the data analysis, interpreted the findings and provided critical revision of the manuscript. All authors approve the final version of the manuscript.

Funding The research was funded by the BUPA Foundation (grants numbers 45NOV06 and TBF-M09-05) and the Alzheimer's Research UK (grant number ART/PPG2007B/2) for the two cohorts' data collection.

Disclaimer The funders had no role in the study design, data analysis and interpretation, the writing of the manuscript and the decision to submit the manuscript for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethical approvals for the Anhui cohort study and the four-province study were obtained from the Ethics Committees of University College London, and School of Health and Wellbeing at University of Wolverhampton, UK, and the Research Ethics Committee of Anhui Medical University.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Weiju Zhou <http://orcid.org/0000-0001-5077-3202>
Eric Brunner <http://orcid.org/0000-0002-0595-4474>

REFERENCES

- 1 Kyu HH, Abate D, Abate KH, *et al*. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2018;392:1859-922.
- 2 Addo J, Ayerbe L, Mohan KM, *et al*. Socioeconomic status and stroke: an updated review. *Stroke* 2012;43:1186-91.
- 3 Avendano M, Kawachi I, Van Lenthe F, *et al*. Socioeconomic status and stroke incidence in the US elderly. *Stroke* 2006;37:1368-73.
- 4 McFadden E, Luben R, Wareham N, *et al*. Social class, risk factors, and stroke incidence in men and women. *Stroke* 2009;40:1070-7.
- 5 Johnson CO, Nguyen M, Roth GA, *et al*. Global, regional, and national burden of stroke, 1990-2016: a systematic analysis for the global burden of disease study 2016. *Lancet Neurol* 2019;18:439-58.
- 6 Feigin VL, Nguyen G, Cercy K, *et al*. Global, regional, and Country-Specific lifetime risks of stroke, 1990 and 2016. *N Engl J Med* 2018;379:2429-37.
- 7 Sicular T, Ximing Y, Gustafsson B, *et al*. The urban-rural income gap and inequality in China. *Rev Income Wealth* 2007;53:93-126.
- 8 Avendano M, Glymour MM. Stroke disparities in older Americans. *Stroke* 2008;39:1533-40.
- 9 Grimaud O, Dufouil C, Alperovitch A, *et al*. Incidence of ischaemic stroke according to income level among older people: the 3C study. *Age Ageing* 2011;40:116-21.
- 10 Han TS, Wang HH-X, Wei L, *et al*. Impacts of undetected and inadequately treated hypertension on incident stroke in China. *BMJ Open* 2017;7:e016581.
- 11 Chen R, Hu Z, Chen R-L, *et al*. Determinants for undetected dementia and late-life depression. *Br J Psychiatry* 2013;203:203-8.
- 12 Chen R, Hu Z, Wei L, *et al*. Socioeconomic status and survival among older adults with dementia and depression. *Br J Psychiatry* 2014;204:436-40.
- 13 Chen R, Hu Z, Chen R-L, *et al*. Socioeconomic deprivation and survival after stroke in China: a systematic literature review and a new population-based cohort study. *BMJ Open* 2015;5:e005688.
- 14 Engstad T, Bonna KH, Viitanen M. Validity of self-reported stroke: the Tromsø study. *Stroke* 2000;31:1602-7.
- 15 Qureshi AI, Suri MFK, Saad M, *et al*. Educational attainment and risk of stroke and myocardial infarction. *Med Sci Monit* 2003;9:CR466-73.
- 16 Jackson CA, Sudlow CLM, Mishra GD. Education, sex and risk of stroke: a prospective cohort study in New South Wales, Australia. *BMJ Open* 2018;8:e024070.
- 17 Wang Q, Shen JJ, Sotero M, *et al*. Income, occupation and education: are they related to smoking behaviors in China? *PLoS One* 2018;13:e0192571.
- 18 Wu B, Mao Z-F, Rockett IRH, *et al*. Socioeconomic status and alcohol use among urban and rural residents in China. *Subst Use Misuse* 2008;43:952-66.
- 19 Chen M, Wu Y, Narimatsu H, *et al*. Socioeconomic status and physical activity in Chinese adults: a report from a community-based survey in Jiaying, China. *PLoS One* 2015;10:e0132918.
- 20 Yang T, Rockett IRH, Lv Q, *et al*. Stress status and related characteristics among urban residents: a six-province capital cities study in China. *PLoS One* 2012;7:e30521.
- 21 Kim S, Symons M, Popkin BM. Contrasting socioeconomic profiles related to healthier lifestyles in China and the United States. *Am J Epidemiol* 2004;159:184-91.

- 22 Chen R, Ma Y, Wilson K, et al. A multicentre community-based study of dementia cases and subcases in older people in China—the GMS-AGECAT prevalence and socio-economic correlates. *Int J Geriatr Psychiatry* 2012;27:692–702.
- 23 Chen R, Song Y, Hu Z, et al. Predictors of diabetes in older people in urban China. *PLoS One* 2012;7:e50957.
- 24 Li C, Hedblad B, Rosvall M, et al. Stroke incidence, recurrence, and case-fatality in relation to socioeconomic position: a population-based study of middle-aged Swedish men and women. *Stroke* 2008;39:2191–6.
- 25 Correia M, Silva MR, Matos I, et al. Prospective community-based study of stroke in northern Portugal: incidence and case fatality in rural and urban populations. *Stroke* 2004;35:2048–53.
- 26 Wang W, Jiang B, Sun H, et al. Prevalence, Incidence, and Mortality of Stroke in China: Results from a Nationwide Population-Based Survey of 480 687 Adults. *Circulation* 2017;135:759–71.
- 27 Walker R, Whiting D, Unwin N, et al. Stroke incidence in rural and urban Tanzania: a prospective, community-based study. *Lancet Neurol* 2010;9:786–92.
- 28 Li W, Gu H, Teo KK, et al. Hypertension prevalence, awareness, treatment, and control in 115 rural and urban communities involving 47000 people from China. *J Hypertens* 2016;34:39–46.
- 29 Huang C, Liu C-J, Pan X-F, et al. Correlates of unequal access to preventive care in China: a multilevel analysis of national data from the 2011 China health and nutrition survey. *BMC Health Serv Res* 2016;16:177.
- 30 Liu M, Wu B, Wang W-Z, et al. Stroke in China: epidemiology, prevention, and management strategies. *Lancet Neurol* 2007;6:456–64.
- 31 Li J, Shi L, Liang H, et al. Urban-Rural disparities in health care utilization among Chinese adults from 1993 to 2011. *BMC Health Serv Res* 2018;18:102.
- 32 Luo J, Zhang X, Jin C, et al. Inequality of access to health care among the urban elderly in northwestern China. *Health Policy* 2009;93:111–7.
- 33 Yu MY, Sarri R. Women's health status and gender inequality in China. *Soc Sci Med* 1997;45:1885–98.
- 34 Li J. Women's status in a rural Chinese setting. *Rural Sociol* 2005;70:229–52.
- 35 Seo SR, Kim SY, Lee S-Y, et al. The incidence of stroke by socioeconomic status, age, sex, and stroke subtype: a nationwide study in Korea. *J Prev Med Public Health* 2014;47:104–12.
- 36 Pujades-Rodríguez M, Timmis A, Stogiannis D, et al. Socioeconomic deprivation and the incidence of 12 cardiovascular diseases in 1.9 million women and men: implications for risk prediction and prevention. *PLoS One* 2014;9:e104671.
- 37 Feigin VL, Lawes CMM, Bennett DA, et al. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *Lancet Neurol* 2009;8:355–69.
- 38 O'Donnell MJ, Xavier D, Liu L, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010;376:112–23.
- 39 Vega S, Benito-León J, Bermejo-Pareja F, et al. Several factors influenced attrition in a population-based elderly cohort: neurological disorders in central Spain study. *J Clin Epidemiol* 2010;63:215–22.