

Low rate of bacterial co-infection in patients with COVID-19

We agree with Michael J Cox and colleagues¹ that clinical management of COVID-19 would be enhanced by further characterisation of bacterial co-infections. A few case reports have described examples of such co-infections.^{2,3,4} However, national⁵ and international⁶ guidelines recommend empirical antibiotics for all patients who are severely ill with suspected COVID-19, and that cessation of therapy is left to the clinicians' discretion. Pending the widespread availability of metagenomic sequencing as envisaged by Cox and colleagues,¹ we argue that traditional diagnostics still have a role.

We reviewed all microbiology results for patients admitted to Whiston hospital (Prescot, UK) with PCR-confirmed COVID-19 between March 6, 2020, and April 7, 2020. Hospital policy for patients admitted with community-acquired pneumonia, including suspected COVID-19 cases, recommends blood cultures and pneumococcal and *Legionella* urinary antigen tests based on clinical severity, in line with national guidelines.⁷ We collected the data to inform and update the hospital's antimicrobial policy, with approval from the Trust Quality Improvement and Clinical Audit department. We recorded results for 7 days from the positive COVID-19 test because positive samples collected after this time period might represent hospital-acquired infections. Samples unequivocally consistent with contamination were considered negative.

We identified 195 patients (for demographics and microbiology see

appendix p 1). Five (3% of 195, or 4% of 137 patients specifically tested), had pneumococcal co-infection and all survived to hospital discharge. One of 31 patients tested was positive for the *Legionella* antigen without lower respiratory tract samples to confirm legionellosis. Bacteria grew from four of 26 sputum samples (appendix p 1). All bacteria were Gram-negative bacilli more typically associated with oropharyngeal colonisation than community-acquired pneumonia.

Our findings suggest that bacterial co-infection is uncommon in patients with COVID-19 who are newly admitted to hospital. The co-prevalence of pneumococcus and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was low, and *Staphylococcus aureus* was not detected. By contrast, in influenza infection the prevalence of bacterial co-infection in hospitalised patients can exceed 30%.^{8,9} These results suggest that routine antibiotics might not be indicated in patients with COVID-19. If superimposed bacterial community-acquired pneumonia is suspected, coverage for typical pathogens such as pneumococcus might suffice, unless there is specific clinical concern for infection with atypical agents.

The main strength of our report is the correlation of microbiology results with all consecutive COVID-19 admissions. The main limitation is the variability of microbiological sampling. Our results might not be generalisable to other geographical settings.

Future studies should implement standardised microbiological sampling for all COVID-19 admissions and prospectively correlate the prevalence of co-infection with mortality rates. Such studies could also correlate clinical and laboratory findings with the presence of

co-infection to support rational prescribing of antibiotics.

We declare no competing interests.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

*Hugh Adler†, Robert Ball†,
Michael Fisher, Kalani Mortimer,
Madhur S Vardhan
hugh.adler@sthk.nhs.uk

†Joint first authors

Department of Microbiology, St Helens and Knowsley Teaching Hospitals NHS Trust, Prescot, L35 5DR, UK (HA, RB, MF, KM, MSV); and Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK (HA)

- Cox MJ, Loman N, Bogaert D, O'Grady J. Co-infections: potentially lethal and unexplored in COVID-19. *Lancet Microbe* 2020; **1**: e11.
- Duployez C, Le Guern R, Tinez C, et al. Pantovaleurine leukocidin-secreting *Staphylococcus aureus* pneumonia complicating COVID-19. *Emerg Infect Dis* 2020; published online April 16. DOI:10.3201/eid2608.201413.
- Edrara EM, Lopez EB, Villarama JB, et al. First COVID-19 infections in the Philippines: a case report. *Trop Med Health* 2020; **48**: 21.
- Khaddour K, Sikora A, Tahir N, Nepomuceno D, Huang T. Case report: the importance of novel coronavirus disease (COVID-19) and coinfection with other respiratory pathogens in the current pandemic. *Am J Trop Med Hyg* 2020; published online April 17. DOI:10.4269/ajtmh.20-0266.
- NHS. Clinical management of persons admitted to hospital with suspected COVID-19 infection. 2020. <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/clinical-management-of-persons-admitted-to-hospita-v1-19-march-2020.pdf> (accessed April 14, 2020).
- WHO. Clinical management of severe acute respiratory infection when COVID-19 is suspected. 2020. [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected) (accessed April 14, 2020).
- Lim WS, Baudouin SV, George RC, et al. BTS guidelines for the management of community acquired pneumonia in adults: update 2009. *Thorax* 2009; **64**: iii1-iii55.
- Chertow DS, Memoli MJ. Bacterial coinfection in influenza: a grand rounds review. *JAMA* 2013; **309**: 275-82.
- Bisno AL, Griffin JP, Van Epps KA, Niell HB, Rytel MW. Pneumonia and Hong Kong influenza: a prospective study of the 1968-69 epidemic. *Am J Med Sci* 1971; **261**: 251-63.



See Online for appendix