

## Review article

# Psychosocial interventions on perinatal depression in China: A systematic review and meta-analysis

Juan Yin<sup>a</sup>, Anum Nisar<sup>b</sup>, Ahmed Waqas<sup>c,d</sup>, Yan Guo<sup>a</sup>, Wen Li Qi<sup>e</sup>, Duolao Wang<sup>f</sup>, Atif Rahman<sup>g,1</sup>, Xiaomei Li<sup>a,1,\*</sup>

<sup>a</sup> School of nursing, Xi'an Jiaotong University, 76 West Yanta Road, 710061 Xi'an, China

<sup>b</sup> School of Public Health, Xi'an Jiaotong University, Xi'an, China

<sup>c</sup> Human Development Research Foundation, Rawalpindi, Pakistan

<sup>d</sup> University of Liverpool, Liverpool, UK

<sup>e</sup> Department of obstetrics, Xi'an Aerospace general hospital, Xi'an, China

<sup>f</sup> Liverpool School of Tropical Medicine, Liverpool, UK

<sup>g</sup> Institute of Life and Human Sciences, University of Liverpool, Liverpool, UK

## ARTICLE INFO

## Keywords:

Perinatal depression

China

Psychosocial interventions

Systematic review and meta-analysis

## ABSTRACT

**Background:** The prevalence of perinatal depression is 16.3% in China and has shown a rising trend in the last decade. However, few studies summarized psychosocial interventions for perinatal depression in this country. This study aimed at evaluating and characterizing psychosocial interventions for perinatal depression in Mainland China.

**Methods:** Ten major English and Chinese language electronic bibliographic databases were searched for RCTs examining the effect of psychosocial interventions for perinatal depressed women in Mainland China. Studies meeting eligibility criteria and published before 25th February 2019 were included, while those focusing on a very specific sub-population or reporting non-psychosocial interventions were excluded. Data was extracted by a standard form. Meta-analysis was conducted to obtain a summary measure of the effectiveness of the interventions in reducing perinatal depressive symptoms. The theoretical underpinnings and implementation processes of the interventions were also characterised.

**Results:** A total of 6857 articles were identified in the initial database searching, of which, 26 studies were eligible for data analysis, representing a sample size of 4673. Meta-analysis indicated that psychosocial interventions in China significantly reduced perinatal depressive symptoms (standard difference in means 0.81, 95% confidence intervals -1.03 to -0.58,  $P < 0.001$ ). However, the overall evidence presented substantial heterogeneity ( $I^2 = 91.12\%$ ). Most interventions were implemented in hospitals in urban areas by non-specialist health care providers. Few studies reported details of implementation procedures or scale-up strategies.

**Limitations:** The evidence in this review is of moderate to low quality and therefore, should be interpreted with caution. Some of the trials were inadequately powered and tended to overestimate effect sizes.

**Conclusions:** Current psychosocial interventions in China are somewhat effective in reducing perinatal depressive symptoms. High quality RCTs on scale-up interventions are required, especially in rural areas.

## 1. Introduction

Perinatal depression is typically defined as major or mild depressive episodes, which occurs during pregnancy or within 12 months after delivery (Gynecologists, 2018). Associated with a number of cultural, social and economic stressors, pregnancy and the postpartum period put women at a greater risk of depressive disorder (Biaggi *et al.*, 2016).

The prevalence of perinatal depression has been estimated at between 10% to 15% in High-Income Countries (HIC) and 19–25% in Low- and Middle Income countries (LMIC) (Gelaye *et al.*, 2016; Woody *et al.*, 2017).

Untreated perinatal depression is reported to be associated with various adverse health outcomes for the mothers, including poor nutrition, substance misuse, increased obstetric complications and suicide

\* Corresponding author.

E-mail address: [roselee@xjtu.edu.cn](mailto:roselee@xjtu.edu.cn) (X. Li).

<sup>1</sup> Atif Rahman and Xiaomei Li are joint senior authors

(Bauer *et al.*, 2016; Evans *et al.*, 2001). It has also been linked with impaired health of infants, including poor physical and mental development and increased risk for later common mental disorders in the offspring (Gelaye *et al.*, 2016; Herba *et al.*, 2016; Weissman *et al.*, 2006). Besides, perinatal depression causes a significant medical cost to the family and society. It is estimated that the total lifetime costs of perinatal depression in UK were £75,728 per woman, aggregating to £6.6 billion (Bauer *et al.*, 2016).

China, the world's most populous country, has seen remarkable improvement in maternal and infant health indicators in recent decades (Ministry of Foreign Affairs, P. R. C. July, 2015). In contrast, the prevalence of perinatal depression remains high (15–18%) and has increased during the last decades (Nisar *et al.*, 2019). The high prevalence of perinatal depression indicates an urgent need for interventions in China, particularly psychosocial interventions that are acceptable, feasible and affordable (Richter *et al.*, 2017). There has been some research exploring the management of perinatal depression in China. However, to our knowledge, there are no systematic reviews and meta-analyses providing a synthesis of psychological and psychosocial interventions in mainland China. This report aims to quantize evidence pertaining to effectiveness of psychological and psychosocial interventions for perinatal depression in China. In addition, we also want to summarize and critically evaluate the theoretical underpinnings of these interventions and their implementation processes.

## 2. Methods

We documented the research process following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

### 2.1. Search strategy and selection criteria

A bilingual team of two researchers systematically searched the following ten English and Chinese language electronic bibliographic databases: Medline, EMBASE, Scopus, CINAHL, PsycINFO, Web of Science, Cochrane Central Register, the China National Knowledge Infrastructure, the VIP Database for Chinese Technical Periodicals, and the Wan Fang Database for Chinese Periodicals, using a pretested search strategy (Appendix Table A1). There was no restriction of publication date and all the studies available from the inception of the databases till 25th February, 2019 were incorporated. The inclusion criteria are as follows:

**Participants:** Studies focused on pregnant women or women within 12 months after delivery in Mainland China.

**Intervention:** Studies examining the effect of psychosocial interventions on perinatal depression, either prevention or treatment.

**Control:** Routine care or blank control.

**Outcomes:** Measurement and reporting of perinatal depression scores using a validated self-report scale (e.g., EPDS, SDS, PHQ-9) or clinician-administered measure.

**Study design:** Studies with randomized controlled trials (RCTs) or cluster RCTs design were included.

The exclusion criteria are: Studies with participants from a very specific sub-population (e.g., people with a specific disease or condition or epidemiological character); Studies with insufficient outcome data for meta-analysis. The protocol of this review can be found at <https://www.crd.york.ac.uk/prospero/#record> (CRD42018115934).

Duplicate articles were identified using EndNote, and the papers with more complete data were included for repeat publications. Two investigators worked independently screening the titles and abstracts for inclusion, followed by the full-text screening. Manual searching of the bibliography of all included articles was conducted lastly.

Consensus on papers to be included was reached after discussion and a third researcher was involved when necessary.

### 2.2. Data extraction

Two independent reviewers extracted the following information with a standard form based on Cochrane handbook: (1)General information including study design, sample characteristics, geographical setting, details of intervention, type and training of delivery agent, outcome measures; (2)Data for meta-analysis including sample size, primary outcome, effect size; (3)Therapeutic techniques used in the intervention. Risk of bias of individual research was evaluated by two reviewers according to Cochrane risk of bias tool. Differences in data extraction and risk of bias were resolved by discussion between the reviewers and the principle author.

Besides the above information, the psychosocial interventions were thoroughly audited to identify the broader elements and techniques employed in them. For the phase of data extraction, taxonomy of elements of psychosocial interventions employed in common mental problems was adopted after discussion with two psychologists (Singla *et al.*, 2017). The finalized taxonomy included 58 most commonly utilized treatment elements that could be treatment specific or nonspecific. Definitions of therapeutic elements employed in psychosocial interventions are summarized in Appendix Table A2.

### 2.3. Meta-analyses

The effect sizes of individual study were listed by standard difference in means and 95% confidence intervals (95% CIs) in a forest plot. A fixed-effect model was used if no statistical difference in heterogeneity was found; otherwise, a random-effect model was adopted. The statistical heterogeneity was quantified by the  $I^2$  statistic and formally tested by Cochrane's Q statistic. Publication bias was visually judged by the symmetry of a funnel plot and the result of Egger's linear regression test. Post-hoc sensitivity analyses were performed (using leave-one-out analysis) to test the impact of exclusion of a single trial with a disproportionately large effect. Subgroup analyses were used to explore the sources of heterogeneity according to quality of the study, scale, theoretical orientation, specificity, scope of intervention and elements of interventions. Studies with two or less items ranked as high risk were classified as high quality, otherwise were classified as low quality (Naveed, S. *et al.*, 2019). Scope of intervention was classified as treatment and prevention. Reference to justify perinatal depression had to be defined according to a diagnostic interview based on clinical criteria (CCMD-3, DSM-5 or ICD-10), or as a score above a cut-off on a self-rating depression scale (PHQ-9, SDS, HAMD, et al). Studies focusing on perinatal depressed mothers with a clear diagnosis of "perinatal depression" (based on ICD-10, CCMD-3 or DSM-5) was classified as "treatment". Studies focusing on high risk mothers or mothers without perinatal depression (identified by screening with a validated scale, e.g., EPDS, SDS, PHQ-9, HAMD) was classified as "prevention". Studies focusing on all mothers (without diagnosis or screening for perinatal depression) was classified as others. Statistical significance was set at  $P$  value < 0.05 (two-tailed test). All the analyses were carried out by the Comprehensive meta-analysis software (version 3, BIOSTATS, 2014).

## 3. Results

A total of 10,311 articles were found through database searching and 3454 duplications were removed. After title and abstract screening, 198 articles were identified eligible for full text screen, and 26 studies were finally included in the data analysis. Study selection process is illustrated as a PRISMA flowchart in Fig. 1.

A total of 26 trials of psychosocial interventions representing a sample size of 4673 perinatal women were included. The age of perinatal women included in the studies ranged from 17 to 45 years old




Fig. 1. Study selection process for psychosocial interventions of perinatal depression in women in China.

(29.37 ± 1.87). For the study design, 25 studies (Cheng *et al.*, 2016; Dou, 2018; Gao *et al.*, 2010; Gao *et al.*, 2015; Guan *et al.*, 2015; Ho *et al.*, 2009; Huang *et al.*, 2015; Jia *et al.*, 2017; Jiang *et al.*, 2014; Leung *et al.*, 2016; Leung and Lam, 2012; Li, 2018; Liu *et al.*, 2012; Liu *et al.*, 2018; Lu and Fan, 2017; Lu *et al.*, 2016; Mao *et al.*, 2012; Ngai *et al.*, 2015; Sun *et al.*, 2011; Wu, 2018; Wu, 2017; Yang *et al.*, 2019; Zhang *et al.*, 2018; Zhang *et al.*, 2016; Zhao, 2018) were RCTs and one (Ngai *et al.*, 2009) reported cluster RCT. Nineteen studies focused on postnatal depression, three on prenatal depression and two on both prenatal and postnatal depression. The most commonly used psychometric scales for assessment of perinatal depression were the Edinburgh Postnatal Depression Scale (14 studies), followed by Zung Self-Rating Depression Scale (six studies), Hamilton depression scale (three studies) and Patient Health Questionnaire-9 (three studies). The primary time point of outcome assessment varied from after intervention to four-month post intervention, with ten studies reported follow-up data. Details can be found in Table 1 and Appendix Table A3.

Regarding to the intervention delivery, 22 studies were conducted in urban areas, two in semi-urban areas, and two did not indicate the research region. All studies delivered the interventions in hospitals or health care centers. The interventions were mostly provided by non-specialist health care workers, with six by nurses, four by midwives, and eight by multidiscipline health care workers (seven studies did not report the

delivery agent). As for the type of interventions, 17 studies focused on preventive interventions while the rest on treatment. The duration of overall intervention program varied from one day to eight weeks. Nineteen out of the 26 studies had more than one intervention sessions. Two studies delivered brief interventions—a one-session education program before discharge with a follow-up session (Gao *et al.*, 2015; Ho *et al.*, 2009). Five studies did not report detailed information regarding to the duration or the number of sessions of the whole program. The interventions were delivered in a variety of ways. The most commonly used format was face-to-face or group. Two studies delivered the intervention by talking with women on the telephone (Cheng *et al.*, 2016), (Ngai *et al.*, 2015). Yang 2019 (Yang *et al.*, 2019) and Zhao 2018 (Zhao, 2018) conducted the intervention on a popular Chinese smartphone APP named WeChat, with all sessions recorded and uploaded to the platform. Half of the studies integrated interventions into an existing health education program or routine care. With respect to the fidelity of implementation, only two studies (Leung and Lam, 2012; Ngai *et al.*, 2015) reported detailed information regarding supervision of the intervention, with intervention sessions videotaped and reviewed (Leung and Lam, 2012; Ngai *et al.*, 2015). Detailed information is listed in Table 2.

The most commonly employed therapeutic approaches included cognitive behavioral (CBT, seven studies), psychoeducation (PE, seven studies), interpersonal therapy (IPT, three studies), mindfulness (two

**Table 1**  
Summary of the included articles ( $n = 26$ ).

Author, year	Study design	Theoretical basis	Sample size	Intervention	Delivery agent	Control	Outcome
Dou, 2018	Prevention RCT	CBT	100	Cognitive behavior treatment consisted of 30-min sessions twice a week from inclusion until 6 weeks postpartum.	NA	Routine care: antepartum health education	Postnatal depressive symptoms: 1, 1.5, 3 and 6 months postpartum using SDS
Huang, 2015	Prevention RCT	CBT	186	Emotional management based on CBT, delivered during pregnancy.	Obstetrician psychiatrist	Routine prenatal care and education on nutrition and health knowledge during pregnancy; various modes of delivery; breastfeeding benefits	Postnatal depressive symptoms: after intervention, 7 and 42 days postpartum using PHQ-9
Leung, 2016	Treatment RCT	CBT	164	Six-session group intervention aimed to change cognitions and subsequently reinforce coping skills to enhance psychological resources and responses, delivered during postpartum.	NA	A booklet that contained comprehensive information and education material about perinatal depression and a list of community resources	Postnatal depressive symptoms: 3 and 6 months after intervention using EPDS
Li, 2018	Treatment RCT	CBT	88	Empathy in nursing, identifying the auto-thoughts, discuss the unhealthy thoughts, change the way of thinking, assigning homework, delivered during postpartum.	Nurse	Routine nursing care	Postnatal depressive symptoms: after intervention using HAMD
Lu, 2017	Treatment RCT	CBT	88	Acceptance and commitment therapy consisted of 90 min session per week for 6 weeks, delivered during postpartum.	Researcher, Nurse, Psychiatrist	Routine care and education	Postnatal depressive symptoms: 6 weeks after intervention using HAMD-24
Mao, 2012	Prevention RCT	CBT	240	Emotional self-manage group training program consisted of four weekly group sessions and one individual counseling session, delivered from 32 weeks of pregnancy.	Nurses	Standard antenatal education focusing on childbirth consisted of four 90-min sessions.	Postnatal depressive symptoms: after intervention using EPDS and 6 weeks postpartum using PHQ-9
Ngai, 2015	Treatment RCT	CBT	397	Telephone-Based Cognitive-Behavioral Therapy involved 5-week telephone-administered CBT delivered weekly from 1 to 5 weeks postpartum.	Midwives	6-week postpartum follow-up at public hospitals and maternal and child health centers, respectively.	Postnatal depressive symptoms: 6 weeks and 6 months postpartum using EPDS
Ho, 2009	Prevention RCT	Psychoeducation	175	A discharge education program: postpartum depression informational booklet plus discussion with primary care nurses on the second day after delivery.	Nurses	General postpartum education	Postnatal depressive symptoms: 6 weeks and 3 months postpartum using EPDS
Jia, 2017	Treatment RCT	Psychoeducation	94	Recall positive feelings; build up positive thinking; social support, delivered during postpartum.	Nurse	Routine care: antepartum health education	Postnatal depressive symptoms: 3 days after intervention using EPDS
Lu, 2016	Treatment RCT	Psychoeducation	78	Health education on postpartum depression, exercise consisted of 60 min sessions for 8 weeks, delivered during postpartum	Nurses Doctors	Routine care: answering the questions	Postnatal depressive symptoms: 8 weeks after intervention using EPDS
Sun, 2011	Prevention RCT	Psychoeducation	194	Health education based on the principles of interpersonal psychotherapy, consisted of two 90 min antenatal class during late pregnancy and telephone follow-up within two weeks after delivery.	Nurse	Routine care: antepartum health education	Postnatal depressive symptoms: 6 weeks and 3 months postpartum using EPDS
Wu, 2017	Treatment RCT	Psychoeducation	110	Follow up education on recovery, newborn nursing, and breastfeeding through telephone call, peer group discussion, delivered during postpartum.	Doctors Nurses	Routine care: postpartum health education	Postnatal depressive symptoms: after intervention using SDS
Zhao, 2018	Treatment RCT	Psychoeducation	296	Follow-up education on perinatal depression, diet, emotion management by WeChat APP, delivered during pregnancy until postpartum.	NA	Routine care	Postnatal depressive symptoms: after intervention using SDS
Ngai, 2009	Prevention Cluster RCT	Psychoeducation	184	Childbirth psychoeducation program focused on cognitive restructuring, problem-solving and efficacy enhancement, consisted of three 1 h sessions, delivered during pregnancy.	Midwives	Routine childbirth education programs consisted of 6-weekly 2 h sessions focusing on prenatal care, labor, pain relief, breastfeeding, postnatal and infant care.	Prenatal and postnatal depressive symptoms: after intervention, 6 weeks and 6 months postpartum using EPDS
Gao, 2010	Prevention RCT	Interpersonal	194	Interpersonal-psychotherapy-oriented childbirth education program consisted of two 2 h group sessions during pregnancy and one telephone follow-up in the postpartum period.	Midwives	Routine antenatal education in the study venue, which consisted of two 90-min sessions conducted by midwives.	Postnatal depressive symptoms: 6 weeks postpartum using EPDS
Gao, 2015	Prevention RCT	Interpersonal	180	Interpersonal-psychotherapy-oriented postnatal psychoeducation program consisted of a 1 h	Midwives		Postnatal depressive symptoms: 6 weeks postpartum using EPDS

(continued on next page)

Table 1 (continued)

Author, year	Study design	Theoretical basis	Sample size	Intervention	Delivery agent	Control	Outcome
Leung, 2012	Treatment RCT	Interpersonal	156	education session before discharge and one telephone follow-up session after delivery. Interpersonal psychotherapy oriented group intervention targeted interpersonal issues consisted of 4 weekly group sessions lasting 1.5 h per session, delivered during pregnancy.	First author Interventionist	A brief visit from a nurse to give them a pamphlet on sources of assistance for mothers on discharge from hospital Routine antenatal care including a brief individual interview with a midwife during which participants could raise any health or pregnancy related questions or concerns.	Prenatal and postnatal depressive symptoms: after intervention and 6–8 weeks postpartum using EPDS
Yang, 2019	Prevention RCT	Mindfulness	123	8-weeks online mindfulness intervention based on WeChat APP focusing on attention monitoring and acceptance, delivered during pregnancy.	Nurses midwife	Routine care: antepartum health education related to childbirth, breastfeeding, nutrition, and parenting.	Prenatal depressive symptoms: after intervention using PHQ-9
Zhang, 2018	Prevention RCT	Mindfulness	63	Mindfulness-based stress reduction program consisted of weekly 90 min sessions for 8 weeks, delivered during pregnancy.	Psychologist	Routine care: Prenatal care knowledge	Prenatal depressive symptoms: after intervention using SDS
Jiang, 2014	Treatment RCT	Counseling	729	Mailing postpartum depression prevention and treatment knowledge manual, face-to-face counseling, and telephone psychological counseling interventions, delivered during postpartum.	NA	Conventional methods	Postnatal depressive symptoms: 6 months postpartum using EPDS
Guan, 2015	Treatment RCT	Traditional Chinese medicine	60	Massage acupoint “thirteen ghost points”, once daily for 2 weeks, delivered during postpartum.	NA	Routine nursing care	Postnatal depressive symptoms: after intervention using HAMD
Liu, 2018	Prevention RCT	Traditional Chinese medicine	112	Relaxation, acupoint massage, psychological hint, 15 min per time, 3 times per day for 4 weeks, delivered during pregnancy.	NA	Routine care: antepartum health education	Prenatal depressive symptoms: after intervention using SDS
Wu, 2018	Treatment RCT	Traditional Chinese medicine	106	Diet guidance, music therapy and massage acupoint based on traditional Chinese medicine theory, delivered during postpartum.	NA	Routine nursing care	Postnatal depressive symptoms: after intervention using EPDS
Zhang, 2016	Prevention RCT	Music therapy; Relaxation therapy	100	Music therapy and relaxation therapy consisted of 30 min session, twice a day, delivered during postpartum.	Psychiatrist; Nurse	Routine care	Postnatal depressive symptoms: discharged from hospital using SDS
Liu, 2012	Prevention RCT	Exercise therapy Psychoeducation	326	Exercise under the guidance of specialist for 40 min, 3 times a week from pregnancy until delivery. Education on healthcare, nutrition, delivery and baby-caring.	Nurses; Doctors; Specialist	NA	Postnatal depressive symptoms: 7, 42 days and 3 months postpartum using EPDS
Cheng, 2016	Prevention RCT	Social support	130	Mobile application program communicating with postpartum women and providing information and emotional social support twice a week for about 10–15 min each time, delivered from 4 weeks after delivery.	Nurses	No intervention	Postnatal depressive symptoms: 1 months postpartum using EPDS

Note: NA, not available; RCT, Random controlled trial; EPDS, Edinburgh Postnatal Depression Scale; PHQ-9, Patient Health Questionnaire-9; CBT, cognitive behavior treatment; SDS, Zung Self-Rating Depression Scale; HAMD, Hamilton depression scale.



**Table 2**  
Information of intervention delivery and delivery agent ( $n = 26$ ).

Author, year	Intervention delivery Integration	Technology	Location	Format	Duration	Fidelity	Delivery agent Profession	Qualification	Training	Supervision
Gao, 2010	Integrated into hospital routine childbirth education sessions	Face-to-face Telephone	hospital	Group	6 weeks	NA	Midwives	NA	Yes	Yes
Gao, 2015	Integrated into hospital discharge education	Face-to-face Telephone	hospital	Individual	2 weeks	NA	Midwives	NA	NA	NA
Guan, 2015	Integrated into routine care.	Face-to-face	hospital	Individual	2 weeks	NA	NA	NA	NA	NA
Ho, 2009	Integrated into hospital discharge education	Face-to-face	hospital	Individual	1 day	NA	Nurses	NA	NA	NA
Huang, 2015	Integrated into routine prenatal care and education	Face-to-face	hospital	NA	6 weeks	NA	Obstetrician psychiatrist	NA	NA	NA
Jia, 2017	Integrated into routine care.	Face-to-face	hospital	Group	NA	NA	Nurse	NA	NA	NA
Jiang, 2014	Integrated into hospital visits	Mail Face-to-face Telephone	Home visits	Individual	NA	NA	NA	NA	NA	NA
Leung, 2012	Integrated into a group community-based family centered parenting program	Face-to-face	Maternal and Child Health Center	Group	4 weeks	Yes	First author, Interventionist	Yes	Yes	NA
Li, 2018	Integrated into routine care.	Face-to-face	hospital	Individual	NA	NA	Nurse	NA	NA	NA
Mao, 2012	Integrated into hospital visits	Face-to-face	hospital	Group	4 weeks	NA	Nurses	NA	Yes	NA
Wu, 2018	Integrated into routine care.	Face-to-face	hospital	Individual	NA	NA	NA	NA	NA	NA
Zhang, 2016	Integrated into routine care.	Face-to-face	hospital	Individual	NA	NA	Psychiatrist, Nurse	NA	NA	NA
Ngai, 2009	Integrated into routine childbirth education program	Face-to-face	hospital	NA	3 weeks	NA	Midwives	NA	Yes	NA
Cheng, 2016	NA	Telephone	hospital	Individual	4 weeks	NA	Nurses	NA	Yes	NA
Dou, 2018	NA	Face-to-face	Community health care center	Individual	NA	NA	NA	NA	NA	NA
Leung, 2016	NA	Face-to-face	hospital	Group	6 weeks	NA	NA	NA	NA	NA
Liu, 2012	NA	Face-to-face	hospital	Individual	NA	NA	Nurses; Doctors; Specialist	NA	NA	NA
Liu, 2018	NA	Face-to-face	hospital	Individual	NA	NA	NA	NA	NA	NA
Lu, 2016	NA	Face-to-face	hospital	Individual	8 weeks	NA	Nurses Doctors	NA	NA	NA
Lu, 2017	NA	Face-to-face	hospital	NA	6 weeks	NA	Researcher, Nurse, Psychiatrist	Yes	Yes	NA
Ngai, 2015	NA	Telephone	hospital	Individual	5 weeks	Yes	Midwives	Yes	Yes	Yes
Sun, 2011	NA	Face-to-face Telephone	hospital	Group	NA	NA	Nurse	NA	NA	NA
Wu, 2017	NA	Face-to-face Telephone	hospital	Individual	NA	NA	Doctors Nurses	NA	NA	NA
Yang, 2019	NA	Mobile APP	hospital	Individual	8 weeks	NA	Nurses midwife	NA	Yes	NA
Zhang, 2018	NA	Face-to-face	hospital	Group	8 weeks	NA	Psychologist	NA	NA	NA
Zhao, 2018	NA	Mobile APP	hospital	Individual	NA	NA	NA	NA	NA	NA


Note: NA, not available.

studies) and person-centered counseling (one study, combined counseling with education). Among cognitive strategies, the commonest elements were identifying thoughts, behaviors, and cognitive restructuring. Among behavioral theory-based interventions, the predominant elements included problem solving, relaxation techniques and emotional regulation. Interpersonal strategies included communication skills and inciting social support. Caregiver coping was the most frequently reported element among parental skills-based strategies and birth procedures were most common psychoeducational component. The therapeutic elements comprising these interventions were heterogeneous and overlapped across different interventions. Other strategies not based on any established psychological theories included traditional Chinese practices (Liu (2018): acupoint massage, education and emotion release therapy; Wu (2018): acupoint massage, music therapy and diet nursing; Guan (2015): acupoint massage, music therapy and education), social support (one study), music therapy (one study, combined exercise with relaxation therapy) and exercise (one study, combined exercise with psychoeducation). The strategies included many non-specific elements such as prompting social support, involvement of family, active listening, involvement of significant others and so on. The most commonly used traditional Chinese practice was acupoint massage. Description of principles and rationale behind these interventions has been detailed in Fig. 2 and Appendix Table A4.


The summary measure corresponded to a strong overall effect size of

0.81 (95% CI:  $-1.03$  to  $-0.58$ ,  $P < 0.001$ ), with five individual studies showing no significant difference in the effect size (Fig. 3). The overall evidence, however, presented substantial heterogeneity ( $I^2 = 91.12\%$ ,  $P < 0.001$ ,  $Q = 282.36$ ). Sensitivity analysis did not reveal any changes in the pooled effect size when individual study was removed from the meta-analysis. The effectiveness of these interventions ( $n = 10$ ) remained significant (effect size:  $-0.87$ , 95% CI:  $-1.49$  to  $-2.73$ ,  $I^2 = 97.45\%$ ) at follow-up (from 7 days to 6 months).

A series of subgroup analyses were conducted to ascertain differences in effect sizes among a priori defined groups of interventions (Appendix Table A5). High quality studies (Gao et al., 2010; Gao et al., 2015; Leung and Lam, 2012; Mao et al., 2012; Ngai et al., 2009; Ngai et al., 2015; Yang et al., 2019) yielded weaker effect sizes and performed better on heterogeneity measures than their counterparts. There was a significant difference in the effect sizes of different theoretical orientations, with exercise ( $n = 1$ ) showing the largest ( $-2.49$ , 95%CI:  $-3.26$  to  $-1.71$ ) and interpersonal therapy ( $n = 3$ ) the smallest ( $-0.22$ , 95%CI:  $-0.48$  to  $0.03$ ). Though the size effect of treatment studies was higher than that of prevention studies, no statistical difference was found ( $\chi^2 = 1.65$ ,  $p = 0.2$ ). And no difference of effect size was found between interventions delivered by specialists and non-specialists. Multivariate meta-regression was used to assess differences in effect sizes of subgroups of studies with different theoretical orientations, while accounting for quality of the studies, and employing



(a). Specific therapeutic ingredients of the included studies.



(b). Non-specific therapeutic ingredients of the included studies.

Fig. 2. (a). Specific therapeutic ingredients of the included studies. (b). Non-specific therapeutic ingredients of the included studies.

exercise-based therapy as a reference group. It yielded a significant model ( $P < 0.001$ ) that explained 69% of variation in heterogeneity in the effect size of studies, where exercise-based therapy yielded the highest effect size, followed by traditional Chinese practice, counseling, music therapy, social support, psychoeducation, cognitive behavioral therapy, mindfulness and interpersonal therapy. Details were shown in Appendix Fig. A1 & Appendix Table A6. When the analysis was restricted to psychotherapies including CBT, IPT, PE and mindfulness, no significant differences in effect sizes were noted ( $P > 0.05$ ) (Appendix Fig. A1 & Appendix Table A7). Maternal age did not account for variation in heterogeneity and effect sizes among the included trials ( $\beta = 0.06$ ;  $P = 0.35$ ).


The result demonstrated that most RCT studies (73.1%) included in the meta-analysis are of low quality. Visualization of funnel plot (Fig. 4) and assessment of Egger's regression statistic ( $P = 0.98$ ) did not reveal any significant publication bias. The highest proportion of unclear risk of bias was exhibited in blinding of participants and personnel (24 studies). While five studies were rated as having a high risk of bias in reporting of incomplete outcome data. Detailed ratings of risk of bias have been presented in Appendix Fig. A2.

#### 4. Discussion

To the best of our knowledge, this is the first systematic review to


provide an overview of psychosocial interventions, including traditional Chinese practices, for perinatal depression in China. The studies included in this review varied in the form, duration, theoretical orientation and delivery-agent of the interventions. Despite the diversity, it can be concluded that psychosocial interventions are relatively effective in reducing perinatal depressive symptoms. This result is consistent with the findings of other recent global reviews of evidence in the area (Dennis and Dowswell, 2013).

The analyses exhibited the highest effect size for exercise-based intervention, followed by psychological therapies embedded in traditional Chinese practices. This may indicate a cultural preference for interventions based on physical activity rather than conventional 'talking therapies'. However, these analyses may not present a true picture of the real treatment effects of these interventions, which may be a reflection of the quality and numbers of studies. There was only one intervention pertaining to exercise (Liu et al., 2012) and three to traditional Chinese practices (Guan et al., 2015; Liu et al., 2018; Wu, 2018). Besides, all the four above-mentioned studies were poorly designed according to the Cochrane risk of bias tool. Though classified as exercise-based intervention, the study (Liu et al., 2012) actually combined psychoeducation, which surely will exaggerate the size effect of exercise. In comparison, CBT (Mao et al., 2012; Ngai et al., 2015) and IPT based (Gao et al., 2010; Gao et al., 2015; Leung and Lam, 2012) studies showed low risk of bias and thus giving more precise estimates



**Fig. 3.** Meta-analysis of psychosocial interventions of perinatal depression in women in China

Note: \* represents no significance was found ( $p < 0.1$ ).



**Fig. 4.** Funnel plot.



of treatment effects. Furthermore, interventions classified as traditional Chinese practices also combined other non-specific elements such as support and relaxation. Therefore, it is difficult to identify the sole effectiveness of these traditional Chinese practices. However, traditional Chinese practices in the management of perinatal depression should be an area of interest. For example, acupuncture, a traditional Chinese treatment, has been shown to significantly reduce postpartum depressive symptoms (Li *et al.*, 2019). Thus, the role of traditional medicine as an adjunct to conventional approaches should be explored further and could be an important area for future research. Though classified as counseling and music therapy, Jiang (2014) and Zhang (2016) actually used multiple interventions in their studies. This may exaggerate the size effect and lead a bias to the result.

In the subgroup analyses, high quality studies yielded a weaker effect size than their low quality counterparts. A large proportion of studies included in this review were rated as low quality and tended to overestimate treatment effect sizes. Small-scale trials lacking proper randomization procedures contribute to selection bias and systematic differences between the intervention and placebo arms. Similar biases are associated with poor or no allocation concealment because allocation to intervention and control groups can be related to prognosis and responsiveness to treatment (Kunz *et al.*, 2007). Future trials should focus on removing these potential sources of biases. The subgroup analyses also demonstrated that the effect of interventions delivered by non-specialists was comparable to that by specialists, which is consistent with other literature reports (Liu *et al.*, 2017). No significant difference was found between the size effect of treatment and prevention studies. However, the result should be interpreted with caution. Participants undergoing treatment interventions have higher scores on depression screening instruments at baseline than the counterparts in preventive interventions. Thus, the former group experiences a larger decrease in their scale scores post-intervention, reflecting higher effect size estimates. In this review, several studies employed only screening instruments such as the EPDS to classify intervention recipients as having depression. The process of diagnoses should be done using diagnostic interviews based on DSM/ICD criteria for diagnosis or SCID. The use of screening instruments for diagnosis without employing a diagnostic interview can add potential bias in the meta-analysis.

It is notable that almost all studies in this review delivered their interventions in hospitals in urban areas of China. However, perinatal depression is more prevalent in underdeveloped regions and associated with poor socioeconomic and living conditions (Nisar *et al.*, 2019). This result indicated that more research should be done on how to bridge the urgent need of mental health care in rural areas and the shortage of medical human resource. It has been proven that mental health interventions delivered by non-specialist under supervision are more beneficial than routine care and can reduce the disease burden of perinatal depression (Liu *et al.*, 2017). Though half studies in this review reported integrating their interventions into existing services, few reported the process, cost and scale-up strategies. Studies in other LMICs proved that maternal mental health interventions can be added to existing health care services at little additional cost (Richter *et al.*, 2017). Based on the above evidence, task shifting from mental health specialists to the community healthcare system, which was successfully demonstrated in many studies from LMICs (Rahman *et al.*, 2013), could be a potential way forward in integration of mental health interventions and services into existing healthcare system in China.

The application of technology could be another way to improve access to mental health care. According to a recent systematic review (Ashford *et al.*, 2016), computer- or web- based interventions could be a promising approach for accessing treatment for perinatal depression. In this review, two studies (Yang *et al.*, 2019; Zhao, 2018) delivered mindfulness

interventions through a popular Chinese mobile application named We-Chat and both reported a significant positive result. However, adherence to intervention and quality control could limit the effectiveness of computer- or web-based interventions. Also, given that perinatal depression is more prevalent in under-developed regions, technological solutions must ensure equity for those who do not have access to this technology.

## 5. Limitation

There are some limitations in this review. Firstly, the evidence in this review is of moderate to low quality and therefore, should be interpreted with caution. The majority of the trials had high risk of bias with poor randomization and allocation concealment procedures. Secondly, some of the trials were inadequately powered and tended to overestimate effect sizes as demonstrated by a wide confidence interval for standard mean differences for those ten follow-up data. Thirdly, the implications of this review is limited in China as we only include psychosocial interventions on perinatal depression in China. Lastly, we did not find any statistical heterogeneity based on different assessment tools for assessment of depressive symptoms among the intervention recipients in different studies. Nonetheless, use of different psychometric instruments with different psychometric properties introduces methodological heterogeneity in meta-analysis. Therefore, these results should be interpreted with caution.

## 6. Conclusion

This review indicates that psychosocial interventions, particularly those involving psychoeducational and cognitive behavioral elements, are effective in reducing perinatal depressive symptoms. Traditional Chinese practices and exercise may add to the effectiveness of interventions. However, future high quality RCTs are required to supplement the evidence. Concurrently, strategies to scale-up effective interventions should be further developed and evaluated in a range of settings to reduce the treatment gap for perinatal depression.

## Contributors

Juan Yin, Atif Rahman and Xiaomei Li conceived and designed the study; Anum Nisar and Juan Yin searched, screened and did the data extraction of the English studies; Yan Guo and Wenli Qi searched, screened and did the data extraction of the Chinese studies; Ahmed Waqas and Duolao Wang did the meta-analysis; Juan Yin first draft the paper and all the authors revised and approved it.

## Funding

This study was supported by the Chinese Nursing Association (ZHKY201809). The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

## Declaration of Competing Interest

None.

## Acknowledgement

We would like to thank the Chinese Nursing Association for supporting our research.

Appendix

Figs. A1 and A2.






Fig. A1. Meta-regression analysis for theoretical orientation.



(a) Review authors' judgements about each risk of bias item presented as percentages across all included studies.



(b) Risk of bias items for each included study.

Fig. A2. Risk of bias.

## Tables A1,A2,A3,A4,A5,A6,A7,A8,A9,A10.

**Table A1**

Search strategy.

Search strategy in English databases	Search strategy in Chinese databases
(Perinatal[ti/ab] OR maternal[ti/ab] OR mother*[ti/ab] OR pregnancy[ti/ab] OR pregnant[ti/ab] OR intrapartum[ti/ab] OR postpartum[ti/ab] OR prenatal[ti/ab] OR postnatal[ti/ab] OR antenatal[ti/ab] OR antepartum[ti/ab]) AND (depression[ti/ab] OR depressive[ti/ab] OR depress*[ti/ab] OR "mental health"[ti/ab]) AND (China OR Chinese OR Guangxi OR "inner Mongolia" OR Ningxia OR Tibet OR Xizang OR Xinjiang OR Heilongjiang OR Jilin OR Liaoning OR Hebei OR Shanxi OR Shandong OR Shaanxi OR Gansu OR Qinghai OR Sichuan OR Hubei OR Hunan OR Henan OR Anhui OR Zhejiang OR Jiangsu OR Guangdong OR Jiangxi OR Fujian OR Guizhou OR Yunnan OR Hainan OR Han OR Hui OR Mongolia OR Zhuang) AND (trial[ti/ab] OR RCT[ti/ab] OR randomized-controlled[ti/ab] OR intervention[ti/ab] OR controlled-trial[ti/ab] OR effectiveness[ti/ab] OR efficacy[ti/ab])	1. China National Knowledge Infrastructure (CNKI) (TI=围产期 OR TI=产妇 OR TI=妊娠 OR TI=孕妇 OR TI=孕期 OR TI=产前 OR TI=产后 OR TI=分娩前 OR TI=分娩后 OR TI=产褥期) AND (TI=抑郁 OR TI=精神 OR TI=心理) AND (AB=试验 OR AB=RCT OR AB=随机对照 OR AB=干预 OR AB=对照 OR AB=效果) 2. China National Knowledge Infrastructure (CNKI) (TI=围产期 OR TI=产妇 OR TI=妊娠 OR TI=孕妇 OR TI=孕期 OR TI=产前 OR TI=产后 OR TI=分娩前 OR TI=分娩后 OR TI=产褥期) AND (TI=抑郁 OR TI=精神 OR TI=心理) AND (AB=试验 OR AB=RCT OR AB=随机对照 OR AB=干预 OR AB=对照 OR AB=效果) 3. VIP Database for Chinese Technical Periodicals (M=围产期 OR M=产妇 OR M=妊娠 OR M=孕妇 OR M=孕期 OR M=产前 OR M=产后 OR M=分娩前 OR M=分娩后 OR M=产褥期) AND (M=抑郁 OR M=精神 OR M=心理) AND (M=试验 OR M=RCT OR M=随机对照 OR M=干预 OR M=对照 OR M=效果) 4. Wan Fang Database Title or Key words:((围产期 OR 产妇 OR 妊娠 OR 孕妇 OR 孕期 OR 产前 OR 产后 OR 分娩前 OR 分娩后 OR 产褥期) AND (抑郁 OR 精神 OR 心理) AND (试验 OR RCT OR 随机对照 OR 干预 OR 对照 OR 效果))

**Table A2**

Definitions of therapeutic techniques employed in psychosocial interventions.

Name of Skill	Definition of skill
1. Involvement of family	The family members are involved in the intervention.
2. Involvement of significant other	The significant other or spouse is involved in the intervention.
3. Active listening	The skillful listens to the speaker with full concentration to understand what is being said [1].
4. Collaboration	Working with others
5. Inciting social support	Providing insight of how others can provide help or support [2].
6. Case management	Planning, facilitating and coordination of different options regarding the patient [3].
7. Normalization	To communicate that the person's experiences also happen to other people [4].
8. eliciting commitment	Motivating the client to take active part in intervention.
9. Discussing advantages	Identifying advantages to take the intervention.
10. Discussing barriers	Identifying difficulties to take the intervention.
11. Identifying affect	Identifying feeling or emotion [5].
12. identifying and eliciting social support	Providing consideration of how others could change their behavior to offer the person help or support [6].
13. Communication skills	These are set of skills which improve the dissemination, reception and exchange of information, opinions or ideas making sure that the intended message is completely understood by those involved [7].
14. Assertiveness training	To train people for effective communication without being passive or aggressive.
15. Assessing relationships	To assess the relationships with people around.
16. Problem solving	To solve a problem by carefully defining problem and weighing different options to solve the problem.
17. Relaxation	To apply various techniques for relaxation.
18. Awarding positive behavior	Applying different methods to encourage recurrence of a positive behavior [8].
19. Exposure	Confronting previously avoided objects, situations, unwanted thoughts or feelings while not avoiding or escaping from it [9].
20. Emotional regulation	Tendency to manage and responding to emotional experience [10].
21. Stress management	May involve a variety of techniques that do not target a specific behavior but seek to reduce anxiety and stress [6].
22. Decision making	Process of making choices by identifying decision, gathering information, and assessing alternative resolutions [11].
23. self-monitoring	The person keeps a record of one's own behavior [6].
24. Delay awards	A process of deferring initial reward for a better reward later [12].
25. Caregiver coping (e.g., management skills for the parents of children)	Mechanism through which the caregiver manages the stress [13].
26. Parent-child Interaction Coaching (e.g., positive 1-on-1, attending to children modeling, etc)	The parent applies a new skill with the child and the therapist provides immediate feedback [14].
27. Birth procedures	Knowledge about different ways of giving birth [15].
28. Specific health areas of children (e.g., nutrition, breastfeeding, SRH, etc)	If the intervention targeted the specific dimensions of health areas in children.
29. Nutrition	Education regarding recommended nutrition practices.
30. Breastfeeding	Education regarding recommended breastfeeding practices.
31. Sexual behavior	Avoidance of risky sexual behaviors.
32. Identifying thoughts, behaviors, and their links (e.g., identifying negative thoughts, thought diary, etc.)	Realizing what thoughts cross one's mind by number of ways [16].
33. Cognitive restructuring (e.g., reattribution, weighing evidence, logical questioning, etc.)	Identifying and disputing maladaptive thoughts [2].
34. Distraction	Paying attention to some other stimuli rather than the unhelpful thoughts [17].
35. Self-talk	Use of self-instruction and self-encouragement to support action [6].
36. Self-praise	To boast one's self esteem by expressing approval or admiration [18].
37. Mood monitoring	Paying attention toward one's mood states by means of different methods [19].
38. Mindfulness	Paying attention towards experiences in the present moment within body and mind and accepting the happenings [20].
39. Self-awareness	Conscious awareness of becoming the object of one's own awareness [21].
40. Aerobics	Aerobic exercise also known as "cardio" exercises include running, swimming, walking, hiking, aerobics classes, dancing, cross country skiing, and kickboxing.

(continued on next page)

Table A2 (continued)

Name of Skill	Definition of skill
41. Non-aerobic exercise	Anaerobic ("without oxygen") exercise is any physical activity that causes you to be quickly out of breath, like sprinting or lifting a heavy weight.
42. Motivational enhancement	To increase internal motivation in order to make long lasting change [22].
43. Praise	Praising for positive behavior or accomplishment.
44. Role play	Performing role of a person in a situation [23].
45. Behavioral contracting	Agreement of a contract specifying behavior to be performed so that there is a written record of the person's resolution witnessed by another [6].
46. Assigning homework	Assigning tasks pertaining to interventions, to be performed at home.
47. Interpersonal focus	Focus on maintaining relationships with other people [24].
48. Behavioral experiments	To test out the negative thoughts and re-evaluate underlying beliefs by performing an action [25].
49. Motivational interviewing	Prompting the person to provide self-motivating statements and evaluation of their own behavior to minimize resistance [6].
50. Direct suggestions	Providing directions of how to act, behave or handle a situation [26, 27].
51. Goal setting	Identifying an aim or goal to achieve in a session or therapy [28].
52. Giving sick role	Assigning role of a sick person to the participant, to understand the circumstances, particular rights and responsibilities of those who are ill.
53. Empathy	To understand and share other's feeling or situation [29].
54. Macronutrients	Food groups needed in large amounts [30].
55. Micronutrient	Food groups needed in small amounts [30].
56. Eating behaviors	The food choices and eating practices [31].
57. Alcohol use	Education about minimal use and harms of alcohol misuse.
58. Substance misuse	Harmful use of substances for non-medical purposes [32].

## References

- Rogers, C.R. and R.E. Farson, *Active listening*. 1957: Industrial Relations Center of the University of Chicago.
- Chambless, D.L. and S.D. Hollon, *Defining empirically supported therapies*. Journal of consulting and clinical psychology, 1998. 66(1): p. 7.
- Marshall, M., et al., *Case management for people with severe mental disorders*. The Cochrane database of systematic reviews, 2000(2): p. CD000050-CD000050.
- Dovis, S., et al., *Can motivation normalize working memory and task persistence in children with attention-deficit/hyperactivity disorder? The effects of money and computer-gaming*. Journal of abnormal child psychology, 2012. 40(5): p. 669-681.
- Dube, S., *Enchantments of modernity: empire, nation, globalization*. 2012: Routledge.
- Abraham, C. and S. Michie, *A taxonomy of behaviour change techniques used in interventions*. Health psychology, 2008. 27(3): p. 379.
- Baran, S.J., E. McDonald, and J. Engberg, *Introduction to mass communication: Media literacy and culture*. 2004: McGraw-Hill.
- Cregor, M., *The building blocks of positive behaviour. teaching tolerance*, 2008. 34(4): p. 18-21.
- Forsyth, J.P., V. Barrios, and D.T. Acheson, *Exposure therapy and cognitive interventions for the anxiety disorders: overview and newer third-generation perspectives*, in *Handbook of exposure therapies*. 2007, Elsevier. p. 61-108.
- Rolston, A. and E. Lloyd-Richardson, *What is emotion regulation and how do we do it?* Cornell Research Program on Self-Injury and Recovery.[on-line].[cit. 2017-22-09] Available at: <http://www.selfinjury.bctr.cornell.edu/perch/resources/what-is-emotion-regulationsinfo-brief.pdf>. Google Scholar, 2017.
- Janis, I.L. and L. Mann, *Decision making: A psychological analysis of conflict, choice, and commitment*. 1977: free press.
- Metcalfe, J. and W. Mischel, *A hot/cool-system analysis of delay of gratification: dynamics of willpower*. Psychological review, 1999. 106(1): p. 3.
- Zeiss, A.M., et al., *Self-efficacy as a mediator of caregiver coping: Development and testing of an assessment model*. Journal of clinical Geropsychology, 1999. 5(3): p. 221-230.
- McNeil, C.B. and T.L. Hembree-Kigin, *Parent-child interaction therapy*. 2010: Springer Science & Business Media.
- Thompson, R. and Y.D. Miller, *Birth control: to what extent do women report being informed and involved in decisions about pregnancy and birth procedures?* BMC pregnancy and childbirth, 2014. 14(1): p. 62.
- Longmore, R.J. and M. Worrell, *Do we need to challenge thoughts in cognitive behaviour therapy?* Clinical psychology review, 2007. 27(2): p. 173-187.
- Fennell, M.J. and J.D. Teasdale, *Effects of distraction on thinking and affect in depressed patients*. British Journal of Clinical Psychology, 1984. 23(1): p. 65-66.
- Abdel-Khalek, A.M., *INTRODUCTION TO THE PSYCHOLOGY OF SELF-ESTEEM*. SELF-ESTEEM: p. 1.
- Swinkels, A. and T.A. Giuliano, *The measurement and conceptualization of mood awareness: Monitoring and labeling one's mood states*. Personality and social psychology bulletin, 1995. 21(9): p. 934-949.
- Langer, E.J., *Mindfulness*. 1989: Addison-Wesley/Addison Wesley Longman. 88
- Morin, A., *Self-awareness part 1: Definition, measures, effects, functions, and antecedents*. Social and personality psychology compass, 2011. 5(10): p. 807-823.
- Miller, W.R., *Motivational enhancement therapy manual: A clinical research guide for therapists treating individuals with alcohol abuse and dependence*. 1995: DIANE Publishing.
- Van Ments, M., *The effective use of role-play: Practical techniques for improving learning*. 1999: Kogan Page Publishers.
- Leary, M.R. and R.F. Baumeister, *The need to belong: Desire for interpersonal attachments as a fundamental human motivation*, in *Interpersonal Development*. 2017, Routledge. p. 57-89.
- Rouf, K., et al., *Devising effective behavioural experiments*. Oxford guide to behavioural experiments in cognitive therapy, 2004: p. 21-58.
- Rossi, E.L. and K.L. Rossi, *What is a suggestion? The neuroscience of implicit processing heuristics in therapeutic hypnosis and psychotherapy*. American Journal of Clinical Hypnosis, 2007. 49(4): p. 267-281.
- Podikunju-Hussain, S., *Working with Muslims: Perspectives and suggestions for counseling*. Vistas: Compelling perspectives on counseling, 2006: p. 103-106.
- Locke, E.A. and G.P. Latham, *A theory of goal setting & task performance*. 1990: Prentice-Hall, Inc.
- Chinsky, J.M. and J. Rappaport, *Brief critique of the meaning and reliability of "accurate empathy" ratings*. Psychological Bulletin, 1970. 73(5): p. 379.
- Hadigan, C.M., et al., *Assessment of macronutrient and micronutrient intake in women with anorexia nervosa*. International Journal of Eating Disorders, 2000. 28(3): p. 284-292.
- Turconi, G., et al., *Eating habits and behaviours, physical activity, nutritional and food safety knowledge and beliefs in an adolescent Italian population*. Journal of the American College of Nutrition, 2008. 27(1): p. 31-43.
- Lingford-Hughes, A., S. Welch, and D. Nutt, *Evidence-based guidelines for the pharmacological management of substance misuse, addiction and comorbidity: recommendations from the British Association for Psychopharmacology*. Journal of Psychopharmacology, 2004. 18(3): p. 293-335.

**Table A3**  
Demographic characteristics of participants in the researches included.

Author, year	Age of mothers	Mean age (SD) (years)	Principle of intervention	Geographical scope	Study Settings	Outcome	Outcome measures
Gao, 2010	28.47 (2.8)		Prevention	NA	hospital	Postnatal depression	EPDS
Gao, 2015	28.49 (2.7)		Prevention	NA	hospital	Postnatal depression	EPDS
Ho, 2009	intervention group: 29.2 (3.4) control group: 29.4 (3.2)		Prevention	urban	hospital	Postnatal depression	EPDS
Jiang, 2014	27.9 (4.36)		Treatment	urban	Home visits	Postnatal depression	EPDS
Mao, 2012	intervention group: 28.5 (2.4) control group: 28.8 (2.5)		Prevention	urban	hospital	Postnatal depression	EPDS, PHQ-9
Cheng, 2016	intervention group: 33.48 (4.12) control group: 32.82 (4.93)		Prevention	urban	hospital	Postnatal depression	EPDS
Huang, 2015	intervention group: 28.5 (2.4) control group: 28.8 (2.5)		Prevention	urban	hospital	Postnatal depression	PHQ-9
Leung, 2016	21 to 45		Treatment	urban	hospital	Postnatal depression	EPDS
Leung, 2012	intervention group: 31.3 (4.02) control group: 31.15 (4.12)		Treatment	urban	Maternal and Child Health Centers	Prenatal and postnatal depression	EPDS
Ngai, 2009	intervention group: 32.1 (3.7) control group: 30.5 (3.7)		Prevention	urban	hospital	Prenatal and postnatal depression	EPDS
Ngai, 2015	intervention group: 31.1 (3.8) control group: 30.4 (4.4)		Treatment	urban	hospital	Postnatal depression	EPDS
Yang, 2019	intervention group: 31.31(4.97) control group: 30.38 (3.91)		Prevention	urban	hospital	Prenatal depression	PHQ-9
Zhang, 2018	intervention group: 25.7 (2.79) control group: 25.58 (2.33)		Prevention	urban	hospital	Prenatal depression	SDS
Lu, 2016	20–38		Treatment	urban	hospital	Postnatal depression	EPDS
Sun, 2011	intervention group: 28.38 (2.73) control group: 28.55 (2.82)		Prevention	urban	hospital	Postnatal depression	EPDS
Liu, 2018	20–44		Prevention	urban	hospital	Prenatal depression	SDS
Wu, 2017	21–39		Treatment	urban	hospital	Postnatal depression	SDS
Li, 2018	22–38		Treatment	urban	hospital	Postnatal depression	HAMD
Jia, 2017	intervention group: 27.62 (3.15) control group: 27.94 (3.28)		Treatment	urban	hospital	Postnatal depression	EPDS
Dou, 2018	intervention group: 28.81 (4.25) control group: 29.04 (4.61)		Prevention	semi-urban	Community health care center	Postnatal depression	SDS
Zhao, 2018	20–42		Treatment	urban	hospital	Postnatal depression	SDS
Lu, 2017	intervention group: 28.5 (4.7) control group: 28.9 (4.6)		Treatment	urban	hospital	Postnatal depression	HAMD;
Zhang, 2016	24–39		Prevention	urban	hospital	Postnatal depression	SDS
Guan, 2015	intervention group: 28.8 (15.8) control group: 29.6 (16.1)		Treatment	semi-urban	hospital	Postnatal depression	SDS
Wu, 2018	20–40		Treatment	urban	hospital	Postnatal depression	HAMD
Liu, 2012	intervention group: 28 (5) control group: 27 (6)		Prevention	urban	hospital	Postnatal depression	EPDS

Note: NA, not available; EPDS, Edinburgh Postnatal Depression Scale; PHQ-9, Patient Health Questionnaire-9; SDS, Zung Self-Rating Depression Scale; HAMD, Hamilton depression scale.

**Table A4**  
Scope and taxonomy of interventions.

Author year	Type	Major elements	Elements
Gao, 2010	Prevention	Engagement Interpersonal skills Behavior coping Parenting skill Psychoeducation Techniques used by delivery agent	Active listening, inciting social support Identifying and eliciting social support, communication skill, assessing relationships Problem solving, stress management Caregiver coping, parent-child interaction coaching Birth procedure, nutrition, breastfeeding, sexual behavior Motivational enhancement, role play, interpersonal focus, direct suggestion, goal setting
Gao, 2015	Prevention	Engagement Interpersonal skill Behavior coping Parenting skill Psychoeducation Techniques used by delivery agent	Active listening, inciting social support Identifying affect, identifying and eliciting social support, Communication skill, assessing relationships Problem solving Parent-child interaction coaching Breastfeeding Interpersonal focus
Ho, 2009	Prevention	Cognitive coping	Mood monitoring, self-awareness
Mao, 2012	Prevention	Engagement Interpersonal skill Behavior coping Psychoeducation Cognitive coping Techniques used by delivery agent	Involvement of significant others, inciting social support Communication skill Problem solving, relaxation, self-monitoring Birth procedure Identifying thoughts, cognitive restructuring, self-talk, self-praise, mood monitoring, self-awareness Role paly, assigning homework, interpersonal focus, reviewing homework
Jiang, 2014	Treatment	–	–
Cheng, 2016	Prevention	Engagement Interpersonal skills Psychoeducation Techniques used by delivery agent	Empathy Identifying and eliciting social support Birth procedure, breast feeding Empathy
Huang, 2015	Prevention	Engagement Interpersonal skills Behavior coping Psychoeducation Cognitive coping	Involvement of family, inciting social support Identifying and eliciting social support Relaxation, exposure, emotional regulation, eliciting social support Birth procedure Cognitive restructuring
Leung, 2016	Treatment	Cognitive coping techniques used by delivery agent	Identifying thoughts, cognitive restructuring, distraction Assigning homework
Leung, 2012	Treatment	Interpersonal skills techniques used by delivery agent	Communication skills, assertiveness training, assessing relationship Role play, assigning homework, interpersonal focus, reviewing homework, goal setting
Ngai, 2009	Prevention	Behavior coping Reinforcement Parenting skill Cognitive coping Techniques used by delivery agent	Problem solving, stress management, decision making Awarding positive behavior Caregiver coping Cognitive restructuring Goal setting
Ngai, 2015	Treatment	Interpersonal skills Behavior coping Reinforcement Parenting skill Cognitive coping Techniques used by delivery agent	Communication skills Problem solving, exposure, stress management, decision making Awarding positive behavior Caregiver coping Identifying thoughts, cognitive restructuring Goal setting
Yang, 2019	Prevention	Engagement Interpersonal skills Behavior coping Cognitive coping Techniques used by delivery agent	Discussing barriers to treatment Identifying affect Relaxation, stress management Identifying thoughts, mood monitoring, mindfulness Assigning homework, reviewing homework
Zhang, 2018	Prevention	Engagement Cognitive coping Exercise Techniques used by delivery agent	Discussing barriers to treatment Mindfulness Aerobics Assigning homework, reviewing homework
Lu, 2016	Treatment	Engagement Exercise Techniques used by delivery agent	Involvement of family, active listening, empathy Others Empathy
Sun, 2011	Prevention	Engagement Interpersonal skills Behavior coping Parenting skill	Inciting social support Identifying and eliciting social support, communication skills Stress management Caregiver coping
Liu, 2018	Prevention	–	–
Wu, 2017	Treatment	Parenting skill Psychoeducation	Caregiver coping Birth procedure, breast feeding
Li, 2018	Treatment	Engagement Cognitive coping Techniques used by delivery agent	Active listening, empathy Identifying thoughts, cognitive restructuring Assigning homework, empathy
Jia, 2017	Treatment	Engagement Behavior coping Psychoeducation Cognitive coping	Involvement of family, involvement of significant others Emotional regulation Birth procedure Identifying thoughts
Dou, 2018	Prevention	Engagement Behavior coping Cognitive coping	Inciting social support Relaxation Identifying thoughts
Zhao, 2018	Treatment	Engagement	Involvement of family

(continued on next page)



Table A4 (continued)

Author year	Type	Major elements	Elements
Lu, 2017	Treatment	Behavior coping	Emotional regulation
		Psychoeducation	Nutrition
		Engagement	Empathy, normalization, eliciting commitment
		Cognitive coping	Identifying thoughts, mindfulness
Zhang, 2016	Prevention	Techniques used by delivery agent	Empathy
Guan, 2015	Treatment	Behavior coping	Relaxation
Wu, 2018	Treatment	–	–
Liu, 2012	Prevention	Psychoeducation	Nutrition
		Engagement	Involvement of family
		Psychoeducation	Birth procedure, nutrition
		Exercise	Aerobics

Table A5

Subgroup analysis of the psychosocial interventions.

Moderators	No• of studies	Point estimate (95% CI)	I <sup>2</sup>	χ <sup>2</sup> (Q statistic)	P
Quality					
High	7	-0.45 (-0.85 to -0.05)	61.83%	4.13	0.04
Low	19	-0.94 (-1.19 to -0.69)	91.78%		
Scale					
EPDS	14	-0.83 (-1.19 to -0.48)	94.67%	1.32	0.73
SDS	6	-0.66 (-0.93 to -0.38)	60.28%		
HAMD	3	-1.13 (-1.84 to -0.41)	84.67%	29.17	<0.001
PHQ-9	3	-0.68 (-0.87 to -0.48)	23.81%		
Theoretical orientation					
CBT	7	-0.72 (-1.00 to -0.43)	81.30%	0.11	0.74
Psychoeducation	7	-0.70 (-0.98 to -0.42)	73.72%		
TCM	3	-1.24 (-2.05 to -0.40)	89.38%	1.65	0.20
Interpersonal treatment	3	-0.22 (-0.48 to 0.03)	31.46%		
Mindfulness	2	-0.53 (-1.37 to 0.31)	86.43%	0.78	0.38
Counseling	1	-1.06 (-1.79 to -0.32)	0%		
Exercise	1	-2.49 (-3.26 to -1.71)	0%	3.78	0.05
Social support	1	-0.82 (-1.62 to -0.02)	0%		
Music Therapy	1	-0.90 (-1.73 to -0.08)	0%	2.47	0.12
Specificity					
Non-specificity	7	-0.87 (-1.14 to -0.60)	74.66%	0.77	0.39
specificity	19	-0.78 (-1.08 to -0.49)	92.85%		
Scope of intervention					
Treatment	12	-0.97(-1.31 to -0.63)	92.97%	0.10	0.76
Prevention	14	-0.67(-0.98 to -0.36)	87.59%		
Elements of interventions					
Engagement					
No	11	-0.69 (-0.98 to -0.40)	88.89%	1.05	0.31
Yes	15	-0.90 (-0.99 to -0.55)	92.74%		
Interpersonal skills					
No	17	-0.96 (-1.29 to -0.64)	92.65%	1.44	0.23
Yes	9	-0.53 (-0.69 to -0.37)	57.52%		
Behavioral coping					
No	14	-0.97 (-1.37 to -0.57)	94.26%	0.41	0.52
Yes	12	-0.60 (-0.71 to -0.48)	31.24%		
Reinforcement					
No	24	-0.84 (-1.08 to -0.59)	91.43%	0.10	0.76
Yes	2	-0.49 (-0.67 to -0.31)	0%		
Parenting skills					
No	19	-0.83 (-1.04 to -0.61)	85.23%	1.05	0.31
Yes	7	-0.75 (-1.35 to -0.15)	96.26%		
Psychoeducation					
No	16	-0.71 (-0.93 to -0.49)	85.02%	1.44	0.23
Yes	10	-0.96 (-1.44 to -0.47)	94.73%		
Cognitive coping					
No	14	-0.93 (-1.30 to -0.57)	93.61%	0.41	0.52
Yes	12	-0.65 (-0.86 to -0.44)	76.85%		
Delivery agents					
Non-specialist*	15	-0.89(-1.23 to -.56)	93.26%	0.41	0.52
Specialist	5	-0.67(-1.26 to -0.08)	90.45%		

Note: EPDS, Edinburgh Postnatal Depression Scale; HAMD, Hamilton depression scale; PHQ-9, Patient Health Questionnaire-9; SDS, Zung Self-Rating Depression Scale; CBT, Cognitive behavior treatment; TCM, Traditional Chinese medicine.

\* includes multidisciplinary teams with no specialists

^ includes multidisciplinary teams with at least one specialist

**Table A6**  
Meta-regression analyses for theoretical orientation of psychosocial therapies.

Covariate	Coefficient	S-E	95% CI		Z-value	p
			Lower	Upper		
Intercept	-2.37	0.39	-3.13	-1.60	-6.05	0
Quality	-0.12	0.19	-0.49	0.26	-0.62	0.53
CBT	1.74	0.37	1.01	2.47	4.67	<0.001
Counseling	1.43	0.47	0.51	2.34	3.06	<0.001
IPT	2.17	0.42	1.35	3.00	5.14	<0.001
Mindfulness	1.87	0.45	0.10	2.74	4.19	<0.01
Music therapy	1.58	0.51	0.59	2.57	3.12	<0.01
Psychoeducation	1.77	0.37	1.04	2.49	4.79	<0.001
Social support	1.66	0.49	0.69	2.63	3.36	<0.001
TCM	1.24	0.41	0.44	2.04	3.04	<0.01

Q = 31.20, df = 8, p = 0.0001;  $R^2$  = 69%; n = 26 studies

**Table A7**  
Meta-regression analyses for psychotherapies only.

Covariate	Beta	Standard Error	95% CI		Z-value	2-sided P-value
			Lower	Upper		
Intercept	-0.71	0.14	-0.99	-0.44	-5.11	0
Interpersonal therapy	0.49	0.26	-0.02	0.99	1.88	0.06
Mindfulness	0.16	0.31	-0.44	0.77	0.53	0.60
Psychoeducation	0.01	0.20	-0.38	0.41	0.07	0.95

Q = 4.08, df = 3, p = 0.2529;  $R^2$  = 24%

**Table A8**  
Inter-coder agreement of the taxonomy of interventions.

	N	Range	Minimum	Maximum	Mean	Std. Deviation
Elements	61	0.35	0.65	1.00	0.9924	0.04706
Valid N (listwise)	61					

**Table A9**  
Inter-coder agreement of risk of bias.

	N	Range	Minimum	Maximum	Mean	Std. Deviation
bias	7	0.16	0.84	1.00	0.9504	0.06591
Valid N (listwise)	7					

**Table A10**  
PRISMA 2009 check list.

Section/topic	#	Checklist item
<i>Title</i>		
Title	1	Identify the report as a systematic review, meta-analysis, or both.
<i>Abstract</i>		
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.
<i>Introduction</i>		
Rationale	3	Describe the rationale for the review in the context of what is already known.
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).
<i>Methods</i>		
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).

(continued on next page)

Table A10 (continued)

Section/topic	#	Checklist item
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis.
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.
<b>Results</b>		
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).
<b>Discussion</b>		
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.
<b>Funding</b>		
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.

## Reference

- Ashford, M.T., Olander, E.K., Ayers, S., 2016. Computer- or web-based interventions for perinatal mental health: a systematic review. *J. Affect. Disord.* 197, 134–146.
- Bauer, A., Knapp, M., Parsonage, M., 2016. Lifetime costs of perinatal anxiety and depression. *J. Affect. Disord.* 192, 83–90.
- Biaggi, A., Conroy, S., Pawlby, S., Pariante, C.M., 2016. Identifying the women at risk of antenatal anxiety and depression: a systematic review. *J. Affect. Disord.* 191, 62–77.
- Cheng, H.Y., Huang, T.Y., Chien, L.Y., Cheng, Y.F., Chen, F.J., 2016. The effects of a mobile application social support program on postpartum perceived stress and depression. *J. Nurs. (Chinese)* 63, 52–60.
- Dennis, C.L., Dowswell, T., 2013. Psychosocial and psychological interventions for preventing postpartum depression. The Cochrane database of systematic reviews, CD001134.
- Dou, J., 2018. Preventive effect of perioperative psychological guidance on postpartum depression. *Psychol. Doc. (Chinese)* 24, 81–82.
- Evans, J., Heron, J., Francomb, H., Oke, S., Golding, J., 2001. Cohort study of depressed mood during pregnancy and after childbirth. *BMJ* 323, 257–260.
- Gao, L.L., Chan, S.W., Li, X., Chen, S., Hao, Y., 2010. Evaluation of an interpersonal-psychotherapy-oriented childbirth education programme for Chinese first-time childbearing women: a randomised controlled trial. *Int. J. Nurs. Stud.* 47, 1208–1216.
- Gao, L.L., Xie, W., Yang, X., Chan, S.W., 2015. Effects of an interpersonal-psychotherapy-oriented postnatal programme for Chinese first-time mothers: a randomized controlled trial. *Int. J. Nurs. Stud.* 52, 22–29.
- Gelaye, B., Rondon, M.B., Araya, R., Williams, M.A., 2016. Epidemiology of maternal depression, risk factors, and child outcomes in low-income and middle-income countries. *Lancet Psychiatry* 3, 973–982.
- Guan, L., Sun, J., Zheng, H., Pan, N., Shi, H., 2015. Evaluation on the effect of seven emotion theory of traditional Chinese medicine in nursing care of patients with postpartum depression. *J. Sichuan Tradit. Chin. Med. (Chinese)* 33, 172–175.
- American College of Obstetricians and Gynecologists, 2018. Screening for perinatal depression. Committee opinion no. 757. *Obstet Gynecol.* 132, 1314–1316.
- Herba, C.M., Glover, V., Ramchandani, P.G., Rondon, M.B., 2016. Maternal depression and mental health in early childhood: an examination of underlying mechanisms in low-income and middle-income countries. *Lancet Psychiatry* 3, 983–992.
- Ho, S.M., Heh, S.S., Jevitt, C.M., Huang, L.H., Fu, Y.Y., Wang, L.L., 2009. Effectiveness of a discharge education program in reducing the severity of postpartum depression: a randomized controlled evaluation study. *Patient Educ. Couns.* 77, 68–71.
- Huang, J., Li, H., Wang, J., Mao, H., Jiang, W., Zhou, H., Chen, S., 2015. Prenatal emotion management improves obstetric outcomes: a randomized control study. *Int. J. Clin. Exp. Med.* 8, 9667–9675.
- Jia, D., Liang, Y., Tang, C., 2017. Effect of group positive experience on preventing postpartum depression. *Chin. J. Mod. Nurs. (Chinese)* 23, 1201–1204.
- Jiang, L., Wang, Z.Z., Qiu, L.R., Wan, G.B., Lin, Y., Wei, Z., 2014. Psychological intervention for postpartum depression. *J. Huazhong Univ. Sci. Technol. Med. Sci. (Chinese)* 34, 437–442.
- Kunz, R., Vist, G., Oxman, A.D., 2007. Randomisation to protect against selection bias in healthcare trials. The Cochrane database of systematic reviews, Mr000012.
- Leung, S., Lee, A., Wong, D., Wong, C., Leung, K., Chiang, V., Yung, W., Chan, S., Chung, K., 2016. A brief group intervention using a cognitive-behavioral approach to reduce postnatal depressive symptoms: a randomised controlled trial. *Hong Kong Med. J. (Chinese)* 22, S4–S8.
- Leung, S.S., Lam, T.H., 2012. Group antenatal intervention to reduce perinatal stress and depressive symptoms related to intergenerational conflicts: a randomized controlled trial. *Int. J. Nurs. Stud.* 49, 1391–1402.
- Li, A., 2018. Effect of empathy nursing combined with cognitive behavior intervention on HAMD score and quality of life in patients with postpartum depression. *Henan Med. Res. (Chinese)* 27, 1326–1327.
- Li, W., Yin, P., Lao, L., Xu, S., 2019. Effectiveness of Acupuncture Used for the Management of Postpartum Depression: A Systematic Review and Meta-Analysis. *Biomed. Res. Int.* 2019, 6597503.
- Liu, H., Sun, G., Liu, Y., Fu, J., 2012. Observation on the clinical effect of health exercises and health education during pregnancy on the prevention of postpartum depression. *Matern. Child Health Care China (Chinese)* 27, 5851–5852.
- Liu, S., Yan, Y., Gao, X., Xiang, S., Sha, T., Zeng, G., He, Q., 2017. Risk factors for postpartum depression among Chinese women: path model analysis. *BMC Pregnancy Childbirth* 17, 133.
- Liu, Z., Na, L., Deng, Z., Shao, X., 2018. Effect of emotional release therapy in prenatal care on mental health, self-efficacy and parenting cognition of pregnant and puerpera. *Nurs. Pract. Res. (Chinese)* 15, 105–107.
- Lu, D., Fan, J., 2017. Effect of routine nursing combined with Acceptance and Commitment Therapy on the depression mood, self efficacy and self-esteem of patients with postpartum depression. *Sichuan Ment. Health (Chinese)* 30, 253–255.
- Lu, X., Chen, W., Wu, H., 2016. Effect of Roy adaptation model implementation of psychological comfort care in patients with postpartum depression. *China Med. Herald (Chinese)* 13, 168–171.
- Mao, H.J., Li, H.J., Chiu, H., Chan, W.C., Chen, S.L., 2012. Effectiveness of antenatal emotional self-management training program in prevention of postnatal depression in Chinese women. *Perspect. Psychiatr. Care* 48, 218–224.
- Ministry of Foreign Affairs, P. R. C., 2015. Report on China's Implementation of the Millennium Development Goals (2000–2015).
- Naveed, S., Waqas, A., Amray, A.N., et al., 2019. Implementation and effectiveness of non-specialist mediated interventions for children with Autism spectrum disorder: a systematic review and metaanalysis. *PLoS One* 14 (11), e0224362.
- Ngai, F.W., Chan, S.W., Ip, W.Y., 2009. The effects of a childbirth psychoeducation program on learned resourcefulness, maternal role competence and perinatal depression: a quasi-experiment. *Int. J. Nurs. Stud.* 46, 1298–1306.
- Ngai, F.W., Wong, P.W., Leung, K.Y., Chau, P.H., Chung, K.F., 2015. The effect of

- telephone-based cognitive-behavioral therapy on postnatal depression: a randomized controlled trial. *Psychother. Psychosom.* 84, 294–303.
- Nisar, A., Yin, J., Waqas, A., Xue, B., Wang, D., Rahman, A., Li, X., 2019. Prevalence of Perinatal Depression and its Determinants in Mainland China: A Systematic Review and Meta-Analysis (unpublished). Xi'an Jiaotong University, pp. 1–15.
- Rahman, A., Fisher, J., Bower, P., Luchters, S., Tran, T., Yasamy, M.T., Saxena, S., Waheed, W., 2013. Interventions for common perinatal mental disorders in women in low- and middle-income countries: a systematic review and meta-analysis. *Bull. World Health Organ.* 91, 593–601.
- Richter, L.M., Daelmans, B., Lombardi, J., Heymann, J., Boo, F.L., Behrman, J.R., Lu, C., Lucas, J.E., Perez-Escamilla, R., Dua, T., Bhutta, Z.A., Stenberg, K., Gertler, P., Darmstadt, G.L., 2017. Investing in the foundation of sustainable development: pathways to scale up for early childhood development. *Lancet North Am. Ed.* 389, 103–118.
- Singla, D.R., Kohrt, B.A., Murray, L.K., Anand, A., Chorpita, B.F., Patel, V., 2017. Psychological Treatments for the world: lessons from low- and middle-income countries. *Ann. Rev. Clin. Psychol.* 13, 149–181.
- Sun, K., Gao, L., Li, Y., Gao, L., Luo, S., 2011. Effects of psychoeducation program on social support and symptomatology of depression among chinese first-time mothers. *J. Sun Yat-sen Univ. (Chinese)* 32, 90–95.
- Weissman, M.M., Wickramaratne, P., Nomura, Y., Warner, V., Pilowsky, D., Verdelli, H., 2006. Off spring of depressed parents: 20 years later. *Am. J. Psychiatry* 163, 1001–1008.
- Woody, C.A., Ferrari, A.J., Siskind, D.J., Whiteford, H.A., Harris, M.G., 2017. A systematic review and meta-regression of the prevalence and incidence of perinatal depression. *J. Affect. Disord.* 219, 86–92.
- Wu, D., 2018. Discussion on nursing intervention methods of traditional Chinese medicine syndrome differentiation of postpartum depression. *Inner Mongolia Med. J. (Chinese)* 50, 638–639.
- Wu, H., 2017. Influence of continuous nursing on postpartum depression and quality of life in primiparae. *China Modern Med. (Chinese)* 24, 151–153.
- Yang, M., Jia, G., Sun, S., Ye, C., Zhang, R., Yu, X., 2019. Effects of an online mindfulness intervention focusing on attention monitoring and acceptance in pregnant women: a randomized controlled trial. *J. Midwifery Women's Health* 64, 68–77.
- Zhang, J.Y., Cui, Y.X., Zhou, Y.Q., Li, Y.L., 2018. Effects of mindfulness-based stress reduction on prenatal stress, anxiety and depression. *Psychol. Health Med.* 24, 51–58.
- Zhang, Y., Zhou, M., Dang, W., Zhang, L., 2016. Factors affecting the mental health of puerperal women and their treatment strategies. *J. Int. Psychiatry* 43, 182–185.
- Zhao, H., 2018. Effect of continuous nursing intervention based on WeChat platform on postpartum anxiety, depression and self-efficacy of parturients. *Inner Mongolia Med. J. (Chinese)* 50, 616–618.