

RESEARCH ARTICLE

Open Access



Clinical characteristics and risk factors of patients with severe COVID-19 in Jiangsu province, China: a retrospective multicentre cohort study

Songqiao Liu^{1†}, Huanyuan Luo^{2†}, Yuancheng Wang^{3†}, Luis E. Cuevas², Duolao Wang², Shenghong Ju³ and Yi Yang^{1*} 

Abstract

Background: Coronavirus Disease-2019 (COVID-19) pandemic has become a major health event that endangers people health throughout China and the world. Understanding the factors associated with COVID-19 disease severity could support the early identification of patients with high risk for disease progression, inform prevention and control activities, and potentially reduce mortality. This study aims to describe the characteristics of patients with COVID-19 and factors associated with severe or critically ill presentation in Jiangsu province, China.

Methods: Multicentre retrospective cohort study of all individuals with confirmed Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) infections diagnosed at 24 COVID-19-designated hospitals in Jiangsu province between the 10th January and 15th March 2020. Demographic, clinical, laboratory, and radiological data were collected at hospital admission and data on disease severity were collected during follow-up. Patients were categorised as asymptomatic/mild/moderate, and severe/critically ill according to the worst level of COVID-19 recorded during hospitalisation.

Results: A total of 625 patients, 64 (10.2%) were severe/critically ill and 561 (89.8%) were asymptomatic/mild/moderate. All patients were discharged and no patients died. Patients with severe/critically ill COVID-19 were more likely to be older, to be single onset (i.e. not belong to a cluster of cases in a family/community, etc.), to have a medical history of hypertension and diabetes; had higher temperature, faster respiratory rates, lower peripheral capillary oxygen saturation (SpO₂), and higher computer tomography (CT) image quadrant scores and pulmonary opacity percentage; had increased C-reactive protein, fibrinogen, and D-dimer on admission; and had lower white blood cells, lymphocyte, and platelet counts and albumin on admission than asymptomatic/mild/moderate cases. Multivariable regression showed that odds of being a severe/critically ill case were associated with age (year) (OR (Continued on next page)

* Correspondence: yyiyang2004@163.com

[†]Songqiao Liu, Huanyuan Luo and Yuancheng Wang contributed equally to this work.

¹Department of Critical Care Medicine, Jiangsu Provincial Key Laboratory of Critical Care Medicine, Zhongda Hospital, School of Medicine, Southeast University, Nanjing 210009, China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

1.06, 95%CI 1.03–1.09), lymphocyte count ($10^9/L$) (OR 0.25, 95%CI 0.08–0.74), and pulmonary opacity in CT (per 5%) on admission (OR 1.31, 95%CI 1.15–1.51).

Conclusions: Severe or critically ill patients with COVID-19 is about one-tenths of patients in Jiangsu. Age, lymphocyte count, and pulmonary opacity in CT on admission were associated with risk of severe or critically ill COVID-19.

Keywords: COVID-19, SARS-CoV-2, Pneumonia, Severity, Severe, Critically ill

Highlights

- About one-tenths of patients with COVID-19 in Jiangsu had severe or critically ill presentation
- All patients were discharged and no patients died by the end of study
- Risk of severe or critically ill COVID-19 patients increased with age
- Low lymphocyte count and high pulmonary opacity score in CT on admission were associated with high risk of severe or critically ill COVID-19

Background

Coronavirus Disease 2019 (COVID-19), caused by the etiological agent Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), was first reported from Wuhan, Hubei province, China, in December 2019. The World Health Organization (WHO) declared a pandemic the 11th March 2020 [1]. The COVID-19 pandemic have spread quickly from a focal outbreak to over 7410,000 cases with more than 400,000 deaths affecting more than 140 countries by the 12th June 2020 [2].

China had reported over 80,000 confirmed cases by the 13th March 2020 [2]. Although the original epicentre was located in Wuhan, other provinces became affected in the following weeks. In a case series of the first 44,672 confirmed cases, 1023 patients had died, with a crude case fatality rate (CFR) of 2.3%, and mortality was higher among critically ill patients, who had a CFR of 49% [3]. In Hubei, the proportion of health workers with severe or critical ill COVID-19 was higher than in other provinces (10.4 and 7.0%, respectively) [3]. Compared to China, the crude CFR in South Korea is lower among both males (1.1%) and females (0.4%) [4], and the crude CFR in Australia (1.4%) [5] and in countries of the European Union (EU) and European Economic Area (EEA) was also lower (1.5%) [6], while the crude CFR in Italy was much higher (7.2%) [7].

COVID-19 initial symptoms are not specific, presenting with fever, and cough, which can then resolve spontaneously or progress to shortness of breath, dyspnoea, and pneumonia, leading to acute respiratory distress syndrome (ARDS), renal failure, coagulation dysfunction, multiple organ failure and death [8–13]. Demographic

features (eg age and gender); epidemiological features (eg smoking and comorbidities); laboratory parameters (eg C-reactive protein, albumin, cytokine, lactate dehydrogenase, D-dimer, platelet, lymphocyte, and neutrophil); clinical manifestations (eg cough, expectoration, chest pain, and dyspnea); computer tomography (CT) image test results; and others, have been reported to be associated with the severity of COVID-19 in some small studies in China [14–20], and other countries [13, 21, 22].

Understanding the factors associated with COVID-19 disease severity could support the early identification of patients with high risk for severe/critically ill COVID-19 presentation and inform prevention and control activities and reduce mortality. Hubei was the COVID-19 epicentre of China at the time of data collection for this study (from the 10th January 2020 to the 15th March 2020), so patients in other parts of China, outside Hubei, may have different profiles of demographic, epidemiological, clinical characteristics, laboratory parameters, and image test results. Knowing those profiles may help predict the severity of COVID-19.

Jiangsu, a province in China over 600 km from Hubei without common geographical borders and 80 million population, reported over 600 patients infected with COVID-19. We report here an analysis of nearly all cases in Jiangsu province from the 10th January 2020 to the 15th March 2020 to describe the demographic, epidemiological, clinical, laboratory, and imaging characteristics of cases and to identify risk factors for severe/critically ill COVID-19 presentation. Previous studies were mostly small and descriptive in nature. Our study will use data from the whole Jiangsu province and make inferential statistical analyses by means of multivariate regression model to control for possible confounding factors and identify independent risk factors of becoming a severe/critically ill case.

Methods

Study design and population

This is a multicentre retrospective cohort study. All patients were included if they (1) were clinically diagnosed and then confirmed to have COVID-19 in Jiangsu province from the 10th January 2020 up to the 15th March 2020, and (2) fulfilled the diagnostic criteria for the

“Diagnosis and Treatment Protocol for Novel Coronavirus Pneumonia (Trial Version 7)” released by National Health Commission & National Administration of Traditional Chinese Medicine of China [23]. The diagnosis of COVID-19 was based on epidemiological history, clinical manifestations, imaging manifestations of pneumonia in CT scans, and laboratory confirmation [23, 24]. Patients without medical records were excluded.

The Ethics Committee of Zhongda Hospital, Affiliated to Southeast University, approved the study protocols (2020ZDSYLL013–P01 and 2020ZDSYLL019–P01). Patient informed consent was waived due to the retrospective study design.

Data measures

The primary outcome was severe or critically ill within the follow up period. Patients were categorised by disease severity into (1) asymptomatic or mild, or moderate, and (2) severe or critically ill, according to “Diagnosis and Treatment Protocol for Novel Coronavirus Pneumonia (Trial Version 7)” [23]. Asymptomatic infections were defined as the absence of clinical symptoms with a positive nucleic acid test (real-time reverse transcriptase–polymerase chain reaction assay, RT-PCR, for SARS-CoV-2). Mild COVID-19 disease was defined as the presence of mild clinical symptoms without respiratory distress and the absence of imaging manifestations of pneumonia. Moderate disease was defined as the presence of fever, with respiratory symptoms and an image of pneumonia in CT scans. Severe disease was defined as the presence of at least one of the three conditions: respiratory distress, a respiratory rate ≥ 30 beats / min; oxygen saturation in resting state $\leq 93\%$; or an arterial blood oxygen partial pressure / oxygen concentration ≤ 300 mmHg (1 mmHg = 0.133 kPa). Critically ill was defined as having respiratory failure requiring mechanical ventilation, shock or combined organ failure requiring intensive care unit (ICU) monitoring and treatment. A clustering onset was defined as the occurrence of two or more confirmed COVID-19 cases in the same cluster/group within 14 days, such as family, community, hospital, working place or public place, etc. A clustering onset could occur due to interpersonal transmission via close contact with or joint exposure to a confirmed COVID-19 case. Others cases not meeting the conditions of the clustering onset were classified as a single onset. All patients were followed up to the 15th March 2020.

Data were collected using case record forms and electronic medical record systems and included demographic, epidemiological, clinical, laboratory, and imaging information provided by the Data Center of Jiangsu Provincial Health Commission without any patient’s personal information. The day of admission to a

COVID-19-designated hospital was considered the first day of hospitalisation. The severity of illness was assessed by two physicians. Severity was assessed at days 1, 2, 3, 4, 5, 6, 7 and 14 after admission and patients were followed for up to discharge. ICU admissions were recorded. Imaging grading was performed by two independent radiologists with more than 5 years’ experience in pulmonary imaging. Chest CT axial sections were divided into quadrants (left and right, anterior and posterior) by drawing horizontal and vertical lines through the centre of the chest. Quadrant scores were estimated as the sum of quadrants with pulmonary opacities extending from the proximal to the distal end of the chest and ranged from 0 to 4. Pulmonary opacity was visually estimated and assigned a percentage of pulmonary opacity area in the area of bilateral lungs, rounded to the nearest 5%.

Statistical analysis

Variables

Primary outcome variable was the occurrence of severe/critically ill case. Predictive variables included sex, age, exposure type, types of disease onset, initial symptoms, medical history, vital signs (including body temperature, heart rate (HR), respiratory rate, and peripheral capillary oxygen saturation (SpO_2)), CT image parameters (including quadrant score and pulmonary opacity), and laboratory parameters (including white blood cells count, neutrophil, lymphocyte, platelet, albumin, creatinine, C-reactive protein, activated partial thromboplastin time, fibrinogen, and D-dimer). Those variables were measured at hospital admission. The other variables including supportive treatments and medical drugs were collected during the follow-up period.

Statistical analysis

Continuous variables were described using means (standard deviations, SD) or medians (with inter-quartile range, IQR) by disease severity and were compared using ANOVA or Kruskal-Wallis tests as appropriate. Categorical variables were summarised using frequencies and percentages and compared using Fisher exact tests.

Logistic regression models were used to identify the risk factors for having a severe or critically ill status. Analysis was performed in 2 steps. Firstly, a univariate logistic regression model was fitted for each variable based on completed cases. As there are many potential predictors, we chose the variables for univariate regression analysis if a variable is significant at 5%. Respiratory rate and SpO_2 were not included in the regression analyses since they were part of criterion for classifying the disease severity. All variables selected for univariate regression analysis were also included in the second stage of the multivariate logistic regression. Missing covariates

at admission were imputed with multiple imputation using a Markov Chain Monte Carlo simulation method with 10 iterations. In the logistic regression analysis, odds ratios (ORs) for having a severe or critically ill status for each variable were calculated along with 95% confidence intervals (CIs). A sensitivity analysis was performed on the completed cases. The 2-tailed $P < 0.05$ was considered as statistically significant for all analyses. The analyses were performed using SAS 9.4 (SAS Institute).

Results

From the 10th January 2020 to 15th March 2020, 721 suspected cases with possible COVID-19 were admitted in 24 hospitals in Jiangsu province, China, while 90 cases were excluded because of negative reverse transcriptase–polymerase chain reaction assay. Six hundred thirty-one cases were diagnosed with COVID-19 totally. Of these, 625 (99.0%) had retrievable medical records and were included in the analysis (Fig. 1).

Table 1 describes the characteristics of patients at the time of admission by disease severity and Table S1 provides a more detailed description of the five severity categories. 561 (89.8%) patients were asymptomatic/mild/moderate and 64 (10.2%) patients were severe or critically ill. Patients with severe/critically ill COVID-19 were more likely to be older and to be single onset (i.e. not belong to a cluster of cases in a family/community, etc.). Patients with severe/critically ill presentation were more likely to have a medical history of hypertension and diabetes. Patients with severe/critically ill COVID-19 on admission had higher temperature, faster respiratory rates, lower SpO₂, and higher CT image quadrant scores and pulmonary opacity percentage (Table 1 and Table S1).

Cases with severe/critically ill presentation were more likely to have increased C-reactive protein, fibrinogen, and D-dimer than asymptomatic/mild/moderate cases. Similarly, severe/critically ill cases had lower white blood cells, lymphocyte, and platelet counts and albumin (Table 1). As expected, severe cases were more likely to use supportive treatments and medical drugs, including antibiotics and antivirals, except interferon (Table 2).

None of the patients died and 625 (100%) of 625 patients were discharged by the end of study (15th March 2020). The results from the univariate and multivariate logistic regression analyses are presented in Table 3. Factors independently associated with severe or critically ill infection included age (year) (OR 1.06, 95%CI 1.03–1.09), lymphocyte count (10⁹/L) (OR 0.25, 95%CI 0.08–0.74), and pulmonary opacity in CT (per 5%) on admission (OR 1.31, 95%CI 1.15–1.51). Sensitivity analysis showed that they remained statistically significant in a logistic model with only above three variables based on the completed cases without missing data.

Discussion

In this large multicentre cohort, 64 (10.2%) of 625 patients were severe or critically ill. This proportion of severe or critically ill cases is lower than the 17.7% reported from Wuhan but similar to the 10.4% in Hubei province and higher than the 7.0% reported for areas outside Hubei province [3]. Specifically, it is lower than the figures reported from several case series from Wuhan including 13 (32%) ICU admission among 41 cases with 6 (15%) deaths [25]; 11 (11%) deaths among 99 cases [9]; and 36 (26.1%) ICU admissions among 138 patients, with 6 deaths (4.3%) [10]. The lower proportion than that in Hubei is likely due to several factors, including more adequate medical resources, better disease




Fig. 1 Study flow diagram

Table 1 Demographic and clinical characteristics of patients with COVID-19 at admission^a

Category	Characteristics	Severe/Critically ill			P-value
		Yes (N = 64)	No (N = 561)	All (N = 625)	
Demographic, n/N (%), N, mean (SD)					
	Male	41/64(64.1%)	288/561(51.3%)	329/625(52.6%)	0.0534
	Female	23/64(35.9%)	273/561(48.7%)	296/625(47.4%)	
	Age (year)	64,59.53(13.43)	561,42.72(16.73)	625,44.44(17.19)	<.0001
	≤18 years	0/64(0.0%)	37/561(6.6%)	37/625(5.9%)	<.0001
	19–44 years	6/64(9.4%)	255/561(45.5%)	261/625(41.8%)	
	45–64 years	32/64(50.0%)	216/561(38.5%)	248/625(39.7%)	
	65+ years	26/64(40.6%)	53/561(9.4%)	79/625(12.6%)	
Exposure type, n/N (%)					
	Imported cases	25/64(39.1%)	194/561(34.6%)	219/625(35.0%)	0.4765
	Local cases	39/64(60.9%)	367/561(65.4%)	406/625(65.0%)	
Types of disease onset, n/N (%)					
	Single onset	40/64(62.5%)	270/561(48.1%)	310/625(49.6%)	0.0294
	Clustering onset	24/64(37.5%)	291/561(51.9%)	315/625(50.4%)	
Initial symptoms, n/N (%)					
	Fever	52/64(81.3%)	360/561(64.2%)	412/625(65.9%)	0.0063
	Cough	44/64(68.8%)	300/561(53.5%)	344/625(55.0%)	0.0200
	Sputum	25/64(39.1%)	141/561(25.1%)	166/625(26.6%)	0.0168
Medical history, n/N (%)					
	Hypertension	19/64(29.7%)	72/561(12.8%)	91/625(14.6%)	0.0003
	Diabetes	10/64(15.6%)	30/561(5.3%)	40/625(6.4%)	0.0015
	Stroke	2/64(3.1%)	8/560(1.4%)	10/624(1.6%)	0.2736
Vital signs, N, mean (SD)					
	Temperature (°C)	64,37.30(0.94)	561,37.02(0.70)	625,37.05(0.73)	0.0040
	HR (bpm)	64,89.98(15.06)	561,86.84(13.25)	625,87.17(13.46)	0.0772
	Respiratory rate (breath per min)	64,20.98(4.87)	561,18.87(2.04)	625,19.08(2.56)	<.0001
	SpO ₂ (%)	64,95.53(4.70)	561,97.92(1.15)	625,97.68(1.99)	<.0001
CT image, N, median (IQR)					
	Quadrant score (1–4)	58,4.0(4.0–4.0)	438,2.0(1.0–4.0)	496,2.0(1.0–4.0)	<.0001
	Pulmonary opacity (%)	58,50.0(35.0–70.0)	438,20.0(5.0–30.0)	496,20.0(5.0–40.0)	<.0001
Laboratory test, N, median (IQR)					
	WBC Count (10 ⁹ /L)	52,4.3(3.4–5.8)	461,5.0(4.0–6.3)	513,4.9(3.9–6.2)	0.0440
	Neutrophil (10 ⁹ /L)	52,2.9(2.0–4.4)	455,3.0(2.2–4.0)	507,3.0(2.2–4.0)	0.7435
	Lymphocyte (10 ⁹ /L)	52,0.7(0.5–1.0)	453,1.4(1.0–1.8)	505,1.3(0.9–1.7)	<.0001
	Platelet (10 ⁹ /L)	48,154.0(118.0–185.0)	446,188.5(154.0–222.0)	494,183.5(151.0–219.0)	<.0001
	Albumin (g/L)	48,39.7(34.2–41.4)	432,42.0(38.0–45.7)	480,41.4(38.0–45.1)	<.0001
	Creatinine (umol/L)	49,64.0(51.0–83.0)	427,63.8(51.0–79.0)	476,63.9(51.0–79.0)	0.6950
	C-reactive protein (mg/L)	44,40.1(8.6–92.7)	430,10.0(2.6–19.3)	474,10.0(2.7–22.6)	<.0001
	Activated partial thromboplastin time (s)	54,32.6(28.5–36.5)	459,32.2(27.9–37.4)	513,32.2(28.0–37.2)	0.8158
	Fibrinogen (g/L)	53,4.3(3.2–5.9)	443,3.4(2.7–4.1)	496,3.5(2.7–4.2)	<.0001
	D-dimer (mg/L)	51,0.3(0.2–1.0)	424,0.2(0.1–0.4)	475,0.2(0.1–0.4)	0.0003

^aContinuous variables: ANOVA or Kruskal-Wallis tests as appropriate; categorical variables: Fisher exact tests

Table 2 Clinical management and outcome

Category	Clinical management and outcome	n (%) or median (IQR)			P-value
		Severe/Critically ill			
		Yes (N = 64)	No (N = 561)	All (N = 625)	
Supportive treatments	Inotropic and vasoconstrictive agents	5(7.8%)	0(0.0%)	5(0.8%)	<.0001
	Nasal cannula	53(82.8%)	168(29.9%)	221(35.4%)	<.0001
	Mask	12(18.8%)	2(0.4%)	14(2.2%)	<.0001
	High-flow nasal cannula oxygen therapy	24(37.5%)	1(0.2%)	25(4.0%)	<.0001
	Non-invasive ventilation	34(53.1%)	0(0.0%)	34(5.4%)	<.0001
	Intermittent mandatory ventilation	5(7.8%)	0(0.0%)	5(0.8%)	<.0001
	Prone position	17(26.6%)	1(0.2%)	18(2.9%)	<.0001
	Continuous renal replacement therapy	1(1.6%)	0(0.0%)	1(0.2%)	0.1024
	Extracorporeal membrane oxygenation	2(3.1%)	0(0.0%)	2(0.3%)	0.0103
	Lung transplantation	2(3.1%)	0(0.0%)	2(0.3%)	0.0103
Medical drugs	Traditional Chinese medicine	29(45.3%)	69(12.3%)	98(15.7%)	<.0001
	Immunoglobulin	50(78.1%)	106(18.9%)	156(25.0%)	<.0001
	Interferon	47(73.4%)	456(81.3%)	503(80.5%)	0.1363
	Antioxidants	35(54.7%)	117(20.9%)	152(24.3%)	<.0001
	Glucocorticoid	52(81.3%)	90(16.0%)	142(22.7%)	<.0001
	Thymosin	43(67.2%)	101(18.0%)	144(23.0%)	<.0001
	Neurotrophic drugs	21(32.8%)	81(14.4%)	102(16.3%)	0.0005
	Any antibiotics	59(92.2%)	277(49.4%)	336(53.8%)	<.0001
	Any antivirals	64(100%)	516(92.0%)	580(92.8%)	0.0098
Clinical outcome	Death	0(0.0%)	0(0.0%)	0(0.0%)	NC
	Hospital stay	21.5(15.0–29.0)	15.0(12.0–21.0)	16.0(12.0–22.0)	<.0001

Table 3 Factors associated with severe/critically ill in patients with COVID-19: Results from logistic regression analysis

Variables	Univariate analysis ^a		Multivariate analysis ^b	
	Odds ratio (95%CI)	P-value	Odds ratio (95%CI)	P-value
Age (year)	1.07(1.05,1.09)	<.0001	1.06(1.03,1.09)	<.0001
Single onset	1.80(1.05,3.06)	0.0311	0.92(0.43,1.96)	0.8275
Fever	2.42(1.26,4.64)	0.0078	1.50(0.64,3.54)	0.3542
Cough	1.91(1.10,3.33)	0.0216	1.24(0.54,2.87)	0.6110
Sputum	1.91(1.12,3.27)	0.0183	1.12(0.48,2.60)	0.7994
Hypertension	2.87(1.59,5.18)	0.0005	1.06(0.47,2.40)	0.8874
Diabetes	3.28(1.52,7.07)	0.0025	1.64(0.52,5.22)	0.4004
Temperature (°C)	1.59(1.15,2.19)	0.0046	0.95(0.61,1.47)	0.8133
Lymphocyte (10 ⁹ /L)	0.03(0.01,0.08)	<.0001	0.25(0.08,0.74)	0.0161
Platelet (10 ⁹ /L)	0.99(0.98,0.99)	0.0003	1.00(0.99,1.00)	0.5147
Albumin (g/L)	0.91(0.87,0.96)	0.0002	0.99(0.92,1.07)	0.8344
C-reactive protein (mg/L)	1.02(1.01,1.02)	<.0001	1.00(0.99,1.01)	0.9789
Fibrinogen (g/L)	1.87(1.50,2.32)	<.0001	1.04(0.72,1.49)	0.8327
D-dimer (mg/L)	1.28(1.06,1.55)	0.0088	1.17(0.83,1.66)	0.3625
Quadrant score (1–4)	2.28(1.71,3.05)	<.0001	0.90(0.56,1.47)	0.6811
Pulmonary opacity (per 5%)	1.38(1.28,1.49)	<.0001	1.31(1.15,1.51)	0.0001

^aUnivariate analysis is based on the complete cases without missing value

^bMultivariate analysis is based on imputed values for missing data in Lymphocyte, Platelet, Albumin, C-reactive protein, Fibrinogen, D-dimer, Quadrant score and Pulmonary opacity using multiple imputation method

recognition and testing capacity, earlier identification of asymptomatic and mild cases and a more informed supportive care in COVID-19-designated hospitals. The proportion of critically ill cases in early stage of COVID-19 outbreak in New York City, USA was much higher (22% [257]) [26] than in China, with 14.2% treated in the ICU [27] and 23.6% required mechanical ventilation reported in other studies [28]. Singapore also reported a higher proportion of severe cases in an early study (33.3% [6]) [29]. A more recent study from USA reported 5.8% (7162) of cases suffered from severe COVID-19 [30] which is lower than in China. The proportions of severe COVID-19 in different countries vary a lot, which may result from quite different study designs, the COVID-19 outbreak stages when data were collected, population characteristics, health resources, government response measurements, et al.

Despite this being a hospital-based study, some patients had no symptoms. This is likely due to testing of contacts after the identification of an index case and the policy of hospitalization of all infected individuals at the initial stages of the epidemic, independently of the presence of symptoms. This study found that fever, cough, and sputum were very common among patients with COVID-19 and more frequent in patients with severe or critically ill COVID-19. Similar to COVID-19, fever and cough are the most common symptoms of the other two diseases caused by coronavirus, i.e. severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) [31, 32]. Fever is a primary symptom for cytokine storms, with the production of high concentrations of cytokines stimulating abnormally excessive immune responses and inflammation [33–35]. Vital signs showed severe or critically ill patients had higher body temperature and respiratory rate, and lower SpO₂ on admission. SpO₂ < 90% has been used as a marker for the use of glucocorticoids during the outbreak [36], and the oxygenation saturation index is associated with ARDS severity and increased mortality [37, 38].

Gender had no effect on severity of COVID-19 patients. Although early reports from Wuhan indicated more men than women had severe COVID-19, recent studies reported similar proportions of men and women admitted to ICUs [9, 11, 25], suggesting gender differences disappeared with higher incidence. Earlier reports may have included more males due to a higher occupational infection risk for males in the markets and congregation places [10].

Our study found that age was independently associated with severe or critically ill presentation. Age is a well-established factor for severe/critically ill COVID-19 for individuals > 60, and especially over 80 years old [39, 40]. Similarly, previous reports have indicated patients in ICUs are older than non-ICU patients [10], and that

CFRs are higher among older individuals [9, 11, 25]. Older patients also have faster disease progression than younger patients [41], which is similar to the MERS and SARS presentations, in which, older age (> 60 or 45) is associated with disease severity (MERS) [42, 43], and mortality [44, 45]. Older age reflects a greater likelihood of underlying medical conditions such as hypertension and diabetes, which predisposes to immunological vulnerabilities. Also, age-related immunosenescence may also contribute to the severe disease [46].

In Wuhan, many asymptomatic patients had abnormal lung CT findings on admission, which then progressed to diffuse ground-glass opacities and consolidation [47]. In Jiangsu, several asymptomatic cases also had radiological changes presented as low quadrant scores and pulmonary opacity scores on admission and severe/critically-ill cases had higher CT quadrant and pulmonary opacity scores than moderate cases. Our study also identified pulmonary opacity as an independent predictor of severe/critical illness. This is consistent with the previous study reporting that the CT visual quantitative evaluation of acute lung inflammatory lesions involving each lobe in severe or critical cases was significantly higher than less severe cases [48].

We found severe or critically ill patients had more obvious damage of white blood cells and immune cells such as lymphocytes with lymphocytes identified as an independent predictor of more severe disease. COVID-19 may cause the reduced T lymphocytes, especially CD4 + T and CD8 + T cells, leading to reduced IFN- γ production, which may be related to the severity of disease [49]. In addition, severe or critically ill patients showed more serious organ dysfunction like reduced albumin on admission which may be a sign of reduced liver production and increased gastrointestinal or renal loss, and increased fibrinogen on admission responding to systemic inflammation and tissue damage, and more fierce inflammatory response presented as much higher level of inflammatory markers, such as C-reactive protein [50–52].

This study has several strengths. Firstly, this is one of the largest studies describing the clinical characteristics of patients with COVID-19 and risk factors for severe/critically ill infection outside the Wuhan, epicentre of the epidemic in China at the time of data collection for this study (from the 10th January 2020 to the 15th March 2020). Secondly, the cohort includes almost all COVID-19 cases in the province, which may have reduced selection bias. Thirdly, Jiangsu province, which is far from Hubei, provides an opportunity to assess the demographic, epidemiological, clinical, laboratory, and imaging features of cases imported from other provinces and local cases. Fourthly, asymptomatic and mild cases were included, which provides a more comprehensive

description of the characteristics of COVID-19 cases with a broad spectrum of disease severity.

There are also limitations that need mentioning. Firstly, laboratory and radiological data had a large amount of missing data preventing their integration in the analysis. Secondly, the predictive factors identified may be subject to uncontrolled confounders by unknown/unmeasured factors such as occupation and pregnancy. Medical staff and pregnant women may have different severity profiles. Thirdly, this is a retrospective observational study and data may be susceptible to measurement and information bias.

Conclusion

In conclusion, this study demonstrates that patients with COVID-19 in Jiangsu had a low rate of severe or critically ill presentation, with no deaths recorded. The COVID-19 severity is associated with epidemiological and clinical characteristics, laboratory test, and radiological findings. Age, lymphocyte count, and pulmonary opacity in CT on admission were independently associated with risk of severe or critically ill COVID-19 presentation.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12879-020-05314-x>.

Additional file 1: Table S1. Demographic and clinical characteristics of patients at admission by disease severity*.

Abbreviations

COVID-19: Coronavirus Disease-2019; SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus-2; SpO₂: Peripheral capillary oxygen saturation; CT: Computer tomography; WHO: World Health Organization; CFR: Case fatality rate; EU: The European Union; EEA: European Economic Area; ARDS: Acute respiratory distress syndrome; RT-PCR: Reverse transcriptase–polymerase chain reaction assay; ICU: Intensive care unit; HR: Heart rate; SD: Standard deviation; IQR: Inter-quartile range; OR: Odds ratio; CI: Confidence interval; SARS: Severe acute respiratory syndrome; MERS: Middle East respiratory syndrome

Acknowledgements

We thank all patients and their families involved in the study. We thank all the doctors and nurses who are working to combat COVID-19.

Authors' contributions

YY, SJ and DW conceived and designed the study. SL, HL and YW contributed to the literature search. SL, YW, SJ, YY, and DW contributed to data collection, quality checks and data management. DW, SJ, SL, HL, LEC, YW and YY contributed to data analysis and results presentation. DW, SJ, SL, LEC, HL and YY were responsible for results interpretation. SL, HL, YW, DW, LEC, SJ, and YY contributed in the drafting and review of the manuscript. The authors read and approved the final manuscript.

Funding

This work was supported, in part, by the National Natural Science Foundation of China (81971885, 81971888) and the Scientific Research Project of Jiangsu Health Committee (H2018048). The funders had no role in the study design, data collection, data analysis, interpretation of data and in writing the manuscript.

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This study was conducted in accordance with the International Council for Harmonisation Guidelines for Good Clinical Practice and the Declaration of Helsinki. All the data in this study were anonymised and provided with administrative permissions from Jiangsu Provincial Health Commission. The Ethics Committee of Zhongda Hospital, Affiliated to Southeast University, approved the study protocols (2020ZDZSYLL013–P01 and 2020ZDZSYLL019–P01). Patient informed consent was waived due to the retrospective study design.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Critical Care Medicine, Jiangsu Provincial Key Laboratory of Critical Care Medicine, Zhongda Hospital, School of Medicine, Southeast University, Nanjing 210009, China. ²Department of Clinical Sciences, Liverpool School of Tropical Medicine, L3 5QA, Liverpool, UK. ³Department of Radiology, Zhongda Hospital, School of Medicine, Southeast University, Nanjing 210009, China.

Received: 19 April 2020 Accepted: 30 July 2020

Published online: 06 August 2020

References

- World Health Organization. WHO director-General's opening remarks at the media briefing on COVID-19; 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Accessed 13 June 2020.
- World Health Organization. Coronavirus disease (COVID-19) situation reports. Accessed 13 June 2020. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>.
- Chinese Center for Disease Control and Prevention. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020;41(2):145–51.
- Shim E, Tariq A, Choi W, Lee Y, Chowell G. Transmission potential and severity of COVID-19 in South Korea. *Int J Infect Dis*. 2020;93:339–44.
- COVID-19 National Incident Room Surveillance Team. COVID-19, Australia: epidemiology report 17 (fortnightly reporting period ending 24 may 2020). *Commun Dis Intell*. 2018;2020:44.
- European Centre for Disease Prevention and Control. Coronavirus disease 2019 (COVID-19) in the EU/EEA and the UK – eighth update. Accessed 13 June 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-rapid-risk-assessment-coronavirus-disease-2019-eighth-update-8-april-2020.pdf>.
- Livingston E, Bucher K. Coronavirus disease 2019 (COVID-19) in Italy. *JAMA*. 2020. <https://doi.org/10.1001/jama.2020.4344> Online ahead of print.
- Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382(18):1708–20.
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507–13.
- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061–9.
- Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020;8(5):475–81.
- Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of COVID-19 in new York City. *N Engl J Med*. 2020;382(24):2372–4.
- Mitra AR, Fergusson NA, Lloyd-Smith E, et al. Baseline characteristics and outcomes of patients with COVID-19 admitted to intensive care units in Vancouver, Canada: a case series. *CMAJ*. 2020;1:200794.

14. Li K, Wu J, Wu F, et al. The clinical and chest CT features associated with severe and critical covid-19 pneumonia. *Investig Radiol.* 2020;55(6):327–31.
15. Chen W, Zheng KI, Liu S, Yan Z, Xu C, Qiao Z. Plasma CRP level is positively associated with the severity of COVID-19. *Ann Clin Microbiol Antimicrob.* 2020;19(1):18.
16. Gao F, Zheng KI, Wang XB, et al. Obesity is a risk factor for greater COVID-19 severity. *Diabetes Care.* 2020;1:dc200682.
17. Li J, Li M, Zheng S, et al. Plasma albumin levels predict risk for nonsurvivors in critically ill patients with COVID-19. *Biomark Med.* 2020. <https://doi.org/10.2217/bmm-2020-0254>.
18. Li X, Xu S, Yu M, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol.* 2020;50091:30495.
19. Li Y, Zhao K, Wei H, et al. Dynamic relationship between D-dimer and COVID-19 severity. *Br J Haematol.* 2020. <https://doi.org/10.1111/bjh.16811>.
20. Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: a meta-analysis. *Clin Chim Acta.* 2020;506:145–8.
21. Simonnet A, Chetboun M, Poissy J, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity (Silver Spring).* 2020. <https://doi.org/10.1002/oby.22831>.
22. Bhatraju PK, Ghassemieh BJ, Nichols M, et al. Covid-19 in critically ill patients in the Seattle region - case series. *N Engl J Med.* 2020;382(21):2012–22.
23. National Health Commission & National Administration of Traditional Chinese Medicine. Diagnosis and treatment protocol for novel coronavirus pneumonia (trial version 7). *Chin Med J.* 2020;1:133. <https://doi.org/10.1097/CM9.0000000000000819>.
24. Sun Q, Qiu H, Huang M, Yang Y. Lower mortality of COVID-19 by early recognition and intervention: experience from Jiangsu Province. *Ann Intensive Care.* 2020;10(1):33.
25. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497–506.
26. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in new York City: a prospective cohort study. *Lancet.* 2020;395(10239):1763–70.
27. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the new York City area. *JAMA.* 2020;323(20):2052–9.
28. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in new York City: prospective cohort study. *BMJ.* 2020;369:m1966.
29. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA.* 2020; 323(15):1488–94.
30. CDC COVID-19 Response Team. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019 - United States, February 12–march 28, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(13):382–6. <https://doi.org/10.15585/mmwr.mm6913e2>.
31. Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. *BMJ.* 2020;368:m606.
32. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA.* 2020;323(13):1239–42.
33. Zhang W, Zhao Y, Zhang F, et al. The use of anti-inflammatory drugs in the treatment of people with severe coronavirus disease 2019 (COVID-19): the perspectives of clinical immunologists from China. *Clin Immunol.* 2020;214: 108393.
34. Lee DW, Gardner R, Porter DL, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood.* 2014;124(2):188–95.
35. Cron R, Behrens EM. Cytokine storm syndrome. 1st ed: Springer International Publishing; 2019.
36. Zhang J, Zhou L, Yang Y, Peng W, Wang W, Chen X. Therapeutic and triage strategies for 2019 novel coronavirus disease in fever clinics. *Lancet Respir Med.* 2020;8(3):e11–2.
37. DesPrez K, McNeil JB, Wang C, Bastarache JA, Shaver CM, Ware LB. Oxygenation saturation index predicts clinical outcomes in ARDS. *Chest.* 2017;152(6):1151–8.
38. Chen WL, Lin WT, Kung SC, Lai CC, Chao CM. The value of oxygenation saturation index in predicting the outcomes of patients with acute respiratory distress syndrome. *J Clin Med.* 2018;7:8.
39. Gralinski LE, Menachery VD. Return of the coronavirus: 2019-nCoV. *Viruses.* 2020;12:2.
40. Wu J, Li W, Shi X, et al. Early antiviral treatment contributes to alleviate the severity and improve the prognosis of patients with novel coronavirus disease (COVID-19). *J Intern Med.* 2020. <https://doi.org/10.1111/joim.13063>.
41. Wang W, Tang J, Wei F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol.* 2020;92(4): 441–7.
42. Feikin DR, Alraddadi B, Qutub M, et al. Association of higher MERS-CoV virus load with severe disease and death, Saudi Arabia, 2014. *Emerg Infect Dis.* 2015;21(11):2029–35.
43. Ko JH, Park GE, Lee JY, et al. Predictive factors for pneumonia development and progression to respiratory failure in MERS-CoV infected patients. *J Inf Secur.* 2016;73(5):468–75.
44. Majumder MS, Klueberg SA, Mekaru SR, Brownstein JS. Mortality risk factors for middle east respiratory syndrome outbreak, South Korea, 2015. *Emerg Infect Dis.* 2015;21(11):2088–90.
45. Chan JWM, Ng CK, Chan YH, et al. Short term outcome and risk factors for adverse clinical outcomes in adults with severe acute respiratory syndrome (SARS). *Thorax.* 2003;58(8):686–9.
46. Maue AC, Yager EJ, Swain SL, Woodland DL, Blackman MA, Haynes L. T-cell immunosenescence: lessons learned from mouse models of aging. *Trends Immunol.* 2009;30(7):301–5.
47. Shi H, Han X, Jiang N, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis.* 2020;20(4):425–34.
48. Li K, Fang Y, Li W, et al. CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19). *Eur Radiol.* 2020;1:1–10.
49. Chen G, Wu D, Guo W, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest.* 2020;130(5):2620–9.
50. Sproston NR, Ashworth JJ. Role of C-reactive protein at sites of inflammation and infection. *Front Immunol.* 2018;9:754.
51. Singh T, Newman AB. Inflammatory markers in population studies of aging. *Ageing Res Rev.* 2011;10(3):319–29.
52. Jousilahti P, Salomaa V, Rasi V, Vahtera E, Palosuo T. The association of C-reactive protein, serum amyloid a and fibrinogen with prevalent coronary heart disease—baseline findings of the PAIS project. *Atherosclerosis.* 2001; 156(2):451–6.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

