

Potential health impact and cost-effectiveness of drug therapy for prehypertension

Tao Chen, PhD, MD^{1,2,3}; Dahai Yu, PhD, MD^{1,4}; Victoria Cornelius⁵, PhD; Rui Qin, MSc⁶; Yamei Cai, MSc¹; Zhixing Jiang, MSc⁷; Zhanzheng Zhao, PhD, MD¹

1, Department of Nephrology, the First Affiliated Hospital, Zhengzhou University, Zhengzhou 450052, China

2, Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool L3 5QA, UK

3, Department of Epidemiology and Biostatistics, Nanjing Medical University, Nanjing, 210029, China

4, Arthritis Research UK Primary Care Centre, Research Institute for Primary Care & Health Sciences, Keele University, Keele ST5 5BG, UK

5, Imperial Clinical Trials Unit, School of Public Health, Imperial College London, London, W12 7RH, UK

6, Jiangsu Province Hospital on Integration of Chinese and Western Medicine, Nanjing, 210028, China

7, Department of Cardiology, Jiangsu Province People's Hospital, Nanjing 210029, China

Authors' contributions

TC and DHY contributed to the conception and design of this study. TC, ZZZ, ZXJ, RQ and DHY carried out the acquisition, analysis and interpretation of data. TC, DHY and RQ wrote the first draft of the manuscript; VC, YMC and ZXJ revised the document for important intellectual content. All authors read and approved the final manuscript.

Corresponding author:

Professor Zhanzheng Zhao, Department of Nephrology, The First Affiliated Hospital Zhengzhou University, 1 Jianshe E Rd, Erqi, Zhengzhou, Henan, 450052, CHINA

Email: zhanzhengzhao@zzu.edu.cn

TEL:+86 139 3852 5666

FAX:+86 371 6698 8753

1 **Funding sources**

2 This study was supported by Ministry of Education of China (20133234120008).

3 **Disclosures**

4 The authors declared no conflict of interest

5 **Key words:**

6 Cost-effectiveness analysis; hypertension; drugs; cardiovascular disease

7

1

2 **Abstract**

3

4 **Background:**

5

6 Studies have reported that pharmacologic interventions with candesartan or ramipril could
7 reduce the risk of hypertension among prehypertensive subjects free of clinical
8 cardiovascular disease(CVD), however, the cost-effectiveness and long-term cardiovascular
9 risk of drug treatment among these population is unclear.

10 **Method:**

11 A Markov state-transition model was developed to simulate a hypothetical cohort of Chinese
12 adults with high-range prehypertension (130-139/85-89 mmHg) but without CVD. Data on
13 the incidence of CVD and hypertension was obtained from corresponding risk equations.
14 Utility and disease-related costs were obtained from published literatures. Robustness and
15 uncertainty was evaluated using deterministic and probabilistic sensitivity analyses.

16 **Results:**

17 Compared with placebo, drug treatment resulted in delaying the development of hypertension
18 by nearly 12 years and reducing the absolute incidence of hypertension by 32.01% over
19 lifetime. The cumulative incidence of coronary heart disease, stroke and heart failure were
20 reduced and survival was improved from 28.46 to 28.80 years. The average incremental cost
21 effectiveness ratio for drug treatment was \$12,994 per quality-adjusted life-year and the value
22 was mostly sensitive to the effect size of treatment and age starting treatment. At a
23 willingness-to-pay threshold of $>3 \times$ China gross domestic product per capita in 2014, there
24 was a 30.48% chance that drug treatment would remain cost-effective and a low chance of
25 being cost-effective if relative risk of treatment on hypertension was larger than 0.64.

26 **Conclusion:** Drug treatment for prehypertension may help stem the current epidemic of
27 hypertension among Chinese adults free of CVD, which may in turn reduce CVD

1 complications and potentially be cost effective.

2 **1. Introduction**

3

4 Hypertension is the leading risk factor for cardiovascular disease (CVD) in China and it
5 contributes to 2.33 million cardiovascular deaths in 2005[1]. Despite tremendous efforts to
6 promote healthful lifestyles and drug treatment rate, the prevalence of hypertension has
7 increased from 20% among men and 17% among women in 2002 to 31.2% and 28.0%,
8 respectively, in 2009[2, 3]. The term of “prehypertension” was introduced in Seventh Report
9 of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High
10 Blood Pressure (JNC-VII) to highlight an at-risk population. It is believed that
11 prehypertension is a precursor of clinical hypertension and successfully control this
12 population would reduce the potential cardiovascular burden [4].

13 Currently, the treatment of prehypertension is primarily nonpharmacological lifestyle changes
14 such as healthy dietary habits, overweight/obesity control [5]. However, an important
15 obstacle to the success of these strategies is that lifestyle modification treatments are difficult
16 to maintain and may require large-scale social and policy changes as well as the patients
17 himself [5]. The trial of preventing hypertension(TROPHY) study was the first trial to
18 perform pharmacological intervention amongst individuals with prehypertension. Results
19 from TROPHY suggested that treatment with Candesartan significantly reduced the incidence
20 of hypertension, even after discontinuing drug therapy for 2 years [6]. This is in agreement
21 with another clinical trial (prevention of hypertension with the angiotensin-converting
22 enzyme inhibitor ramipril in patients with high-normal blood pressure(PHARAO) study)
23 performed in Germany [7]. In this study, Ramipril is proved to be effective in reducing the
24 incidence of office hypertension by 34.4%. However, there are no studies to assess whether
25 early drug treatment of prehypertension can reduce its long-term cardiovascular
26 consequences.

1 Additionally, nearly 30% of the general adult population in China have prehypertension and
2 the health and economic consequences caused by prehypertension are substantial [8-10]. It is
3 important that an effective strategy must be carefully chosen on the basis of both public
4 health and economic reasons. However, we are still unclear whether prescribing drugs among
5 individuals with prehypertension but without clinical CVD is cost-effective, given the
6 relatively low absolute risk amongst this population [11, 12].

7 The aim of our study was to combine data from published clinical trials and epidemiological
8 studies to predict 1) medium and long-term hypertension incidence and its associated
9 cardiovascular risk and 2) the lifetime cost-effectiveness of pharmacological therapy versus
10 placebo intervention among individuals with prehypertension free of CVD.

11 **2. Method**

12 *2.1 Model structure*

13 We adopted a similar approach to develop a state-transition Markov model used by Herman
14 et al for the primary prevention of type 2 diabetes mellitus in adults with impaired glucose
15 tolerance [13]. Our model simulated the progression from prehypertension to onset of
16 hypertension, to hypertension with complications and death. Simultaneously, this model
17 allowed individuals with prehypertension to move to CVD/death directly (**Figure 1**). In total,
18 we modelled the following 10 distinct health states: (1) healthy with prehypertension; (2)
19 hypertension; (3) stroke; (4) post stroke; (5) coronary heart disease(CHD); (6) post CHD; (7)
20 heart failure (HF); (8) post HF; (9) CVD mortality and (10) non-CVD mortality. After each
21 cycle, cohort members were redistributed to one of the above ten health states. Individuals
22 reaching a CVD state by the end of the first year could subsequently transit to another state
23 (e.g., the occurrence of stroke after HF). Subsequent to a primary event, patients might move
24 to a chronic health state (post CHD or post HF or post stroke), which had a higher risk to
25 suffer a new CVD event or death. Due to the lack of data, some states were not modelled (e.g.,

1 myocardial infarction after post-stroke). Subjects transiting to hypertension had a similar
2 pathway as those with prehypertension, but they could not return to prehypertension (linear
3 model without remission) and had a different pre-specified transition probability. The detailed
4 transition pathways could be found in the **online supplementary Figure A1 to Figure A5**.

5 In this study, we simulated two cohorts: one that followed individuals who received placebo
6 treatment (placebo intervention) and the other one which followed individuals who were
7 prescribed antihypertensive treatment (drug treatment). This will keep consistence with the
8 two completed trials[6-7] after considering the fact that no head-to-head clinical trials were
9 performed to compare drug treatment, intensive lifestyle changes and usual care. All
10 prehypertensive individuals entered the model with their pre-specified characteristics and
11 were followed along the disease paths until they turned 100 years old or died. The model
12 cycle length was 1 year, and half-corrections were applied. A more detailed description for
13 the base-case study population could be found in the **online supplementary file**.

14 ***2.2 Transition probability***

15 The values for the transition probabilities and other inputs were mainly obtained from
16 published evidences among Chinese population and summarized in **Table 1**. A prediction
17 model for hypertension risk was used for estimating hypertension incidence rate [14]. The
18 incidence of each primary CVD events (stroke, HF and CHD) was derived from their
19 corresponding risk equations reported in studies [15-17] on the basis of individual data from
20 Nanjing Community Cardiovascular Risk Survey study [18]. This will keep the correlation
21 between each of the risk factors. The distribution for each risk factors among participants
22 with prehypertension and hypertension could be found in **online supplementary Table A3**
23 and **Table A4**. As the age in this study was mainly from 30 to 75 years old, exponential
24 interpolation was employed to smooth the age-specific annual rates to persons older than 75
25 years (**online supplementary Figure A6 and A7**). A detailed description for the estimation of

1 primary CVD event rate could be found in the **online supplementary file**.
2 Each primary event might be followed by secondary events or chronic conditions, and
3 individual who experienced a first non-fatal cardiovascular event would have a higher risk of
4 sequelaes. As there were no suitable risk equations to calculate the risk for each age/sex
5 stratum among Chinese population, transition probabilities for subsequent events (within or
6 more than 1 year after a primary event) were drawn from registries and/or meta-analysis
7 studies [19-24]. These rates were varied in sensitivity analyses (**online supplementary Table**
8 **A5**). However, for the sake of conservative estimation, the probabilities of new events in any
9 of the post-CVD states were not adjusted their different blood pressure (BP) categories.
10 As evidences in the published literature demonstrated that prehypertension may be associated
11 with higher CVD mortality, we separated mortality into CVD and non-CVD mortality [1, 8].
12 Age- and sex-specific risks of all-cause/CVD mortality were derived from Chinese National
13 Statistical Office in 2013[25]. The non-CVD mortality risks were recalculated after exclusion
14 of CVD death from the total mortality and this was assumed to be unaffected by the states in
15 the model.

16 ***2.3 Treatment effect***

17 Antihypertensive treatment could significantly decrease the risk of CVD morbidity or
18 mortality by lowering BP among patients with hypertension as well as those with clinical
19 history of CVD but without hypertension [26-28]. However, trials to date could only indicate
20 a deceasing risk of incident hypertension, rather than a reduction in CVD risk among
21 prehypertensive individuals without prior CVD [6, 7, 11]. Our model only incorporated the
22 benefits of drug treatment in hypertension incidence, and made the conservative assumption
23 of no effect in CVD risk (**Table 1**).

24 ***2.4 Cost***

25 Costs data were obtained from multiple sources, including the China Health Statistics

1 Yearbook report and health economics researches (**Table 1**). For costs of prehypertension, we
2 assumed it was the same as those with hypertension, although the dose for this specific
3 population may be lower. This approach could be justifiable because it would lead to a
4 conservative estimate. In our model, we used the data from a cost-effectiveness and an
5 observational study [29, 30]. For costs of CVD events during year 1, the values were obtained
6 from several recently published articles which included detailed breakdown costs (eg, drug
7 cost, salary costs of primary healthcare providers) [31-34]. Cost of recurrent stroke or MI
8 were captured by applying the same cost of first-year events. The costs were estimated from a
9 health system perspective and were inflated to the 2014 price level using the average rate of
10 inflation in China from 2009 to 2014 and converted to US dollars (6.25 Yuan \approx 1 USD 2014).
11 A more detailed description for cost estimation could be found in the **online supplementary**
12 **file**.

13 *2.5 Utility*

14 The utility measured in a 2008 survey in China was applied to individuals without CVD [35].
15 We derived utility weight of CHD, stroke or HF from a published cost-effectiveness study
16 [36], as the preference weights for them were not available in Chinese population (**Table 1**).
17 Quality-adjusted life-years (QALYs) were calculated by multiplying the length of time spent
18 in a certain health state by the utility associated with that health state.

19 *2.6 Base-case analysis*

20 We used results from the Nanjing Community Cardiovascular Risk Survey study as a proxy
21 for participants with prehypertension [18]. On average, members of the cohort had a mean
22 age of 50.5 years, were 54.4% female, had a mean systolic blood pressure(SBP), diastolic
23 blood pressure(DBP) and body mass index(BMI) of 131.9mmHg, 82.4 mmHg, 24.5 kg/m²,
24 respectively. Using our base-case cohort, we assessed the simulated 5-year, 10-year and
25 lifetime cumulative incidence of hypertension, CVD events and life expectancy by the

1 method of Markov model cohort simulation. Incremental cost-effectiveness ratios (ICERs)
2 were also calculated by dividing the incremental change in total healthcare costs by the
3 incremental change in QALYs. The degree of cost-effectiveness was evaluated based on the
4 WHO Choosing Interventions That Are Cost Effective (highly cost-effective, ICER less than
5 the gross domestic product(GDP) per capita; moderately cost-effective, ICER of 1-3×GDP
6 per capita; and not cost-effective, ICER of >3×GDP per capita) [37]. The gross domestic
7 product(GDP) per capita of China in 2014 was reported to be \$7,593.9 from the World Bank
8 report [38]. Cost and utility would be both discounted at a 3% annual rate in the future for the
9 primary projections. All analyses were conducted using a life-time horizon unless otherwise
10 indicated.

11 *2.7 Analysis of uncertainty*

12 The impact of uncertainty in model variables on the ICER was tested by a series of sensitivity
13 analyses. First, in the 1-way sensitivity analysis, each input parameter was varied over a set
14 of plausible ranges provided in **Table 1** and **online supplementary Table A5**. Second, a two-
15 way sensitivity analysis was performed to assess how large the changes of the first two key
16 inputs must be in order to exceed the cost-effective threshold (ICER of >3×GDP per
17 capita=\$22,782). Third, in order to assess how sensitive the results were to variations in
18 simultaneous changes of several variables, we also conducted a probabilistic sensitivity
19 analysis (PSA). We ran our model for 5,000 repetitions using a Monte Carlo simulation and at
20 each simulation a value for each of the 54 model parameters were randomly selected based
21 on their respective distributions, which were shown in **Tables 1** and **online supplementary**
22 **Table A5**. In our model, probabilities, prevalence rates, and preference weights were assumed
23 to follow a beta distribution, and cost data were assumed to follow a gamma distribution.
24 Additionally, uniform and triangular distributions were used where appropriate. The
25 incremental cost-effectiveness scatter plot and acceptability curve were constructed to test the

- 1 uncertainty of ICER from the derived 5,000 pairs of incremental cost and QALYs between
- 2 these two strategies.
- 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

3. Results

3.1 Model validation

The findings from our model indicated a reasonable concordance between model predictions and national life tables (30.20 vs. 31.31 for women, 26.69 vs. 27.21 for men). Our results showed that the 5-year absolute cardiovascular risk (except HF) was relatively low (2.58%), however, each event rate was found to be within reported ranges (**online supplementary Tables A6**).

3.2 Base-case analysis

Using baseline estimates, our model simulated the lifetime cumulative incidence of hypertension, which was shown in **online supplementary Tables A8**. The figure indicated that approximately 30% of individuals with placebo would develop hypertension within 5 years, however, it would take more than 17 years to achieve that percent for those prescribed antihypertensive medication. Thus, compared to placebo, the drug intervention delayed the onset of hypertension by about 12 years. Over a lifetime, prescribing drugs among individuals with prehypertension would reduce the risk for developing hypertension by 32.01% (**Table 2**).

Table 2 summarizes the potential changes in CVD events and life expectancy as well as costs, which may be caused by the delaying or reduction of hypertension risk. In general, risk reductions in CHD, stroke and HF were observed for all different time horizons. The reduction in mortality or clinical event risk from drug treatment increased life expectancy by 0.34 years and QALYs by 0.22 (undiscounted: 0.09), whereas it increased discounted costs from \$3,690.1 to \$4,859.5 (an increment of \$1,169.4). Therefore, based on the WHO criteria, the discounted lifetime ICERs might be moderately cost-effective (\$12,994 per QALY within ICER of 1-3×GDP per capita).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

3.3 Uncertainty analysis

Our cost-effectiveness analysis was relatively insensitive to a variety of alternative assumptions (**online supplementary Figure A9**), specifically for those relating to secondary CVD events. The 10 individual parameters that were most influential in ICER estimates were RR of hypertension on treatment, age starting treatment, BMI, drug treatment costs, discount rate, SBP, utility of stroke, incidence of stroke among hypertensive persons, utility of non-CVD state, and DBP.

A two-way sensitivity analysis was performed to investigate the relationship of the two most influential variables (RR of hypertension on treatment and age starting treatment). At a willingness-to-pay threshold of \$22,870 per additional QALY (3 times the GDP), our model indicated that the probability of being cost-effective increased with age and decreased with the higher RR of hypertension on treatment, but there was low probability if the RR of treatment was larger than 0.64 (**online supplementary Figure A10**).

To further reflect the uncertainty in the estimates, the incremental cost-effectiveness scatter plot was derived from PSA (**online supplementary Figure A11**). In comparison with placebo, prescribing drug to participants with prehypertension would result in an increase in QALYs in 99.70% of simulations. Regarding the incremental costs, drug treatment for prehypertension would decrease costs in 5.87% of simulations. Overall, the cost-effectiveness for drug treatment was 30.48% with a maximum willingness to pay to be \$22,870 per additional QALYs. Also, we found an increasing probability of being cost-effective if the decision maker was willing to pay more for an additional health gain, and a 5.80% of the probability that drug intervention was cost-saving (**Figure 2**).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

4. Discussion:

This study provides evidence that drug treatment with the angiotensin-converting enzyme inhibitor ramipril or the angiotensin receptor antagonist candesartan for prehypertension, as compared with placebo intervention, could delay and/or prevent hypertension by approximately 12 years over a lifetime, which would result in a reduction in cardiovascular complications and an increase in life expectancy. In addition, when patients' preferences and cost of treatment were both taken into account, we found that prescribing drugs among prehypertension was moderately cost-effective, costing \$12,994 per QALY.

In comparison with other currently accepted medical interventions for hypertension or CVD, our study found a relative smaller life-year or CVD gains from our base-case model [36, 39-42]. For example, our models projected that an antihypertensive intervention would prolong life expectancy of prehypertensive individuals by 0.34 years. By contrast, Tsevat et al [39] predicted that BP control could increase gains in life expectancy of 2.3 and 1.7 years in men and women, respectively. Geisler et al [36] estimated that renal denervation substantially increased median survival from 17.1 years to 18.4 years and reduced cardiovascular mortality by 30% and all-cause mortality by 15%. We also found a comparable benefit showing metformin interventions could only improve survival by 0.2 years among drug therapy in adults with impaired glucose tolerance [13]. We believed that the relative smaller benefits in our study may be ascribed to the relatively low absolute risk of CVD for this specific population (prehypertension but without CVD) and assumption of no effect on CVD incidence by drug treatment.

In general, the overall conclusions of the cost-effectiveness analysis for our study were robust to the ranges of uncertainty surrounding utility weights, and costs as well as treatment effect. Although it seemed that the judgment of cost-effectiveness was most sensitivity to the RR of antihypertensive treatment, it was still important that health policy should be based on what is

1 determined to be an acceptable ICER. The acceptable ICER from our relative low CVD risk
2 population (2.58% of 5-year CVD risk) may support the concept of using predicted baseline
3 CVD risk equations to guide blood pressure-lowering treatment decisions, not just the BP
4 levels [26, 43, 44]. This is because prehypertensive subjects often exhibited several additional
5 risk factors (e.g., dyslipidemia, obesity) and could have the same absolute CVD risk with
6 those with hypertension [10, 12]. The idea of implementing intensive BP control in high-risk
7 patients was confirmed by SPRINT trial, which may partly support our study results.

8 Several limitations need to be acknowledged in our study. First, due to the lack of evidences
9 to show the benefits on CVD risk by drug treatment among prehypertensive subjects without
10 CVD, our model only captured the benefit of hypertension risk and assumed no direct
11 benefits on stroke, CHD, and HF. This limitation might underestimate the benefit and result
12 in conservative estimates. Second, we did not incorporate the rate of drug side effects and
13 their associated costs in our model, however, it is anticipated to have little influence on the
14 final evaluation as both studies reported a comparable and low rate of serious adverse events
15 between groups and no concerns were raised for adverse reactions caused by drug [6, 7].
16 Third, as there are no reports discussing the medication compliance of a life-long drug
17 treatment among these healthy prehypertensive adults, a potential impact on our result may
18 deserve more attentions. Fourth, many key inputs into the model were derived from Chinese
19 population and we did not know whether our results were transferable to other countries,
20 especially considering the large variations in costs from different countries. Finally, as all
21 computer simulation studies, we were clear that average gains in life expectancy or ICER
22 from the prediction models may not apply to the individual patient, and therefore results need
23 to be interpreted with caution.

24 In summary, our model projected that drug treatment in prehypertensive Chinese adults free
25 of CVD could delay and/or prevent onset of hypertension and its complications. This policy

1 may be cost-effective and reasonable, but are highly dependent on the treatment effectiveness.

2 **Acknowledgments**

3 None

4

1

2 Reference:

- 3 [1] He J, Gu D, Chen J, Wu X, Kelly TN, Huang JF, et al. Premature deaths attributable to blood pressure in China:
4 a prospective cohort study. *Lancet*. 2009;374:1765-72.
- 5 [2] Wang JW, Zhang LX, Wang F, Liu LS, Wang HY. Prevalence, Awareness, Treatment, and Control of
6 Hypertension in China: Results From a National Survey. *American Journal of Hypertension*. 2014;27:1355-61.
- 7 [3] Wu Y, Huxley R, Li L, Anna V, Xie G, Yao C, et al. Prevalence, awareness, treatment, and control of
8 hypertension in China: data from the China National Nutrition and Health Survey 2002. *Circulation*.
9 2008;118:2679-86.
- 10 [4] Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. The Seventh Report of the Joint
11 National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure - The JNC 7
12 Report. *Jama-Journal of the American Medical Association*. 2003;289:2560-72.
- 13 [5] Svetkey LP. Management of prehypertension. *Hypertension*. 2005;45:1056-61.
- 14 [6] Julius S, Nesbitt SD, Egan BM, Weber MA, Michelson EL, Kaciroti N, et al. Feasibility of treating
15 prehypertension with an angiotensin-receptor blocker. *N Engl J Med*. 2006;354:1685-97.
- 16 [7] Luders S, Schrader J, Berger J, Unger T, Zidek W, Bohm M, et al. The PHARAO study: prevention of
17 hypertension with the angiotensin-converting enzyme inhibitor ramipril in patients with high-normal blood
18 pressure: a prospective, randomized, controlled prevention trial of the German Hypertension League. *J*
19 *Hypertens*. 2008;26:1487-96.
- 20 [8] Gu DF, Kelly TN, Wu XG, Chen J, Duan XF, Huang JF, et al. Blood pressure and risk of cardiovascular disease in
21 chinese men and women. *American Journal of Hypertension*. 2008;21:265-72.
- 22 [9] Yu DH, Huang JF, Hu DS, Chen JC, Cao J, Li JX, et al. Prevalence and Risk Factors of Prehypertension Among
23 Chinese Adults. *Journal of Cardiovascular Pharmacology*. 2008;52:363-8.
- 24 [10] Yu D, Huang J, Hu D, Chen J, Cao J, Li J, et al. Association between prehypertension and clustering of
25 cardiovascular disease risk factors among Chinese adults. *J Cardiovasc Pharmacol*. 2009;53:388-400.
- 26 [11] Fuchs FD, de Mello RB, Fuchs SC. Preventing the progression of prehypertension to hypertension: role of
27 antihypertensives. *Curr Hypertens Rep*. 2015;17:505.
- 28 [12] Egan BM, Stevens-Fabry S. Prehypertension--prevalence, health risks, and management strategies. *Nat Rev*
29 *Cardiol*. 2015;12:289-300.
- 30 [13] Herman WH, Hoerger TJ, Brandle M, Hicks K, Sorensen S, Zhang P, et al. The cost-effectiveness of lifestyle
31 modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Annals of*
32 *Internal Medicine*. 2005;142:323-32.
- 33 [14] Chien KL, Hsu HC, Su TC, Chang WT, Sung FC, Chen MF, et al. Prediction models for the risk of new-onset
34 hypertension in ethnic Chinese in Taiwan. *J Hum Hypertens*. 2011;25:294-303.
- 35 [15] Kannell WB, D'Agostino RB, Silbershatz H, Belanger AJ, Wilson PWF, Levy D. Profile for estimating risk of
36 heart failure. *Archives of Internal Medicine*. 1999;159:1197-204.
- 37 [16] Liu J, Hong YL, D'Agostino RB, Wu ZS, Wang W, Sun JY, et al. Predictive value for the Chinese population of
38 the Framingham CHD risk assessment tool compared with the Chinese multi-provincial cohort study. *JAMA*.
39 2004;291:2591-9.
- 40 [17] Chien KL, Su TC, Hsu HC, Chang WT, Chen PC, Sung FC, et al. Constructing the Prediction Model for the Risk
41 of Stroke in a Chinese Population Report From a Cohort Study in Taiwan. *Stroke*. 2010;41:1858-64.
- 42 [18] Yu DH, Chen T, Qin R, Cai YM, Jiang ZX, Zhao ZZ, et al. Association between lung capacity and abnormal
43 glucose metabolism: findings from China and Australia. *Clinical Endocrinology*. 2016;85:37-45.
- 44 [19] Bronnum-Hansen H, Davidsen M, Thorvaldsen P, Grp DMS. Long-term survival and causes of death after
45 stroke. *Stroke*. 2001;32:2131-6.
- 46 [20] Sai XY, He Y, Men K, Wang B, Huang JY, Shi QL, et al. All-cause mortality and risk factors in a cohort of
47 retired military male veterans, Xian, China: an 18-year follow up study. *Bmc Public Health*. 2007;7.
- 48 [21] Wang Z, Li JJ, Wang CX, Yao XM, Zhao XQ, Wang YL, et al. Gender Differences in 1-Year Clinical
49 Characteristics and Outcomes after Stroke: Results from the China National Stroke Registry. *Plos One*. 2013;8.
- 50 [22] Liang Y, Zhu J, Tan HQ, Zhang Y, Liu LS, Chinese Coordinating Center of OR. [Risk factors associated with
51 increased end points of patients with non-ST elevation acute coronary syndromes in China: 2 years follow up
52 results of China-OASIS Registry]. *Zhonghua Xin Xue Guan Bing Za Zhi*. 2009;37:580-4.
- 53 [23] Yu S, Zhao Q, Cui H, Qin M, Liu T, Kong B, et al. Long-term prognostic value of atrial fibrillation in patients
54 with chronic systolic heart failure. *Chin J Cardiac Arrhyth*. 2012;16:39-45.


- 1 [24] Sun I, Lv R, Liang T, Ji S, Kang X, Guo J, et al. Prognosis status with its influencing factors in heart failure
2 patients during 1 year of discharged period Chinese Circulation Journal 2013;25:125-8.
- 3 [25] The Ministry of Health
4 of China.<http://www.nhfpc.gov.cn/htmlfiles/zwggkzt/ptjnj/year2013/index2013.html> 2015, Accessed Jan 6, 2015.
- 5 [26] Sundstrom J, Arima H, Woodward M, Jackson R, Karmali K, Lloyd-Jones D, et al. Blood pressure-lowering
6 treatment based on cardiovascular risk: a meta-analysis of individual patient data. *Lancet*. 2014;384:591-8.
- 7 [27] Thompson AM, Hu TA, Eshelbrenner CL, Reynolds K, He JA, Bazzano LA. Antihypertensive Treatment and
8 Secondary Prevention of Cardiovascular Disease Events Among Persons Without Hypertension A Meta-analysis.
9 *JAMA*. 2011;305:913-22.
- 10 [28] Sipahi I, Swaminathan A, Natesan V, Debanne SM, Simon DI, Fang JC. Effect of Antihypertensive Therapy on
11 Incident Stroke in Cohorts With Prehypertensive Blood Pressure Levels A Meta-Analysis of Randomized
12 Controlled Trials. *Stroke*. 2012;43:432-40.
- 13 [29] Liang X, Gu D, Zhang H, Zhu K, Deng Y, Cao J, et al. The analysis of drug cost and direct medical expense in
14 community health management of hypertensive patients. . *Chin J Prev Med* 2011;45:732-6.
- 15 [30] Wang X, Li WQ, Li X, An N, Chen H, Jan S, et al. Effects and cost-effectiveness of a guideline-oriented
16 primary healthcare hypertension management program in Beijing, China: results from a 1-year controlled trial.
17 *Hypertension Research*. 2013;36:313-21.
- 18 [31] Yuanmei P, Yansheng L, Yan L, Ruifang W, Min S, Xuan H, et al. Study on drug cost burdens of standard
19 secondary prevention for ischemic stroke patients in Shanghai. *Shanghai Med J* 2010;33:808-13.
- 20 [32] Wang Y, Li L, Ma W. Factor analysis of costs of hospitalized patients with stroke who participated in urban
21 and rural medical insurance in Tianjin. *Chinese Journal of Hospital Statistics*. 2014;21:401-4.
- 22 [33] Yuan S, Li N, Zhang Y, Gu Q, Liu Y, Ma J. Effect of different health coverage on hospital expenditure of
23 patients with acute myocardial infarction *Journal of Shanghai Jiaotong University(Medical Science)*. 2013;33:214-
24 9.
- 25 [34] Yu SB, Zhao QY, Cui HY, Qin M, Liu T, Kong B, et al. [Investigation on the prevalence and related factors of
26 medicinal therapy in patients with chronic systolic heart failure]. *Zhonghua Liu Xing Bing Xue Za Zhi*.
27 2012;33:229-33.
- 28 [35] Sun S, Chen JY, Johannesson M, Kind P, Xu L, Zhang YG, et al. Population health status in China: EQ-5D
29 results, by age, sex and socio-economic status, from the National Health Services Survey 2008. *Quality of Life*
30 *Research*. 2011;20:309-20.
- 31 [36] Geisler BP, Egan BM, Cohen JT, Garner AM, Akehurst RL, Esler MD, et al. Cost-Effectiveness and Clinical
32 Effectiveness of Catheter-Based Renal Denervation for Resistant Hypertension. *Journal of the American College*
33 *of Cardiology*. 2012;60:1271-7.
- 34 [37] Baltussen R, Tan TT, Hutubessy R, Acharya A, Evans D, Murray C. Generalized Cost-Effectiveness Analysis: A
35 Guide. World Health Organization. Geneva, Switzerland. 2003.
- 36 [38] The World Bank. <http://data.worldbank.org/indicator/NY.GDP.PCAP.CD>, 2015, Accessed Jan 6, 2015.
- 37 [39] Tsevat J, Weinstein MC, Williams LW, Tosteson ANA, Goldman L. Expected Gains in Life Expectancy from
38 Various Coronary Heart-Disease Risk Factor Modifications. *Circulation*. 1991;83:1194-201.
- 39 [40] Grover SA, Paquet S, Levinton C, Coupal L, Zowall H. Estimating the benefits of modifying risk factors of
40 cardiovascular disease - A comparison of primary vs secondary prevention. *Archives of Internal Medicine*.
41 1998;158:655-62.
- 42 [41] Messori A, Trippoli S, Tendi E. Gains in life expectancy from medical interventions. *New England Journal of*
43 *Medicine*. 1998;339:1943-4.
- 44 [42] Gandjour A, Stock S. A national hypertension treatment program in Germany and its estimated impact on
45 costs, life expectancy, and cost-effectiveness. *Health Policy*. 2007;83:257-67.
- 46 [43] Gaziano TA, Steyn K, Cohen DJ, Weinstein MC, Opie LH. Cost-effectiveness analysis of hypertension
47 guidelines in South Africa - Absolute risk versus blood pressure level. *Circulation*. 2005;112:3569-76.
- 48 [44] Cobiac LJ, Magnus A, Barendregt JJ, Carter R, Vos T. Improving the cost-effectiveness of cardiovascular
49 disease prevention in Australia: a modelling study. *BMC Public Health*. 2012;12:398.

Figure titles and legends:

Figure 1: Schematic Depiction of the Model

— Hypertension module; — prehypertension module; — MI module; — HF module; — Stroke module

Figure 2: Cost-effectiveness acceptability curve






Table 1 key model parameters

Data input	Base Case Numeric Value (Range) or Survival Model (Covariates)	Distribution	Reference
Age	50.5 (30-80)years	Triangular	[18]
Sex	54.4(0-100)% female	Beta	[18]
BMI	24.5(20-30)kg/m ²	Normal	[18]
SBP	131.9(120-139)mmHg	Triangular	[18]
DBP	82.4(65-89)mmHg	Triangular	[18]
Hypertension incidence	Weibull(age, sex, BMI, SBP, DBP)	±25% uniform	[14]
CHD incidence	Exponential (age, sex, DBP, SBP,TC, HDL, DM, Smoking)	±25% uniform	[16]
Stroke incidence	Exponential (age, sex, DBP, SBP, FHS, AF, DM)	±25% uniform	[17]
HF incidence	Exponential (age, sex, SBP, DM, LVH, BMI, heart rate, CHD, valve disease)	±25% uniform	[15]
All-cause mortality	Age and sex dependent	±25% uniform	[25]
Stroke mortality	First year: 12.3% for male, 17.8% for female; beyond: SMR 2.23(1.29–3.88) on background mortality	Beta/ Lognormal	[19-21]
CHD mortality	First year: 5.7%; beyond: SMR 1.43(1.18–1.72) on background mortality	Beta/ Lognormal	[20, 22]
HF mortality	First year: 15.9%; beyond: 8%	Beta	[23, 24]
Effect of treatment			
RR of hypertension on treatment	0.34(0.25-0.95)	Triangular	[6, 7]
Cost			
Annual cost for Antihypertensive treatment	\$161.44(\$53.76-\$484.32)	Gamma	[29, 30]
Annual cost for Stroke	\$3358.11(\$1,118.88-\$10,074.24) first year, post \$1,315.16(\$438.24-\$3,945.44)	Gamma	[25, 31, 32]
Annual cost for CHD	\$4,820.89(\$1,606.88-\$14,462.56) first year, post \$382.08(\$160.64-\$1,446.24)	Gamma	[25, 33]
Annual cost for HF	\$1,260.16(\$420-\$3,780.48) first year, post \$293.92(\$97.92-\$881.76)	Gamma	[25, 34]
Quality of life weights(utilities)			
No cardiovascular disease	Age and sex dependent	±25% uniform	[35]
Stroke	0.63 (0.26 to 0.92)	Beta	[36]
CHD	First year: 0.76 (0.5 to 0.87); beyond: 0.88 (0.67 to 0.94)	Beta	[36]
HF	0.71 (0.43 to 0.84)	Beta	[36]
Discount rate	3% (0%-5%)	Uniform	Assumed

AF: atrial fibrillation; BMI: body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure; DM: diabetes mellitus; TC: total cholesterol; HDL: high-density lipoprotein; FHS: family history stroke; HF: heart failure; LVH: left ventricular hypertrophy; CHD; coronary heart disease; RR: relative risk

Table 2 Base-Case Results

Outcomes	5-year time horizon				10-year time horizon				lifetime horizon			
	Placebo Intervention	Drug Treatment	RD	RR	Placebo Intervention	Drug treatment	RD	RR	Placebo Intervention	Drug treatment	RD	RR
Hypertension	31.37	11.80	19.57	0.38	51.17	20.85	30.32	0.41	80.66	48.65	32.01	0.60
CHD,%	0.80	0.79	-0.01	0.99	1.83	1.73	-0.10	0.95	12.53	12.30	-0.23	0.98
Stroke,%	1.46	1.42	-0.04	0.97	3.49	3.21	-0.28	0.92	26.89	25.41	-1.48	0.94
HF,%	0.95	0.90	-0.05	0.95	2.22	2.00	-0.22	0.90	12.82	12.65	-0.17	0.99
CV mortality,%	0.59	0.57	-0.02	0.97	1.64	1.53	-0.11	0.93	22.96	22.56	-0.40	0.98
All-cause mortality,%	3.08	3.06	-0.02	0.99	7.31	7.19	-0.12	0.98				
Life years									28.46	28.80	0.34	
Undiscounted QALYs									20.43	20.65	0.22	
QALYs discounted,3%									13.46	13.55	0.09	
Cost discounted(\$),3%									3690.1	4859.5	1169.4	
Discounted ICER												\$12,994/QALY

CHD; coronary heart disease; HF: Heart failure; QALYs: quality adjusted life years; ICER: incremental cost-effectiveness ratio; RD: risk difference; RR: relative risk