

Parasites & Vectors

Implications of insecticide resistance for malaria vector control with long-lasting insecticidal nets: trends in pyrethroid resistance during a WHO-coordinated multi-country prospective study --Manuscript Draft--

Manuscript Number:	PARV-D-18-00506R2	
Full Title:	Implications of insecticide resistance for malaria vector control with long-lasting insecticidal nets: trends in pyrethroid resistance during a WHO-coordinated multi-country prospective study	
Article Type:	Research	
Section/Category:	Protozoan Biology and Disease	
Funding Information:	Bill and Melinda Gates Foundation	Not applicable
	Department for International Development	Not applicable
	Medical Research Council	Not applicable
Abstract:	<p>Abstract</p> <p>Background: Increasing pyrethroid resistance has been an undesirable correlate of the rapid increase in coverage of insecticide-treated nets (ITNs) since 2000. Whilst monitoring of resistance levels has increased markedly over this period, longitudinal monitoring is still lacking, meaning the temporal and spatial dynamics of phenotypic resistance in the context of increasing ITN coverage are unclear.</p> <p>Methods: As part of a large WHO-co-ordinated epidemiological study investigating the impact of resistance on malaria infection, longitudinal monitoring of phenotypic resistance to pyrethroids was undertaken in 290 clusters across Benin, Cameroon, India, Kenya and Sudan. Mortality in response to pyrethroids in the major anopheline vectors in each location was recorded during consecutive years using standard WHO test procedures. Trends in mosquito mortality were examined using generalised linear mixed-effect models.</p> <p>Results: Insecticide resistance (using the WHO definition of mortality < 90%) was detected in clusters in all countries across the study period. The highest mosquito mortality (lowest resistance frequency) was consistently reported from India, in an area where ITNs had only recently been introduced. Substantial temporal and spatial variation was evident in mortality measures in all countries. Overall, a trend of decreasing mosquito mortality (increasing resistance frequency) was recorded (Odds Ratio per year: 0.79 per year (95% CI: 0.79–0.81, P < 0.001). There was also evidence that higher net usage was associated with lower mosquito mortality in some countries.</p> <p>Discussion: Pyrethroid resistance increased over the study duration in four out of five countries. Insecticide-based vector control may be compromised as a result of ever higher resistance frequencies.</p>	
Corresponding Author:	Jackie Cook, Phd London School of Hygiene and Tropical Medicine UNITED KINGDOM	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	London School of Hygiene and Tropical Medicine	
Corresponding Author's Secondary Institution:		
First Author:	Jackie Cook (on behalf of consortium)	
First Author Secondary Information:		
Order of Authors:	Jackie Cook (on behalf of consortium)	
Order of Authors Secondary Information:		
Response to Reviewers:	Hi, I have adapted the author details as requested following communication with Aneta	

	Kostadinova- please let me know if anything needs changing with them. We sent better quality figures with the last submission- when I download these, they look fine- but in the submission, they are a bit blurry- could you please let me know if you need improved quality on them. Thanks in advance, Best wishes, Jackie
Additional Information:	
Question	Response
Is this study a clinical trial?<hr><i>A clinical trial is defined by the World Health Organisation as 'any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects on health outcomes'.</i>	No
Are you submitting to an Article Collection?	No

[Click here to view linked References](#)

1 **Implications of insecticide resistance for malaria vector control with long-**
2 **lasting insecticidal nets: trends in pyrethroid resistance during a WHO-**
3 **coordinated multi-country prospective study**
4
5

6
7
8 **Implications of Insecticide Resistance Consortium***
9

10 *Correspondence: Jackie.Cook@lshtm.ac.uk

11
12 Email addresses:

13
14 Jackie Cook: Jackie.Cook@lshtm.ac.uk

15
16 Sean Tomlinson: sean.tomlinson@lstmed.ac.uk

17
18 Immo Kleinschmidt: Immo.Kleinschmidt@lshtm.ac.uk

19
20 Martin James Donnelly: martin.donnelly@lstmed.ac.uk

21
22 Martin Akogbetom: akogbetom@yahoo.fr

23
24 Alioun Adechoubou: aliounadechoubou@hotmail.com

25
26 Achile Massougbodji: massougbodjiachille@yahoo.fr

27
28 Mariam Okê-Sopoh: mariamoke@yahoo.fr

29
30 Vincent Corbel: vincent.corbel@ird.fr

31
32 Sylvie Cornélie: sylvie.cornelie@ird.fr

33
34 Aurore Hounto: aurorefel@yahoo.fr

35
36 Josiane Etang: josyet2@gmail.com

37
38 Herman Parfait Awono-Ambene: hpaawono@yahoo.fr

39
40 Jude Bigoga: jbigoga@gmail.com

41
42 Stanislas Elysée Mandeng: mandengelysee@yahoo.fr

43
44 Boris Njeambosay: njoboris@gmail.com

45
46 Raymond Tabue: ttraymon@yahoo.fr

47
48 Celestin Kouambeng: kouambeng@gmail.com
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Etienne Fondjo: fondjoetienne@yahoo.fr
2
3 Kamaraju Raghavendra: kamarajur2000@yahoo.com
4
5 Rajendra M Bhatt: rmbhatt@rediffmail.com
6
7 Mehul Kumar Chourasia: mehul4frnds@gmail.com
8
9
10 Dipak K. Swain: dipakswain@gmail.com
11
12 [Sreehari Uragayala: sreeuragayala2008@gmail.com](mailto:Sreehari.Uragayala:sreeuragayala2008@gmail.com)
13
14
15 Neena Valecha: neenavalecha@gmail.com
16
17 Charles Mbogo: CMbogo@kemri-welcome.org
18
19
20 Nabie Bayoh: NBayoh@kemricdc.org
21
22 Teresa Kinyari: teresakinyari@yahoo.com
23
24
25 Kiambo Njagi: knjagi@domckenya.or.ke
26
27 Lawrence Muthami: muthamilawrence@yahoo.com
28
29
30 Luna Kamau: LKamau@kemri.org
31
32 Evan Mathenge: mathengeevan@gmail.com
33
34
35 Eric Ochomo: EOchomo@kemricdc.org
36
37 Hmooda Toto Kafy: hmoodatuok@gmail.com
38
39
40 Adam Ismail Bashir: bashiradam69@yahoo.co.uk
41
42 Elfatih M Malik: fatihmmalik@gmail.com
43
44
45 Khalid Elmardi: khalid.elmardi@gmail.com
46
47 Jihad Eltaher Sulieman: jihadmia@hotmail.com
48
49
50 Mujahid Abdin: [Mogahid Abdin@live.com](mailto:Mogahid_Abdin@live.com)
51
52 Krishanthi Subramaniam: Krishanthi.Subramaniam@lstmed.ac.uk
53
54
55 Brent Thomas: brent.thomas@lstmed.ac.uk
56
57
58 Philippa West: pippawest@hotmail.co.uk
59
60 John Bradley: John.Bradley@lshtm.ac.uk
61
62
63
64
65

1 Tessa Bellamy Knox: knoxt@who.int

2 Abraham Peter Mnzava: AMnzava@alma2030.org

3
4 Jonathan Lines: Jo.Lines@lshtm.ac.uk

5
6 Michael Macdonald: macdonaldrm@macito.net

7
8
9 Zinga José Nkuni: nkuni50@yahoo.com

10 11 12 13 14 15 16 17 18 19 20 21 22 23 **Abstract**

24
25 **Background:** Increasing pyrethroid resistance has been an undesirable correlate of the rapid
26 increase in coverage of insecticide-treated nets (ITNs) since 2000. Whilst monitoring of
27 resistance levels has increased markedly over this period, longitudinal monitoring is still
28 lacking, meaning the temporal and spatial dynamics of phenotypic resistance in the context of
29 increasing ITN coverage are unclear.

30
31 **Methods:** As part of a large WHO-co-ordinated epidemiological study investigating the
32 impact of resistance on malaria infection, longitudinal monitoring of phenotypic resistance to
33 pyrethroids was undertaken in 290 clusters across Benin, Cameroon, India, Kenya and Sudan.
34 Mortality in response to pyrethroids in the major anopheline vectors in each location was
35 recorded during consecutive years using standard WHO test procedures. Trends in mosquito
36 mortality were examined using generalised linear mixed-effect models.

37
38 **Results:** Insecticide resistance (using the WHO definition of mortality < 90%) was detected
39 in clusters in all countries across the study period. The highest mosquito mortality (lowest
40 resistance frequency) was consistently reported from India, in an area where ITNs had only
41 recently been introduced. Substantial temporal and spatial variation was evident in mortality
42 measures in all countries. Overall, a trend of decreasing mosquito mortality (increasing
43 resistance frequency) was recorded (Odds Ratio per year: 0.79 per year (95% CI: 0.79–0.81,
44 $P < 0.001$). There was also evidence that higher net usage was associated with lower
45 mosquito mortality in some countries.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **Discussion:** Pyrethroid resistance increased over the study duration in four out of five
2 countries. Insecticide-based vector control may be compromised as a result of ever higher
3 resistance frequencies.
4

5 **Keywords:** Malaria, Vector control, Insecticide resistance, Trends, Bednets, Bioassay
6
7
8
9

10 **Background**

11 Vector control using indoor residual spraying (IRS) and insecticide-treated nets (ITNs) are
12 core strategies for malaria control and elimination. The huge scale-up of these interventions
13 in the last 20 years has been associated with major reductions in disease burden [1]. Between
14 2000 and 2015, it is estimated that over 1 billion ITNs were distributed in malaria endemic
15 countries. The proportion of people in sub-Saharan Africa sleeping under a net increased
16 from 30 to 54% between 2010 and 2016, whilst in 2016 an estimated 2.9% of the at-risk
17 population was covered by IRS globally [1]. The increased coverage of vector control is
18 estimated to have been a major contributor to the documented 62% decline in malaria
19 mortality between 2000–2015 [2, 3]. However, between 2015 and 2016, data suggest that
20 malaria mortality have remained the same in the WHO regions of Southeast Asia, the
21 Western Pacific and Africa, and possibly increased in the Eastern Mediterranean and the
22 Americas [1]. There are therefore justified concerns about the emergence and spread of
23 insecticide resistance and the impact this may have on the continued effectiveness of
24 insecticide-based interventions [1, 4].
25
26
27
28
29
30
31
32
33
34
35
36
37

38 Resistance has now been detected in malaria vectors to the four classes of public
39 health insecticides used in malaria vector control (pyrethroids, organochlorines,
40 organophosphates and carbamates) [5], and up to October 2016 had been reported in 71
41 malaria-endemic countries [6]. Until recently, pyrethroids have been the only class used for
42 long-lasting insecticidal nets (LLINs) and accounted for a large proportion of the insecticide
43 used for IRS. This heavy reliance on a single insecticide class prompted the World Health
44 Organization (WHO) to issue a Global Plan for Insecticide Resistance Management (GPIRM)
45 [5] which was subsequently expanded as part of the Global Vector Control Response [7]. The
46 aim of these initiatives is to sustain the advances made in the fight against vector-borne
47 disease through rational use of vector control tools, including insecticide deployment to slow
48 the development of resistance. Country-level implementation of recommended activities and
49 monitoring has been poor due to a combination of limited availability and costs of
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 insecticides with new modes of action; human, financial and infrastructural capacity
2 shortfalls; and insufficient data to determine epidemiological impact of insecticide resistance
3 [8]. To address this latter point the WHO, with funding from the Bill and Melinda Gates
4 Foundation, initiated a multi-country prospective study to assess the impact of insecticide
5 resistance on the effectiveness of LLINs and IRS. The main objectives of the study were: (i)
6 to determine the impact of insecticide resistance in malaria vectors on the protective
7 effectiveness of LLINs and IRS, and hence on malaria disease burden; and (ii) to assess
8 trends in the insecticide resistance status and underlying mechanisms in the main malaria
9 vector species from the study areas in response to different interventions.
10

11 The study was conducted in five countries, Benin, Cameroon, India, Kenya and
12 Sudan, with data collection conducted from 2010–2016. Details of the overall study design
13 are given in Kleinschmidt et al. [9]. Overall epidemiological outcomes, presented in
14 Kleinschmidt et al. [10], showed that nets provided protection against malaria irrespective of
15 resistance frequency, indicating that populations in malaria endemic areas should continue to
16 use LLINs to reduce their risk of infection. A number of country-specific analyses from this
17 and other studies corroborate this finding [11–15]. In addition, several studies have published
18 country-specific entomological data relating to the second objective [16–18], with ranges of
19 resistance to pyrethroids reported. In this paper, we describe temporal and spatial trends in
20 insecticide resistance of the main malaria vector species from across the five study countries.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Methods**

39 **Study design**

40 The overall study design is described in detail in Kleinschmidt et al. [9]. The five countries
41 included in the study were selected to represent areas of varying transmission intensity where
42 resistance had previously been detected in malaria vectors (Table 1). In 279 study clusters
43 (villages or groups of villages) across 16 areas in the five countries pyrethroid susceptibility
44 in malaria vectors, and malaria infection and disease in children were measured
45 simultaneously over several years. We aimed to assess whether higher levels of resistance are
46 associated with loss of effectiveness of LLINs, and to characterise temporal and spatial trends
47 in insecticide resistance. The numbers of clusters chosen per country are shown in Table 1
48 and were based on sample size calculations determined by the epidemiological outcomes [9].
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Clusters were defined as villages or groups of hamlets with no less than 500 houses and were
2 at least 2 km apart to avoid spill over in outcomes between clusters.
3
4

5 **Vector control**

6
7 LLIN mass distributions were carried out routinely in each site to provide universal coverage
8 for each household (one net per two persons). Nets were distributed in Benin in 2011 (Olyset
9 Net®, Sumitomo Chemical, Tokyo, Japan; 1 g/m² permethrin) and 2014 (PermaNet® 2.0,
10 Vestergaard, Lausanne, Switzerland; 55 mg/m² deltamethrin), in Cameroon in 2011 and 2015
11 (PermaNet® 2.0), in India in 2014 (PermaNet® 2.0), in Kenya in 2010 and 2013
12 (PermaNet® 2.0), and in Sudan in 2011 and 2014 (PermaNet® 2.0). Net usage, defined as the
13 proportion of respondents reporting as having slept under an LLIN the previous night, was
14 determined through cross-sectional surveys which took place at least once in each country
15 during the study period [10]. Cross-sectional household surveys, which consisted of sampling
16 children from random households occurred in 2012 (Kenya, Sudan), 2013 (Cameroon,
17 Sudan), 2014 (Sudan), 2015 (Benin, India) and 2016 (India) [10]. We used net usage as a
18 proxy for the level of local mosquito exposure to pyrethroids. In Sudan half of the clusters
19 were randomised to receive two rounds of IRS with bendiocarb (Ficam®80% WP, Bayer,
20 Leverkusen, Germany; 200 mg active ingredient/m². An exception was the Galabat region
21 where clusters received IRS with deltamethrin (25 mg of a.i./m²; Chema Industries,
22 Alexandria, Egypt) before changing to bendiocarb in subsequent years [15].
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Measuring resistance**

39
40 Phenotypic susceptibility to the pyrethroid deltamethrin, in the main local vector(s), was
41 measured annually in each cluster using WHO adult susceptibility tests and recorded as
42 percent mortality [19]. In Benin, Cameroon, Kenya and Sudan larvae were collected from
43 breeding sites within each cluster annually and reared to adulthood in insectaries. In India,
44 where larval sites were difficult to locate, resting females were caught [19]. Adult female
45 mosquitoes [of unknown age (India); 2–5 days-old (all other countries)] were exposed for 60
46 minutes to deltamethrin using WHO impregnated papers at standard concentrations (0.05%
47 deltamethrin). Mosquitoes were kept at temperatures between 23 and 27 °C, with humidity,
48 where measured, between 75–85%. Mortality was measured 24 h post-exposure. In all tests,
49 observed mortality in control mosquitoes was less than 5% therefore Abbott's correction was
50 not applied.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Statistical analysis

Mosquito mortality data were analysed at the level of the individual mosquito, with post-exposure status (dead/alive after 24 h) modelled as the response variable in logistic regression. Explanatory variables of interest were year, years since last LLIN distribution, and cluster- and year-specific LLIN use as measured in cross-sectional household surveys. Susceptibility test data were excluded from the analysis if fewer than 40 mosquitoes were tested. A mortality estimate was calculated per cluster for each time point, with data for each country analysed separately and in an all-country model. Association between cluster mortality estimates was assessed between years using binomial generalised linear models. Separate generalised mixed-effect models were used to assess trends in mortality over time, effect of time since LLIN mass distribution and effect of bednet use, with the cluster specified as the random effect to account for within cluster correlation of responses. Year was modelled as a linear term to investigate trends over time. Where appropriate, a regional identifier was included as a fixed effect to allow for spatial differences in resistance within countries. Where data were available, insectary temperature and humidity during resistance testing were included in country-level models (Cameroon, India, Sudan).

Cluster-level net usage, as a categorical variable (low, < 40%; medium, 40–80%; and high, > 80%), was explored as an explanatory variable in the years where these data were available from concurrent cross-sectional surveys. As bednet usage was only available for some years, a time variable was not included in these models. To investigate whether the impact of bednet distributions waned over time, models using time since bednet distribution (in years) as the key explanatory variable (as opposed to calendar time) were also examined.

Data from all 5 countries were combined to investigate whether there was evidence for an overall temporal trend in phenotypic resistance, with country added as a fixed effect. As the only data available from 2016 were from India, the all-country analysis was undertaken with and without India.

Results are presented in terms of changes in mortality of mosquitoes by year [Odds Ratios (OR) per year] or with increasing cluster-level category of net usage, with a reduction in mortality indicative of increasing resistance frequency.

Results

Estimates of mortality

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

More than 90,000 mosquitoes were tested in 911 separate tests across 5 countries and over 6 years. The median number of mosquitoes exposed per cluster per year was 100 [interquartile range (IQR) 84–104]. Median mortality across all tests was 81% (IQR: 63–94%). Insecticide resistance, classified according to the WHO criteria of < 90% mortality, was detected in all tested species, in all five countries and in 87% ($n = 793$) of tests performed. In only 7% of tests performed ($n = 63$, from 57 clusters) was 100% mortality observed. There were noticeable differences in the proportions of clusters defined as susceptible across countries. For example, in India, $\geq 98\%$ mortality was observed in 28% ($n = 66$) of tests compared to only 1% of tests ($n = 2$) in Sudan. In Benin, Cameroon, Kenya and Sudan, $> 50\%$ mortality was recorded in at least 14% of tests recorded; no tests in India had less than 50% mortality.

Temporal and spatial variation

Cluster-specific mosquito mortality showed limited and inconsistent evidence of year-to-year correlation in all countries (Fig. 1). The strongest association was seen between data points from 2014 and 2015 (Kendall's tau coefficient: 0.42, $P < 0.001$), although this pattern differed by country, with no correlation seen between those years in Benin or India (Sudan ceased data collection in 2014) (Kendall's tau coefficient 0.07, $P = 0.677$, and 0.02, $P = 0.886$, respectively). The strongest correlation between years was seen in Cameroon, with Kendall's tau coefficient > 0.3 for all year pairs ($P < 0.02$), whilst for the other countries correlation was only present in some pairwise comparisons.

Trends in mortality over time

The trends in mortality over the study period differed by country (Table 2, Fig. 2). A decrease in mortality was detected in Benin, Cameroon, Kenya and Sudan. A slight increase in mortality was detected in India (aOR: 1.03 (95% CI: 0.98–1.1), $P = 0.08$). The most substantial yearly decrease was detected in Sudan (aOR: 0.67 (95% CI: 0.64–0.70), $P < 0.001$). With data from all countries combined, a 21% decrease per year in odds of mortality was detected (aOR 0.79 (95% CI: 0.79–0.81), $P < 0.001$). This was not substantially altered with the exclusion of India (aOR 0.77 (95% CI: 0.76–0.79), $P < 0.001$).

Effect of bednet distributions and bednet use

Bednet distributions occurred in all sites during the study period. Associations between bednet usage and cluster specific mosquito mortality was investigated for each year that epidemiological cross-sectional data were available. Mean net use was above 65% in all

1 countries, with Kenya reporting the highest value (94.2%). Benin, India and Kenya had no
2 clusters with less than 40% net usage. Net usage appeared to have differential impact on
3 mosquito mortality in each country with no association found in Benin and Kenya ($P = 0.225$
4 and $P = 0.241$, respectively); higher mortality found in areas with higher net usage in
5 Cameroon (aOR 1.6 and 1.4 for net usage between 40–80% and above 80% respectively,
6 compared to clusters with net use under 40%, $P < 0.001$) and strong negative associations
7 found in India and Sudan (Table 3).
8
9
10
11

12 Time since bednet distribution was also investigated to establish whether changes
13 associated with bednet distributions waned over time. Differential trends were evident with
14 Benin, India and Sudan demonstrating an increase in odds of mortality (decreasing resistance
15 frequency) for each year post-distribution ($P < 0.001$ for each) whereas mosquito mortality in
16 Cameroon (aOR: 0.95, $P = 0.016$) and Kenya (aOR: 0.59; $P < 0.001$) decreased (increased
17 resistance frequency) with each year post-LLIN distribution (Table 4).
18
19
20
21
22
23
24
25
26

27 Discussion

28 Insecticides have been a key component in the public health and agriculture toolbox for over
29 a century, resulting in the inevitable emergence of resistance in mosquito vectors. This study
30 brings together a very large collection of data from a range of transmission settings to
31 investigate the trends in pyrethroid resistance. Whilst year to year variation was substantial,
32 and poor inter-year correlation prevented cluster specific predictions of resistance, a decrease
33 in mosquito mortality was detected in 4 out of the 5 countries over the 5-year period of the
34 study suggesting that resistance to pyrethroids has been gradually increasing in these settings.
35
36
37
38
39
40
41

42 WHO encourages regular monitoring of resistance frequencies to all insecticides used
43 in country. Consequently, the level of reporting has increased dramatically in recent years
44 with over 30,000 data points now entered into global databases such as the WHO Malaria
45 Threats Map [20] and IR-mapper (www.irmapper.com) [21]. The picture that emerges from
46 these summary data [6, 22, 23], as with the present study, is that resistance to pyrethroids is
47 increasing in frequency and geographic extent. However, these global databases often
48 aggregate data with substantially differential sampling effort across years and regions [6]
49 which may obscure the substantial stochasticity in mortality estimates.
50
51
52
53
54
55
56

57 It is assumed that the increase in resistance to pyrethroids over the past decade is due
58 in part to the higher coverage of insecticide-based interventions, such as LLINs. However,
59
60
61
62
63
64
65

1 studies have shown conflicting results with some reports of increasing resistance following
2 bednet distributions [24–27], and other reports of no increases despite sustained insecticidal
3 campaigns [28–30]. Although ascertaining the effect of bednet coverage was not a primary
4 goal of this study, it was possible to investigate the impact of net use through cross-sectional
5 surveys that were conducted concurrently to resistance measurements. Trends were not
6 uniform across countries, perhaps in part reflecting the differing biology of the vector
7 species. *Anopheles arabiensis* (a major vector in Kenya, Cameroon and Sudan study
8 locations) and *An. culicifacies* (primary vector in India study locations) commonly show high
9 rates of zoophily. Obtaining blood meals from sources other than humans means LLINs
10 would potentially have less impact on selective pressure on resistance. However, overall,
11 higher net usage was associated with increasing resistance in mosquitoes. This trend was
12 most evident in Sudan where the widest range of net usage was reported whereas in other
13 settings reported net usage was more uniform, thereby reducing the likelihood of detecting a
14 trend.
15

16 We did not discern a consistent trend in mosquito mortality with increasing time post-
17 net distribution. In Benin, India and Sudan, mortality increased every year post-distribution,
18 suggesting that the initial increased coverage of nets may have been a short-term driver for
19 resistance and that as the insecticide on the nets reduced over time, the selection pressure
20 reduced, in turn reducing the proportion of resistant mosquitoes. However, in Cameroon and
21 Kenya, the opposite effect was observed, with mortality decreasing with every year from the
22 date of the distribution. Data from the *An. gambiae* 1000 genome project has revealed that
23 there appear to be numerous instances of localised adaptation to insecticide pressure [31].
24 The difference we observed in response to LLIN distribution may reflect in part this innate
25 difference of vector populations to respond to insecticide pressure and caution against making
26 generalised predictions.
27

28 Moreover, whilst bednet distributions will have increased selection pressure in the
29 study settings, it is also possible that the insecticide resistance could be linked to ongoing
30 agricultural practices [32–34]. In several African countries, including northern Cameroon, the
31 use of pyrethroids for cotton farming has been implicated as a catalyst for the increase in
32 recorded resistance in *An. gambiae* populations [34–36]. Differences in the use of pyrethroids
33 for agricultural purposes in the study settings could further impact the relationship between
34 time of net distribution and insecticide resistance.
35

36 Previous studies have also shown resistance to be highly focal [16, 37–39], with large
37 variations over small geographical distances. This is exemplified by the range of mortality
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

measures within each country and highlights the need for multiple sentinel monitoring sites per country and reinforces that extrapolating resistance data from few, widely-dispersed sentinel sites to larger areas is untenable. Spatial heterogeneity in insecticide resistance poses challenges for integrated resistance management and suggests that locally tailored vector control and resistance management programmes are required.

There was considerable temporal heterogeneity with high between year variation at cluster level. This phenomenon has also been reported elsewhere [40–43]. There are several reasons why levels of resistance in a mosquito population may fluctuate over time, for instance, resistance can recede if proper resistance management practices are implemented or if resistance drivers reduce and resistance associated genetic variants are deleterious in the absence of selection pressure [5, 44]. It is possible that in our study settings varying exposure to pyrethroids resulted in fluctuating frequency of resistance in the mosquito population with evidence from some areas suggesting that bednet usage resulted in higher resistance frequencies.

As well as genuine fluctuations in the frequency of resistance, it is possible that the different susceptibility recorded is, in part, an artefact of the method of testing. Longitudinal monitoring is easily influenced by any changes in protocol for measuring mortality and the timings of the tests. Some studies have demonstrated fluctuations in mosquito mortality over a transmission season [45] and whilst all efforts were taken to ensure that tests occurred at the same time each year, differences between seasons may have had an impact. In addition, humidity and temperature are known to have an impact on mortality testing [46]; whilst these were controlled for where data were available, it is possible that differing conditions influenced mortality results.

There is mounting evidence that tests recording mosquito mortality after 24 h may not be the best way to record changes in population resistance, particularly when the level of resistance is high [47]. A number of alternative options are now available for monitoring the presence of resistance, including molecular assays, time/dose response assays and increasing the time post-exposure at which mortality is calculated all of which are likely to be more sensitive to resistance trends [48–51], but these methods are also more resource intensive. Although as noted by Churcher et al. [4] the strong association between bioassay data and mortality measured in experimental hut trials supports the use of bioassays as a quantitative test of the impact of resistance on LLIN efficacy. In this multi-country study, to ensure comparability between sites, the test was performed using one insecticide dose and one exposure time, using wild-caught mosquitoes reared in the laboratory. These settings may not

1 reflect adequately the conditions wild mosquitoes experience, such as variations in
2 temperature, food availability and pre-existing pesticide exposure [51]. In addition, the doses
3 used in the resistance tests are not necessarily reflective of the doses mosquitoes would
4 experience in the wild, which can be influenced by age or retreatment of ITN or regularity
5 and coverage of IRS. The dose used for detecting resistance can have a particularly strong
6 effect depending on the prevalence and penetrance of the resistant mechanisms present in the
7 mosquito population. Recording mortality at 24 h may also miss some of the nuances
8 involved with the evolution of resistance which may result in delayed mortality [51]. In
9 addition, mosquito age has been shown to have a big impact on susceptibility, with older
10 mosquitoes showing higher mortality rates compared to their younger counterparts [52]. If
11 insecticides remain effective against mosquitoes old enough to transmit malaria, this may
12 explain why some studies are observing minimal impact on epidemiological outcomes [10–
13 12, 14).
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **Conclusions**

28 This study demonstrated increasing frequency of resistance to pyrethroids in malaria vectors
29 from 4 out of 5 study countries. Although the increase does not appear linear, if the current
30 trend continues, it is likely to result in a reduction of the effectiveness of pyrethroid-based
31 interventions such as ITN and IRS. There was evidence in some countries of increased
32 selection pressure for pyrethroid resistance in clusters where net use was higher. There are a
33 number of strategies presented within GPIRM to mitigate the increase of insecticide
34 resistance in malaria vectors such as rotations, combinations, mosaics and mixtures [5]. In the
35 short term, two trials have demonstrated improved efficacy of dual-active [53] and
36 pyrethroid-PBO treated LLINs [54], suggesting that we are likely to be able to prolong the
37 useful active life of pyrethroid-based interventions. However, the lack of vector control tools
38 with different modes of actions and their increased costs, means that many endemic countries
39 will continue to struggle to develop and implement insecticide resistance management plans.
40 Whilst new products are currently being trialled [55–57], and some have recently come to
41 market, the fine-scale monitoring of resistance phenotypes and mechanisms will be key to
42 mitigating the impacts of insecticide resistance through informed selection of vector control
43 tools.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Declarations**

3
4 **Ethics approval and consent to participate**

5 Ethics approvals for the study were obtained from the relevant national research ethics
6 committees in the study countries. Ethics approval numbers: 007 (25th May 2010; Benin);
7 102/CNE/SE/09 (Cameroon); ECR/NIMR/EC/2010/75 (India); SSC/ERC No.1677 (Kenya);
8 116-12-09 (Sudan).
9
10
11
12

13
14 **Consent for publication**

15 Not applicable.
16
17
18
19

20 **Availability of data and materials**

21 The datasets used and/or analysed during the present study are available from the
22 corresponding author on reasonable request.
23
24
25
26

27 **Competing interests**

28 The authors declare that they have no competing interests.
29
30
31

32 **Funding**

33 This study was funded by Bill & Melinda Gates Foundation, UK Medical Research Council,
34 and UK Department for International Development.
35
36
37
38
39

40 **Acknowledgements**

41 The Implications of Insecticide Resistance Consortium is a multi-country collaboration. All
42 authors contributed substantially either to the study design, fieldwork, analysis or drafting of
43 the manuscript. All authors read and approved the final manuscript. Study country Principal
44 Investigators are denoted by *
45
46
47
48
49
50
51

52 Writing and analysis team: Jackie Cook¹, Sean Tomlinson², Immo Kleinschmidt^{1,3} and Martin James
53 Donnelly².
54
55

56 Benin country team: *Martin Akogbeto⁴, Alioun Adechoubou⁵, Achile Massougbdji⁶, Mariam Okê-
57 Sopoh⁵, Vincent Corbel⁷, Sylvie Cornélie⁷ and Aurore Hounto⁶.
58
59
60
61
62
63
64
65

1 Cameroon study team: *Josiane Etang^{8,9,10}, Herman Parfait Awono-Ambene⁸, Jude Bigoga¹¹, Stanislas
2 Elysée Mandeng^{8,12}, Boris Njeambosay¹¹, Raymond Tabue¹¹, Celestin Kouambeng¹³ and Etienne
3 Fondjo¹³.
4

5 India study team: *Kamaraju Raghavendra¹⁴, Rajendra M Bhatt¹⁴, Mehul Kumar Chourasia¹⁴, Dipak
6 K. Swain¹⁴, Sreehari Uragayala¹⁴ and Neena Valecha¹⁴.
7

8 Kenya study team: *Charles Mbogo¹⁵, Nabie Bayoh¹⁶, Teresa Kinyari¹⁷, Kiambo Njagi¹⁸, Lawrence
9 Muthami¹⁹, Luna Kamau²⁰, Evan Mathenge²¹ and Eric Ochomo¹⁶.
10

11 Sudan study team: *Hmooda Toto Kafy^{22,23}, Adam Ismail Bashir²⁴, Elfatih M Malik²⁵, Khalid
12 Elmardi²², Jihad Eltaher Sulieman²⁶ and Mujahid Abdin²².
13

14 Study support: Krishanthi Subramaniam², Brent Thomas², Philippa West¹ and John Bradley¹.
15

16 WHO co-ordination team: Tessa Bellamy Knox²⁷, Abraham Peter Mnzava²⁷, Jonathan Lines²⁸,
17 Michael Macdonald²⁷ and Zinga José Nkuni²⁷
18

19 Author details

20 ¹MRC Tropical Epidemiology Group, Department of Infectious Disease Epidemiology,
21 London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK.
22

23 ²Department of Vector Biology, Liverpool School of Tropical Medicine, Pembroke Place,
24 Liverpool, L3 5QA, UK. ³School of Public Health, University of the Witwatersrand,
25 Johannesburg, South Africa. ⁴Centre de Recherche Entomologique de Cotonou. Cotonou,
26 Benin. ⁵Programme National de Lutte contre le Paludisme (PNLP), Ministère de la Santé,
27 Benin. ⁶Faculté des Sciences de la Santé, Université d'Abomey Calavi, Cotonou, Benin.
28

29 ⁷Maladies Infectieuses et Vecteurs, Ecologie, Génétique, Evolution et Contrôle (MIVEGEC),
30 Institut de Recherche pour le Développement (IRD), CNRS, University of Montpellier,
31 Montpellier, France. ⁸Organisation de Coordination pour la lutte contre les Endemies en
32 Afrique Centrale (OCEAC), Yaounde, Cameroon. ⁹Faculty of Medicine and Pharmaceutical
33 Sciences, University of Douala, PO Box 2701, Douala, Cameroon. ¹⁰ Institute for Insect
34 Biotechnology, Justus Liebig University Gießen, Winchesterstr. 2, 35394 Gießen, Germany.
35

36 ¹¹National Reference Unit (NRU) for Vector Control, The Biotechnology Center, University
37 of Yaounde I, PO Box 3851, Messa, Yaounde, Cameroon. ¹²Laboratory of General Biology,
38 University of Yaounde I, P.O. Box 812, Yaounde, Cameroon. ¹³National Malaria Control
39 Program, Ministry of Public Health, PO, Box 14386, Yaounde, Cameroon. ¹⁴National
40 Institute of Malaria Research, Department of Health Research, (GoI), Sector 8, Dwarka, New
41 Delhi 110 077, India. ¹⁵KEMRI Centre for Geographic Medicine Research Coast, PO Box
42

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

230 – 80108, Kilifi, Kenya. ¹⁶KEMRI/CDC Research and Public Health Collaboration, PO Box 1578, Kisumu 40100, Kenya. ¹⁷University of Nairobi, School of Medicine, College of Health Sciences, Department of Medical Physiology, Nairobi, Kenya. ¹⁸Ministry of Health, Malaria Control Unit, PO Box 1992–00202, Nairobi, Kenya. ¹⁹KEMRI Centre for Public Health Research, Nairobi, Kenya. ²⁰KEMRI - Centre for Biotechnology and Research Development, Nairobi, Kenya. ²¹KEMRI - Eastern and Southern Africa Centre of International Parasite Control, Nairobi, Kenya. ²²Integrated Vector Management Department, Federal Ministry of Health, PO Box 303, Khartoum, Sudan. ²³School of Biological Sciences, Universiti Sains Malaysia, 11800 USM Pulau Penang, Malaysia. ²⁴Khartoum State Malaria Control Programme, VBDC, PO Box 1517, Khartoum, Sudan. ²⁵Faculty of Medicine, University of Khartoum, Khartoum, Sudan. ²⁶Sennar Malaria Research and Training Centre, Sennar, Sudan. ²⁷Global Malaria Programme, World Health Organization, Avenue Appia 20, 1211 Geneva, Switzerland. ²⁸Department of Disease Control, London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK.

References

1. World Malaria Report 2017. World Health Organisation, 2017.
2. World Malaria Report 2016. Geneva: World Health Organisation, 2016.
3. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature*. 2015;526:207–11.
4. Churcher TS, Lissenden N, Griffin JT, Worrall E, Ranson H. The impact of pyrethroid resistance on the efficacy and effectiveness of bednets for malaria control in Africa. *Elife*. 2016;5:pii:e16090.
5. Global plan for insecticide resistance management in malaria vectors. Geneva: World Health Organisation; 2012.
6. Coleman M, Hemingway J, Gleave KA, Wiebe A, Gething PW, Moyes CL. Developing global maps of insecticide resistance risk to improve vector control. *Malar J*. 2017;16:86.
7. Global Vector Control Response: 2017–2030. Geneva: World Health Organization; 2017.
8. Mnzava AP, Knox TB, Temu EA, Trett A, Fornadel C, Hemingway J, et al. Implementation of the global plan for insecticide resistance management in malaria vectors: progress, challenges and the way forward. *Malar J*. 2015;14:173.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
9. Kleinschmidt I, Mnzava AP, Kafy HT, Mbogo C, Bashir AI, Bigoga J, et al. Design of a study to determine the impact of insecticide resistance on malaria vector control: a multi-country investigation. *Malar J.* 2015;14:282.
 10. Kleinschmidt I, Bradley J, Knox TB, Mnzava AP, Kafy HT, Mbogo C, et al. Implications of insecticide resistance for malaria vector control with long-lasting insecticidal nets: a WHO-coordinated, prospective, international, observational cohort study. *Lancet Infect Dis.* 2018;18:640–649.
 11. Ochomo E, Chahilu M, Cook J, Kinyari T, Bayoh NM, West P, et al. Insecticide-treated nets and protection against insecticide-resistant malaria vectors in Western Kenya. *Emerg Infect Dis.* 2017;23:758–64.
 12. Bradley J, Ogouyemi-Hounto A, Cornelie S, Fassinou J, de Tove YSS, Adeothy AA, et al. Insecticide-treated nets provide protection against malaria to children in an area of insecticide resistance in southern Benin. *Malar J.* 2017;16:225.
 13. Chourasia MK, Kamaraju R, Kleinschmidt I, Bhatt RM, Swain DK, Knox TB, et al. Impact of long-lasting insecticidal nets on prevalence of subclinical malaria among children in the presence of pyrethroid resistance in *Anopheles culicifacies* in central India. *Int J Infect Dis.* 2017;57:123–9.
 14. Lindblade KA, Mwandama D, Mzilahowa T, Steinhardt L, Gimnig J, Shah M, et al. A cohort study of the effectiveness of insecticide-treated bed nets to prevent malaria in an area of moderate pyrethroid resistance, Malawi. *Malar J.* 2015;14:31.
 15. Kafy HT, Ismail BA, Mnzava AP, Lines J, Abdin MSE, Eltahir JS, et al. Impact of insecticide resistance in *Anopheles arabiensis* on malaria incidence and prevalence in Sudan and the costs of mitigation. *Proc Natl Acad Sci USA.* 2017;114:E11267–75.
 16. Ochomo E, Bayoh NM, Kamau L, Atieli F, Vulule J, Ouma C, et al. Pyrethroid susceptibility of malaria vectors in four districts of western Kenya. *Parasit Vectors.* 2014;7:310.
 17. Tabue RN, Awono-Ambene P, Etang J, Atangana J, C AN, Toto JC, et al. Role of *Anopheles (Cellia) rufipes* (Gough, 1910) and other local anophelines in human malaria transmission in the northern savannah of Cameroon: a cross-sectional survey. *Parasit Vectors.* 2017;10:22.
 18. Ismail BA, Kafy HT, Sulieman JE, Subramaniam K, Thomas B, Mnzava A, et al. Temporal and spatial trends in insecticide resistance in *Anopheles arabiensis* in Sudan: outcomes from an evaluation of implications of insecticide resistance for malaria vector control. *Parasit Vectors.* 2018;11:122.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
19. Global Malaria Programme. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes, 2 ed. Geneva: World Health Organization; 2016.
 20. Malaria Threats Map: World Health Organisation. www.who.int/malaria/maps/threats/. Accessed 1 Feb 2018
 21. Knox TB, Juma EO, Ochomo EO, Pates Jamet H, Ndungo L, Chege P, et al. An online tool for mapping insecticide resistance in major *Anopheles* vectors of human malaria parasites and review of resistance status for the Afrotropical region. *Parasit Vectors*. 2014;7:76.
 22. Ranson H, Lissenden N. Insecticide resistance in African *Anopheles* mosquitoes: a worsening situation that needs urgent action to maintain malaria control. *Trends Parasitol*. 2016;32:187–96.
 23. Dialynas E, Topalis P, Vontas J, Louis C. MIRO and IRbase: IT tools for the epidemiological monitoring of insecticide resistance in mosquito disease vectors. *PLoS Negl Trop Dis*. 2009;3:e465.
 24. Stump AD, Atieli FK, Vulule JM, Besansky NJ. Dynamics of the pyrethroid knockdown resistance allele in western Kenyan populations of *Anopheles gambiae* in response to insecticide-treated bed net trials. *Am J Trop Med Hyg*. 2004;70:591–6.
 25. Reimer LJ, Tripet F, Slotman M, Spielman A, Fondjo E, Lanzaro GC. An unusual distribution of the kdr gene among populations of *Anopheles gambiae* on the island of Bioko, Equatorial Guinea. *Insect Mol Biol*. 2005;14:683–8.
 26. Czeher C, Labbo R, Arzika I, Duchemin JB. Evidence of increasing Leu-Phe knockdown resistance mutation in *Anopheles gambiae* from Niger following a nationwide long-lasting insecticide-treated nets implementation. *Malar J*. 2008;7:189.
 27. Protopopoff N, Verhaeghen K, Van Bortel W, Roelants P, Marcotty T, Baza D, et al. A significant increase in kdr in *Anopheles gambiae* is associated with an intensive vector control intervention in Burundi highlands. *Trop Med Int Health*. 2008;13:1479–87.
 28. Kulkarni MA, Malima R, Mosha FW, Msangi S, Mrema E, Kabula B, et al. Efficacy of pyrethroid-treated nets against malaria vectors and nuisance-biting mosquitoes in Tanzania in areas with long-term insecticide-treated net use. *Trop Med Int Health*. 2007;12:1061–73.
 29. Vulule JM, Beach RF, Atieli FK, Mount DL, Roberts JM, Mwangi RW. Long-term use of permethrin-impregnated nets does not increase *Anopheles gambiae* permethrin tolerance. *Med Vet Entomol*. 1996;10:71–9.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
30. Koimbu G, Czeher C, Katusele M, Sakur M, Kilepak L, Tandrapah A, et al. Status of insecticide resistance in Papua New Guinea: an update from nation-wide monitoring of *Anopheles* mosquitoes. *Am J Trop Med Hyg.* 2018;98:162–5.
 31. *Anopheles gambiae* 1000 Genomes Consortium. Genetic diversity of the African malaria vector *Anopheles gambiae*. *Nature.* 2017;552:96–100.
 32. Hemingway J, Jayawardena KG, Herath PR. Pesticide resistance mechanisms produced by field selection pressures on *Anopheles nigerrimus* and *A. culicifacies* in Sri Lanka. *Bull World Health Organ.* 1986;64:753–8.
 33. Abuelmaali SA, Elaagip AH, Basheer MA, Frah EA, Ahmed FT, Elhaj HF, et al. Impacts of agricultural practices on insecticide resistance in the malaria vector *Anopheles arabiensis* in Khartoum State, Sudan. *PLoS One.* 2013;8:e80549.
 34. Chouaibou M, Etang J, Brevault T, Nwane P, Hinzoumbe CK, Mimpfoundi R, et al. Dynamics of insecticide resistance in the malaria vector *Anopheles gambiae* (*s.l.*) from an area of extensive cotton cultivation in northern Cameroon. *Trop Med Int Health.* 2008;13:476–86.
 35. Diabate A, Baldet T, Chandre F, Akoobeto M, Guiguemde TR, Darriet F, et al. The role of agricultural use of insecticides in resistance to pyrethroids in *Anopheles gambiae* (*s.l.*) in Burkina Faso. *Am J Trop Med Hyg.* 2002;67:617–22.
 36. N'Guessan R, Corbel V, Akogbeto M, Rowland M. Reduced efficacy of insecticide-treated nets and indoor residual spraying for malaria control in pyrethroid resistance area, Benin. *Emerg Infect Dis.* 2007;13:199–206.
 37. Kabula B, Tungu P, Matowo J, Kitau J, Mweya C, Emidi B, et al. Susceptibility status of malaria vectors to insecticides commonly used for malaria control in Tanzania. *Trop Med Int Health.* 2012;17:742–50.
 38. Cisse MB, Keita C, Dicko A, Dengela D, Coleman J, Lucas B, et al. Characterizing the insecticide resistance of *Anopheles gambiae* in Mali. *Malar J.* 2015;14:327.
 39. Matowo NS, Munhenga G, Tanner M, Coetzee M, Feringa WF, Ngowo HS, et al. Fine-scale spatial and temporal heterogeneities in insecticide resistance profiles of the malaria vector, *Anopheles arabiensis* in rural south-eastern Tanzania. *Wellcome Open Res.* 2017;2:96.
 40. Foster GM, Coleman M, Thomsen E, Ranson H, Yangalbe-Kalnone E, Moundai T, et al. Spatial and temporal trends in insecticide resistance among malaria vectors in Chad highlight the importance of continual monitoring. *PLoS One.* 2016;11:e0155746.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
41. Badolo A, Traore A, Jones CM, Sanou A, Flood L, Guelbeogo WM, et al. Three years of insecticide resistance monitoring in *Anopheles gambiae* in Burkina Faso: resistance on the rise? *Malar J.* 2012;11:232.
 42. Djegbe I, Boussari O, Sidick A, Martin T, Ranson H, Chandre F, et al. Dynamics of insecticide resistance in malaria vectors in Benin: first evidence of the presence of L1014S kdr mutation in *Anopheles gambiae* from West Africa. *Malar J.* 2011;10:261.
 43. Yahouedo GA, Cornelie S, Djegbe I, Ahlonsou J, Aboubakar S, Soares C, et al. Dynamics of pyrethroid resistance in malaria vectors in southern Benin following a large scale implementation of vector control interventions. *Parasit Vectors.* 2016;9:385.
 44. Raghavendra K, Verma V, Srivastava HC, Gunasekaran K, Sreehari U, Dash AP. Persistence of DDT, malathion & deltamethrin resistance in *Anopheles culicifacies* after their sequential withdrawal from indoor residual spraying in Surat district, India. *Indian J Med Res.* 2010;132:260–4.
 45. Mbepera S, Nkwengulila G, Peter R, Mause EA, Mahande AM, Coetzee M, et al. The influence of age on insecticide susceptibility of *Anopheles arabiensis* during dry and rainy seasons in rice irrigation schemes of northern Tanzania. *Malar J.* 2017;16:364.
 46. Glunt KD, Paaijmans KP, Read AF, Thomas MB. Environmental temperatures significantly change the impact of insecticides measured using WHOPES protocols. *Malar J.* 2014;13:350.
 47. Bagi J, Grisales N, Corkill R, Morgan JC, N'Fale S, Brogdon WG, et al. When a discriminating dose assay is not enough: measuring the intensity of insecticide resistance in malaria vectors. *Malar J.* 2015;14:210.
 48. Muller P, Chouaibou M, Pignatelli P, Etang J, Walker ED, Donnelly MJ, et al. Pyrethroid tolerance is associated with elevated expression of antioxidants and agricultural practice in *Anopheles arabiensis* sampled from an area of cotton fields in Northern Cameroon. *Mol Ecol.* 2008;17:1145–55.
 49. Mawejje HD, Wilding CS, Rippon EJ, Hughes A, Weetman D, Donnelly MJ. Insecticide resistance monitoring of field-collected *Anopheles gambiae* (*s.l.*) populations from Jinja, eastern Uganda, identifies high levels of pyrethroid resistance. *Med Vet Entomol.* 2013;27:276–83.
 50. Toe KH, Jones CM, N'Fale S, Ismail HM, Dabire RK, Ranson H. Increased pyrethroid resistance in malaria vectors and decreased bed net effectiveness, Burkina Faso. *Emerg Infect Dis.* 2014;20:1691–6.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
51. Viana M, Hughes A, Matthiopoulos J, Ranson H, Ferguson HM. Delayed mortality effects cut the malaria transmission potential of insecticide-resistant mosquitoes. *Proc Natl Acad Sci USA*. 2016;113:8975–80.
52. Jones CM, Sanou A, Guelbeogo WM, Sagnon N, Johnson PC, Ranson H. Aging partially restores the efficacy of malaria vector control in insecticide-resistant populations of *Anopheles gambiae s.l.* from Burkina Faso. *Malar J*. 2012;11:24.
53. Tiono AB, Ouedraogo A, Ouattara D, Bougouma EC, Coulibaly S, Diarra A, et al. Efficacy of Olyset Duo, a bednet containing pyriproxyfen and permethrin, *versus* a permethrin-only net against clinical malaria in an area with highly pyrethroid-resistant vectors in rural Burkina Faso: a cluster-randomised controlled trial. *Lancet*. 2018;pii:S0140–6736(18)31711–2.
54. Protopopoff N, Mosha JF, Lukole E, Charlwood JD, Wright A, Mwalimu CD, et al. Effectiveness of a long-lasting piperonyl butoxide-treated insecticidal net and indoor residual spray interventions, separately and together, against malaria transmitted by pyrethroid-resistant mosquitoes: a cluster, randomised controlled, two-by-two factorial design trial. *Lancet*. 2018;391:1577–88.
55. N'Guessan R, Odjo A, Ngufor C, Malone D, Rowland M. A Chlorfenapyr Mixture Net Interceptor(R) G2 shows high efficacy and wash durability against resistant mosquitoes in West Africa. *PLoS One*. 2016;11:e0165925.
56. Sternberg ED, Ng'habi KR, Lyimo IN, Kessy ST, Farenhorst M, Thomas MB, et al. Eave tubes for malaria control in Africa: initial development and semi-field evaluations in Tanzania. *Malar J*. 2016;15:447.
57. Qualls WA, Muller GC, Traore SF, Traore MM, Arheart KL, Doumbia S, et al. Indoor use of attractive toxic sugar bait (ATSB) to effectively control malaria vectors in Mali, West Africa. *Malar J*. 2015;14:301.

49
50

Figure legends

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 1 Association in cluster mortality between years. Scatter diagrams show results for clusters with mortality estimates in consecutive years for each year of the study. The predicted mortality result from binomial generalised linear models is overlaid on each graph with 95% confidence intervals

Fig. 2 Box-and-whisker plots showing the range of cluster-level mortality by year and country. Arrows indicate the timing of bednet distributions within country

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Details of study sampling and sites including vector control coverage and insecticide resistance prevalence at baseline

	Study sampling sites				
	Benin	Cameroon	India	Kenya	Sudan
Malaria transmission intensity	High	High	Low	High	Low
Study locations	Districts of Ifangni, Sakété, Pobé and Kétou (Departement de Plateau)	Districts of Garoua, Pitoa and Mayo Oulo (North region)	Subdistrict of Keshkal (Kondagaon, Chhattisgarh)	Districts of Teso, Rachuonyo, Nyando and Bondo (western Kenya)	El Hoosh and Hag Abdalla (Gezira State); Galabat (Gedarif State); New Halfa (Kassala State)
Number of clusters sampled	32	38	80	61	79
Entomological sampling points (years)	2011–2015	2012–2015	2013–2016	2011–2015	2011–2014
Main malaria vectors	<i>Anopheles gambiae</i> (s.s.) ^a , <i>Anopheles coluzzii</i> ^a	<i>An. arabiensis</i> ^a , <i>An. gambiae</i> (s.s.) ^a , <i>An. funestus</i>	<i>An. culicifacies</i> ^a	<i>An. gambiae</i> (s.s.) ^a , <i>An. arabiensis</i> ^a , <i>An. funestus</i>	<i>An. arabiensis</i> ^a
Vector control interventions	High coverage of ITNs (primarily PermaNet 2.0) in all clusters	High coverage of ITNs (PermaNet 2.0) in all clusters	High coverage of ITNs (PermaNet 2.0) in all clusters	High coverage of ITNs (PermaNet 2.0 and Olyset Net) in all clusters. Rachuonyo and Nyando received IRS with deltamethrin and lambda-cyhalothrin in 2012, but no IRS was carried out subsequently	High coverage of ITNs (PermaNet 2.0) in all study clusters. In each study area half of clusters randomly allocated to receive additional IRS with bendiocarb
Baseline insecticide resistance	Kdr frequency by cluster ranged from 44 to 93%	Kdr frequency by cluster ranged from 9	WHO Bioassay mortality to deltamethrin	WHO Bioassay mortality to deltamethrin	Kdr frequency by cluster ranged from 8.3

information	(2011) WHO	to 65% (2011)	ranged between	ranged between	to 70.8%
(cluster-specific	Bioassay	WHO Bioassay	86–100%	1–100% (2011)	(2010); WHO
range)	mortality to	mortality to			Bioassay
	deltamethrin	deltamethrin			mortality to
	ranged between	ranged between			deltamethrin in
	20–100% (2011)	43–100%			sentinel clusters
		(2012)			ranged between
					47–100%
					(2011)

^aMortality results in paper presented for these species

Table 2 Impact of time on mosquito mortality. Results from generalised linear mixed-effect models examining the impact on mosquito mortality over time (year)

Country	Odds ratio for change in mortality per year (95% CI)	P-value
All five countries combined ^a	0.79 (0.79–0.81)	<0.001
Four countries combined (without India) ^a	0.77 (0.76–0.79)	<0.001
Benin ^b	0.74 (0.72–0.76)	<0.001
Cameroon ^c	0.74 (0.69–0.78)	<0.001
India ^c	1.03 (0.98–1.1)	0.08
Kenya ^b	0.88 (0.86–0.90)	<0.001
Sudan ^c	0.67 (0.64–0.70)	<0.001

^aAdjusted for country

^bAdjusted for district

^cAdjusted for district, temperature and humidity

Results are presented in terms of change in odds of mortality of mosquitoes in WHO bioassays by year. Odds ratios are adjusted for locality and temperature and humidity where indicated. The data are shown for each country, as well as all countries combined (with country included as a covariate). Cluster was included as a random effect in all models

Table 3 Impact of cluster-level bednet usage on mosquito mortality. Results from generalised mixed-effect models examining the impact of cluster-level bednet usage on mosquito mortality

		All countries combined ^a	Benin ^b	Cameroon ^c	India ^c	Kenya
	No. of clusters included (year)	59 (2012); 87 (2013); 143 (2014); 99 (2015); 80 (2016)	19 (2015)	22 (2013); 26 (2014)	80 (2015); 80 (2016)	13(201
	Mean net usage (range) (%)		74.9 (52.5–100)	67.8 (7.0–100)	89.9 (60.9–100)	94.2 (7
Effect of cluster-level net usage on mosquito mortality, OR (95% CI)	< 40%	1 (reference)	–	1(reference)	–	–
	40–80%	1.03 (0.89–1.19)	1 (reference)	1.61 (1.21–2.14)	1 (reference)	1 (refer
	> 80%	0.65 (0.57–0.74)	1.59 (0.75–3.37)	1.40 (1.08–1.82)	0.36 (0.29–0.44)	2.38 (0
	<i>P</i> -value	<0.001	0.225	<0.001	<0.001	0.241

^aAdjusted for country

^bAdjusted for district

^cAdjusted for district, temperature and humidity

Results are presented in terms of change in mortality of mosquitoes for increasing bednet usage category (< 40%; between 40–80%; and above 80%). Bednet usage was calculated for years where cross-sectional survey data was available. Odds ratios are adjusted for locality and temperature and humidity where indicated. The results are shown for each country, as well as all countries combined (with country included as a covariate). Cluster was included as a random effect in all models

Table 4 Impact of time since bednet distribution (years) on mosquito mortality. Results from generalised mixed-effect models examining the impact time since bednet distribution on mosquito mortality

Country	Odds ratio for change in mortality per year (95% CI)	<i>P</i> -value
All countries combined ^a	1.34 (1.31–1.37)	<0.001
Benin ^b	3.20 (3.02–3.39)	<0.001
Cameroon ^c	0.95 (0.90–0.99)	0.016
India ^c	1.62 (1.52–1.73)	<0.001
Kenya ^b	0.59 (0.56–0.62)	<0.001
Sudan ^c	1.60 (1.53–1.67)	<0.001

^aAdjusted for country

^bAdjusted for district

^cAdjusted for district, temperature and humidity

Results are presented in terms of change in mortality of mosquitoes for each year since a mass bednet distribution took place in-country. Odds ratios are adjusted for locality and temperature and humidity where indicated. The results are shown for each country, as well as all countries combined (with country included as a covariate). Cluster was included as a random effect in all models



